

Exact Solutions for Some Singular Linear Ordinary Differential Equations of High Orders via NB1 Polynomials

Abstract

In this study, we numerically solve the singular linear ordinary differential equations (SLODEs) of higher order using the collocation method based on the NB1 polynomial. The form of an operational matrix of the given ordinary differential equations (ODEs), the relations of various solutions and the derivatives are obtained from NB1 polynomials. The linear algebraic equations system appears from transforming the given problem by the proposed method, thus disappearing the singularity of the problem. The inverse matrix method is used to solve the resulting system to obtain the coefficients of NB1 polynomials. The obtained exact solutions to different problems of high orders show the reliability and accuracy of the presented method.

Keywords: Collocation method, NB1 polynomial, operational matrix form, singular differential equations.

1. Introduction

The equations Emden-Fowler with appropriate initial and boundary value problems appear in various fields of engineering and science, for instance, fluid mechanics and quantum, chemical reactors, optimal design, geophysics and so on, see [6, 9, 19, 31, 39, 43] for more details.

In this study, we will consider the singular linear Emden-Fowler of order $(m + 1)$, $m \geq 1$ in the following general form:

$$\omega^{(m+1)}(r) + \frac{\mu_1}{r} \omega^{(m)}(r) + \frac{\mu_2}{r^2} \omega^{(m-1)}(r) + g(r)\omega(r) = h(r), \quad (1)$$

with initial conditions (ICs.)

$$\omega(0) = \omega'(0) = \omega''(0) = \dots = \omega^{(m)}(0) = 0, \quad (2)$$

where μ_1 and μ_2 are appropriate constants, g and h are given real values functions of the variable r and $\omega(r)$ is an unknown function.

We can derive some various order kinds of Emden-Fowler Eq. (1) from the following equation:

$$r^{-t} \frac{d}{dr} \left[r^{t-1} \frac{d^s}{dr^s} (r \omega) \right] + g(r)\omega - h(r) = 0, \quad t, s \geq 1. \quad (3)$$

For $s = 1, 2, 3, \dots, m$, we get the first kind, the second kind, and the third kind up to $(m + 1)th$ kind respectively of the Emden-Fowler equation, as follows:

$$\omega^{(2)} + \frac{t+1}{r} \omega^{(1)} + \frac{t-1}{r^2} \omega + g(r)\omega - h(r) = 0, \quad (4)$$

$$\omega^{(3)} + \frac{t+2}{r} \omega^{(2)} + \frac{2(t-1)}{r^2} \omega^{(1)} + g(r)\omega - h(r) = 0, \quad (5)$$

$$\omega^{(4)} + \frac{t+3}{r} \omega^{(3)} + \frac{3(t-1)}{r^2} \omega^{(2)} + g(r)\omega - h(r) = 0, \quad (6)$$

$$\vdots$$

$$\omega^{(m+1)} + \frac{t+m}{r} \omega^{(m)} + \frac{m(t-1)}{r^2} \omega^{(m-1)} + g(r)\omega - h(r) = 0, \quad (7)$$

Where $\mu_1 = t + s$ and $\mu_2 = s(t - 1)$, $s = 1, 2, \dots, m$.

If $m = 1$ and $\mu_2 = 0$, Eq. (1) becomes

$$\omega'' + \frac{\mu_1}{r} \omega' + g(r)\omega - h(r) = 0, \quad (8)$$

which called the Lane- Emden equation. The standard linear Lane- Emden equation is obtained from Eq. (8) When $\mu_1 = 2$, $g(r) = 1$ and $h(r) = 0$. Many models of Eq. (8) with some appropriate fixed values of

μ_1 as astrophysics phenomena and a mathematical physics you can find in [10, 13, 17, 20, 24, 25, 26, 28, 40, 41, 45]. The singularity at $r = 0$ makes a solution of these types of equations is difficult.

The Emden-Fowler and Lane-Emden equations are solved analytically and numerically by various methods. [15, 16, 27, 39, 42] used the Adomian decomposition method, [3, 7, 32] employed the homotopy analysis method, [36, 44, 46] used the variational iteration method. You can find more different methods in [5, 11, 29, 30, 33, 35, 37, 47]. In the 20th century, the Taylor wavelet method solved the linear and nonlinear Lane-Emden Equations [14]. In [18], Kashem solved the nonlinear Lane-Emden problem by modifying Hermite operational matrix method. Haar wavelet quasilinearization method was used to solve the Emden-Fowler type equations [34]. In [22], Said-Ball Polynomials were employed to solve some linear delay differential equations. The DP-Ball and Wang-Ball polynomials [21, 23] were used to solve singular ordinary differential equations. In [12], Bessel matrix method was used to solve linear and nonlinear singular differential equations. In [38], Wang et al. used the Chebyshev collocation method to solve singular ordinary differential equations. The least square method depends on Wang Ball function to find an approximate solution of ODEs for higher order [4]. Lastly, Said-Ball Polynomials were used to solve singular ordinary differential equations in [2].

The main purpose of this research is to use the method of an operational matrix depending on NB1 Polynomials to solve high order of singular linear ordinary differential equations (SLODEs).

We organize this paper as follows: Some concepts of NB1 polynomials and NB1 monomial formulas are presented in section 2. Section 3 shows the fundamental matrix relations. Numerical examples are presented in section 4. Section 5 is a conclusion of this study.

2. The NB1curve and NB1 monomial formulas:

2.1 Definition: An NB1 polynomial $N_i^n(r)$ of degree n ($n > 3$) is defined as [8]:

$$N_i^n(r) = \begin{cases} \binom{i + \lfloor \frac{n}{2} \rfloor - 1}{i} r^i (-r + 1)^{\lfloor \frac{n}{2} \rfloor}, & 0 \leq i \leq -2 + \lfloor \frac{n}{2} \rfloor, \\ \binom{2(-1 + \lfloor \frac{n}{2} \rfloor)}{-1 + \lfloor \frac{n}{2} \rfloor} r^{\lfloor \frac{n}{2} \rfloor - 1} (-r + 1)^{1 + \lfloor \frac{n}{2} \rfloor}, & i = -1 + \lfloor \frac{n}{2} \rfloor, \\ 2 \binom{2(-1 + \lfloor \frac{n}{2} \rfloor)}{-1 + \lfloor \frac{n}{2} \rfloor} r^{\lfloor \frac{n}{2} \rfloor} (-r + 1)^{\lfloor \frac{n}{2} \rfloor}, & i = \lfloor \frac{n}{2} \rfloor, \\ N_{n-i}^n(-r + 1), & 1 + \lfloor \frac{n}{2} \rfloor \leq i \leq n, \end{cases} \quad (9)$$

where $\lfloor k \rfloor$ denotes the greatest integer less than (GILT) or equal to k and $\lceil k \rceil$ denotes the least integer greater than (LIGT) or equal to k , respectively.

2.2 Definition: An NB1curve $N^n(r)$ of degree n with $n + 1$ control points, denoted by $\{\xi_i\}_{i=0}^n$, can be expressed power basis as [1]:

$$N^n(r) = \sum_{i=0}^n \sum_{j=0}^n \xi_i \cdot \hat{u}_{i,j} \cdot r^j, \quad (10)$$

where

$$\dot{u}_{i,j} = \begin{cases} (-1)^{-i+j} \binom{\lfloor \frac{n}{2} \rfloor + i - 1}{i} \binom{\lfloor \frac{n}{2} \rfloor}{j-i}, & 0 \leq i \leq -2 + \lfloor \frac{n}{2} \rfloor, \\ (-1)^{-i+j} \binom{2i}{i} \binom{i+2}{-i+j}, & i = -1 + \lfloor \frac{n}{2} \rfloor, \\ (-1)^{-i+j} 2 \binom{2(i-1)}{i-1} \binom{-i+n}{-i+j}, & i = \lfloor \frac{n}{2} \rfloor, \\ (-1)^{-i+j} 2 \binom{2(-i+n-1)}{-i+n-1} \binom{-i+n}{-i+j}, & i = \lfloor \frac{n}{2} \rfloor, \\ (-1)^{i+j-n} \binom{2(-i+n)}{-i+n} \binom{-i+n}{i+j-n-2}, & i = \lfloor \frac{n}{2} \rfloor + 1, \\ (-1)^{j-\lfloor \frac{n}{2} \rfloor} \binom{-i+n+\lfloor \frac{n}{2} \rfloor - 1}{-i+n} \binom{-i+n}{j-\lfloor \frac{n}{2} \rfloor}, & \lfloor \frac{n}{2} \rfloor + 2 \leq i \leq n. \end{cases} \quad (11)$$

The NB1 monomial matrix is given by:

$$\Gamma_{(n+1) \times (n+1)} = \begin{bmatrix} \dot{u}_{0,0} & \dot{u}_{0,1} & \dots & \dot{u}_{0,n} \\ \dot{u}_{1,0} & \dot{u}_{1,1} & \dots & \dot{u}_{1,n} \\ \vdots & \vdots & \ddots & \vdots \\ \dot{u}_{n,0} & \dot{u}_{n,1} & \dots & \dot{u}_{n,n} \end{bmatrix} \quad (12)$$

Where $\dot{u}_{i,j}$ is given in Eq. (11).

The following form of NB1 polynomials will be used to get approximate solutions of Eq. (1) and some appropriate ICs :

$$\omega_n(r) = \sum_{i=0}^n b_i N_i^n(r) \quad (13)$$

Where $b_i, i = 0, 1, \dots, n$ are unknown NB1 coefficients to be determined and $N_i^n(r)$ are the NB1 polynomials.

3. Fundamental Matrix Relations

The approximate solution (13) will be written as:

$$\omega_n(r) = \mathbf{N}(r) \mathbf{B} \quad (14)$$

Where $\mathbf{N}(r) = [N_0^n(r) \ N_1^n(r) \ \dots \ N_n^n(r)]$ and $\mathbf{B} = [b_0 \ b_1 \ \dots \ b_n]^T$.

Eq. (14) can be written as:

$$\omega_n(r) = \mathbf{X}_n(r) \Gamma^T \mathbf{B} = \psi(r) \mathbf{B} \quad (15)$$

Where $\psi(r) = \mathbf{X}_n(r) \Gamma^T$, $\mathbf{X}_n(r) = [1 \ r \ r^2 \ \dots \ r^n]$ and Γ is the monomial matrix, that is defined in (12).

3.1. The First Derivative of The Matrix Relation:

We will derive Eq. (14) as:

$$\begin{aligned} \omega_n'(r) &= \mathbf{N}'(r) \mathbf{B} \\ &= \psi'(r) \mathbf{B} \\ &= \mathbf{X}'_n(r) \Gamma^T \mathbf{B} \end{aligned}$$

$$\begin{aligned}
&= \frac{d}{dt} (\mathbf{X}_n(r)) \Gamma^T \mathbf{B} \\
&= [0 \quad 1 \quad 2r \quad \dots \quad nr^{n-1}] \Gamma^T \mathbf{B} \\
&= [1 \quad r \quad r^2 \quad \dots \quad r^n] \begin{bmatrix} 0 & 1 & 0 & \dots & 0 \\ 0 & 0 & 2 & \ddots & 0 \\ \vdots & \vdots & \ddots & \ddots & 0 \\ 0 & 0 & 0 & \dots & n \\ 0 & 0 & 0 & \dots & 0 \end{bmatrix} \Gamma^T \mathbf{B} \\
&= \mathbf{X}_n(r) \mathbf{V} \Gamma^T \mathbf{B},
\end{aligned}$$

where

$$\mathbf{V} = \begin{bmatrix} 0 & 1 & 0 & \dots & 0 \\ 0 & 0 & 2 & \ddots & 0 \\ \vdots & \vdots & \ddots & \ddots & 0 \\ 0 & 0 & 0 & \dots & n \\ 0 & 0 & 0 & \dots & 0 \end{bmatrix}_{(n+1) \times (n+1)}$$

Thus

$$\omega'_n(r) = \mathbf{X}_n(r) \mathbf{V} \Gamma^T \mathbf{B} \quad (16)$$

Similarly,

$$\omega''_n(r) = \mathbf{X}_n(r) \mathbf{V}^2 \Gamma^T \mathbf{B} \quad (17)$$

$$\begin{aligned}
&\vdots \\
\omega_n^{(m)}(r) &= \mathbf{X}_n(r) \mathbf{V}^m \Gamma^T \mathbf{B}, \quad m = 0, 1, 2, \dots \quad (18)
\end{aligned}$$

Where

$$\mathbf{V}^0 = \mathbf{I}_{n+1} = \begin{bmatrix} 1 & 0 & \dots & 0 \\ 0 & 1 & \ddots & \vdots \\ \vdots & \ddots & \ddots & 0 \\ 0 & \dots & 0 & 1 \end{bmatrix}_{(n+1) \times (n+1)} \quad \text{when } m = 0,$$

and $\omega_n^{(0)}(r) = \omega_n(r)$.

By using Equations (15) and (18) in Eq. (1), we get

$$\mathbf{X}_n(r) \mathbf{V}^{m+1} \Gamma^T \mathbf{B} + \frac{\mu_1}{r} \mathbf{X}_n(r) \mathbf{V}^m \Gamma^T \mathbf{B} + \frac{\mu_2}{r^2} \mathbf{X}_n(r) \mathbf{V}^{m-1} \Gamma^T \mathbf{B} + g(r) \mathbf{X}_n(r) \Gamma^T \mathbf{B} = h(r) \quad (19)$$

Eq. (19) can be written as the following:

$$\left(\mathbf{X}_n(r) \mathbf{V}^{m+1} + \frac{\mu_1}{r} \mathbf{X}_n(r) \mathbf{V}^m + \frac{\mu_2}{r^2} \mathbf{X}_n(r) \mathbf{V}^{m-1} + g(r) \mathbf{X}_n(r) \right) \Gamma^T \mathbf{B} = h(r) \quad (20)$$

By using the following collocation points:

$$r_i = \frac{i+1}{n}, \quad i = 1, 2, \dots, n, \quad (21)$$

we obtain

$$(\mathbf{X} \mathbf{V}^{m+1} + \mathbf{P}_0 \mathbf{X} \mathbf{V}^m + \mathbf{P}_1 \mathbf{X} \mathbf{V}^{m-1} + \mathbf{G} \mathbf{X}) \Gamma^T \mathbf{B} = \mathbf{H}, \quad (22)$$

where

$$\mathbf{X} = \begin{bmatrix} 1 & r_0 & r_0^2 & \cdots & r_0^n \\ 1 & r_1 & r_1^2 & \cdots & r_1^n \\ \vdots & \vdots & \vdots & \cdots & \vdots \\ 1 & r_n & r_n^2 & \cdots & r_n^n \end{bmatrix}_{(n+1) \times (n+1)}, \quad \mathbf{G} = \begin{bmatrix} g(r_0) & 0 & \cdots & 0 \\ 0 & g(r_1) & \ddots & \vdots \\ \vdots & \ddots & \ddots & 0 \\ 0 & \cdots & 0 & g(r_n) \end{bmatrix}_{(n+1) \times (n+1)}$$

$$\mathbf{P}_0 = \begin{bmatrix} \frac{\mu_1}{r_0} & 0 & \cdots & 0 \\ 0 & \frac{\mu_1}{r_1} & \ddots & \vdots \\ \vdots & \ddots & \ddots & 0 \\ 0 & \cdots & 0 & \frac{\mu_1}{r_n} \end{bmatrix}_{(n+1) \times (n+1)}, \quad \mathbf{P}_1 = \begin{bmatrix} \frac{\mu_2}{r_0} & 0 & \cdots & 0 \\ 0 & \frac{\mu_2}{r_1} & \ddots & \vdots \\ \vdots & \ddots & \ddots & 0 \\ 0 & \cdots & 0 & \frac{\mu_2}{r_n} \end{bmatrix}_{(n+1) \times (n+1)},$$

and $\mathbf{H} = [h(r_0) \ h(r_1) \ \cdots \ h(r_n)]^T$.

We can write Eq. (22) as the short form:

$$\mathbf{WB} = \mathbf{H} \quad \text{or} \quad [\mathbf{W}; \mathbf{H}], \quad (23)$$

where

$$\mathbf{W} = \mathbf{X} \mathbf{V}^{m+1} \Gamma^T + \mathbf{P}_0 \mathbf{X} \mathbf{V}^m \Gamma^T + \mathbf{P}_1 \mathbf{X} \mathbf{V}^{m-1} \Gamma^T + \mathbf{G} \mathbf{X} \Gamma^T$$

Now, we apply the proposed method to initial conditions, we have

$$\omega^{(k)}(0) = \mathbf{X}(0) \mathbf{V}^{(k)} \Gamma^T \mathbf{B}, \quad k = 0, 1, \dots, m \quad (24)$$

We can write Eq. (24) in the following matrix form:

$$\mathbf{UB} = \mathbf{O} \quad \text{or} \quad [\mathbf{U}; \mathbf{O}] \quad (25)$$

Next, we replace the last rows of the augmented matrix (23) with the rows of the initial conditions matrix (25), then the system (23) becomes

$$\widetilde{\mathbf{W}}\mathbf{B} = \widetilde{\mathbf{H}} \quad \text{or} \quad [\widetilde{\mathbf{W}}; \widetilde{\mathbf{H}}], \quad (26)$$

where

$$[\widetilde{\mathbf{W}}; \widetilde{\mathbf{H}}] = \begin{bmatrix} w_{0,1} & w_{0,2} & \cdots & w_{0,n} & ; & h(r_0) \\ w_{1,1} & w_{1,2} & \cdots & w_{1,n} & ; & h(r_1) \\ \vdots & \vdots & \cdots & \vdots & ; & \vdots \\ w_{n-(m+1),1} & w_{n-(m+1),2} & \cdots & w_{n-(m+1),n} & ; & h(r_n) \\ u_{0,1} & u_{0,2} & \cdots & u_{0,n} & ; & 0 \\ u_{1,1} & u_{1,2} & \cdots & u_{1,n} & ; & 0 \\ \vdots & \vdots & \cdots & \vdots & ; & \vdots \\ u_{m,1} & u_{m,2} & \cdots & u_{m,n} & ; & 0 \end{bmatrix}$$

If $\det(\widetilde{\mathbf{W}}) \neq 0$, we get

$$\mathbf{B} = (\widetilde{\mathbf{W}})^{-1} \widetilde{\mathbf{H}}, \quad (27)$$

which is determined uniquely.

By using Eq. (27), we can solve the linear system (26) of $(n + 1)$ unknown coefficients $b_i, i = 0, 1, \dots, n$ of NB1 polynomials in $(n + 1)$ equations to get the unknowns $b_i, i = 0, 1, \dots, n$. Finally, we obtained the solution of the studied problems.

4. Numerical Examples

Here, we present four examples of the singular linear ordinary differential equations (SLODEs) of various high orders.

Matlab program has carried out the computations of the examples.

Example 1: Consider the Lane-Emden equation [23]

$$\omega'' + \frac{8}{r}\omega' + r\omega = r^5 - r^4 + 44r^2 - 30r \quad (28)$$

With ICs.

$$\omega(0) = 0, \quad \omega'(0) = 0 \quad (29)$$

The exact solution to this problem is $\omega(r) = r^4 - r^3$.

Here, $g(r) = r$ and $h(r) = r^5 - r^4 + 44r^2 - 30r$.

We applied the mentioned method for $n = 4$, the collocation points are $r_0 = \frac{1}{4}, r_1 = \frac{1}{2}, r_2 = \frac{3}{4}, r_3 = 1, r_4 = \frac{5}{4}$, and the matrix equation (22) becomes:

$$\left[\mathbf{X}_4(r) \mathbf{V}^2 \Gamma^T + \mathbf{P}_0 \mathbf{X}_4(r) \mathbf{V} \Gamma^T + \mathbf{G} \mathbf{X}_4(r) \Gamma^T \right] \mathbf{B} = \mathbf{H}, \quad (30)$$

where

$$\Gamma^T = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ -2 & 2 & 0 & 0 & 1 \\ 1 & -6 & 4 & 0 & 0 \\ 0 & 6 & -8 & 2 & 0 \\ 0 & -2 & 4 & -2 & 0 \end{bmatrix}, \quad \mathbf{V} = \begin{bmatrix} 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 2 & 0 & 0 \\ 0 & 0 & 0 & 3 & 0 \\ 0 & 0 & 0 & 0 & 4 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}, \quad \mathbf{V}^2 = \begin{bmatrix} 0 & 0 & 2 & 0 & 0 \\ 0 & 0 & 0 & 6 & 0 \\ 0 & 0 & 0 & 0 & 12 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix},$$

$$\mathbf{P}_0 = \begin{bmatrix} 32 & 0 & 0 & 0 & 0 \\ 0 & 16 & 0 & 0 & 0 \\ 0 & 0 & 32/3 & 0 & 0 \\ 0 & 0 & 0 & 8 & 0 \\ 0 & 0 & 0 & 0 & 32/5 \end{bmatrix}, \quad \mathbf{G} = \begin{bmatrix} 1/4 & 0 & 0 & 0 & 0 \\ 0 & 1/2 & 0 & 0 & 0 \\ 0 & 0 & 3/4 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 5/4 \end{bmatrix}, \quad \mathbf{H} = \begin{bmatrix} -4867/1024 \\ -129/32 \\ 1283/591 \\ 14 \\ 7073/222 \end{bmatrix},$$

$$\mathbf{X}_4 = \begin{bmatrix} 1 & 1/4 & 1/16 & 1/64 & 1/256 \\ 1 & 1/2 & 1/4 & 1/8 & 1/16 \\ 1 & 3/4 & 9/16 & 27/64 & 81/256 \\ 1 & 1 & 1 & 1 & 1 \\ 1 & 5/4 & 25/16 & 125/64 & 625/256 \end{bmatrix}$$

Therefore, the augmented matrix (23) becomes

$$[\mathbf{W}; \mathbf{H}] = \begin{bmatrix} -2935/64 & -2277/512 & 5897/256 & 4867/512 & 513/16 & ; & -4867/1024 \\ -111/8 & -127/16 & -31/8 & 129/16 & 65/4 & ; & -129/32 \\ -631/192 & -1765/1536 & -2277/256 & -2223/512 & 539/48 & ; & 1283/591 \\ 2 & 0 & 8 & -28 & 9 & ; & 14 \\ 1689/320 & -19837/2560 & 12157/256 & -32625/512 & 637/80 & ; & 7073/222 \end{bmatrix} \quad (31)$$

By using initial conditions (29), we have from Eq. (25), the matrix form:

$$[\mathbf{U}; \mathbf{O}] = \begin{bmatrix} 1 & 0 & 0 & 0 & 0; 0 \\ -2 & 2 & 0 & 0 & 1; 0 \end{bmatrix} \quad (32)$$

Now, replace the last two rows of the augmented matrix (31) by matrix (32), and we get

$$[\tilde{W}; \tilde{H}] = \begin{bmatrix} -2935/64 & -2277/512 & 5897/256 & 4867/512 & 513/16 & ; & -4867/1024 \\ -111/8 & -127/16 & -31/8 & 129/16 & 65/4 & ; & -129/32 \\ -631/192 & -1765/1536 & -2277/256 & -2223/512 & 539/48 & ; & 1283/591 \\ 1 & 0 & 0 & 0 & 0 & ; & 0 \\ -2 & 2 & 0 & 0 & 1 & ; & 0 \end{bmatrix} \quad (33)$$

By using Eq. (27), the solution of system (33) gives the coefficients of NB1 polynomials:

$$b_0 = 0.00, b_1 = 0.00, b_2 = 0.00, b_3 = -0.50 \text{ and } b_4 = 0.00.$$

That is,

$$\mathbf{B} = \begin{bmatrix} b_0 \\ b_1 \\ b_2 \\ b_3 \\ b_4 \end{bmatrix} = \begin{bmatrix} 0.00 \\ 0.00 \\ 0.00 \\ -0.50 \\ 0.00 \end{bmatrix}$$

Substituting these values into Eq. (14), then the approximate solution is

$$\begin{aligned} \omega_4(r) &\cong \mathbf{N}(r)\mathbf{B} \\ &= \mathbf{X}_4(r) \Gamma^T \mathbf{B} \\ &= \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ -2 & 2 & 0 & 0 & 1 \\ 1 & -6 & 4 & 0 & 0 \\ 0 & 6 & -8 & 2 & 0 \\ 0 & -2 & 4 & -2 & 0 \end{bmatrix} \begin{bmatrix} 0 \\ 0 \\ 0 \\ -1/2 \\ 0 \end{bmatrix} \\ &= -r^3 + r^4 = \omega(r) \quad (\text{exact solution}) \end{aligned}$$

Example 2: Consider the Emden-Fowler equation [27]

$$\omega^{(4)} + \frac{7}{r}\omega^{(3)} + \frac{9}{r^2}\omega^{(2)} + \omega = 300 + r^4, \quad (34)$$

with ICs.

$$\omega(0) = \omega'(0) = \omega''(0) = \omega'''(0) = 0, \quad (35)$$

$\omega(r) = r^4$ is the exact solution.

As above example, we apply the proposed method for $n = 4$. Then Eq. (34) takes the following form as Eq. (20):

$$\left[\mathbf{X}_4(r)\mathbf{V}^4\Gamma^T + \frac{7}{r}\mathbf{X}_4(r)\mathbf{V}^3\Gamma^T + \frac{9}{r^2}\mathbf{X}_4(r)\mathbf{V}^2\Gamma^T + \mathbf{X}_4(r)\Gamma^T \right] \mathbf{B} = 300 + r^4, \quad (36)$$

By using the collocation points, Eq. (36) becomes

$$\mathbf{WB} = \mathbf{H}, \quad (37)$$

where $\mathbf{W} = \mathbf{X}\mathbf{V}^4\Gamma^T + \mathbf{P}_0\mathbf{X}\mathbf{V}^3\Gamma^T + \mathbf{P}_1\mathbf{X}\mathbf{V}^2\Gamma^T + \mathbf{G}\mathbf{X}\Gamma^T$ such that

$$\mathbf{P}_0 = \text{diag}(28, 14, 28/3, 7, 28/5),$$

$$\mathbf{P}_1 = \text{diag}(144, 36, 16, 9, 144/25),$$

$$\mathbf{G} = \text{diag}(1, 1, 1, 1, 1),$$

$$\mathbf{H} = [76801/256 \quad 4801/16 \quad 23725/79 \quad 301 \quad 10283/34]^T,$$

\mathbf{X} and Γ^T are the same as example 1.

Replacing the last four rows of matrix (37) by the rows of initial conditions matrix (35), and we use Eq. (27) to get the vector matrix \mathbf{B} as the following.

$$\mathbf{B} = [0 \quad -1/2 \quad -3/4 \quad -3/2 \quad 1]^T$$

Thus

$$\begin{aligned}
 \omega_4(r) &\cong N(r)\mathbf{B} \\
 &= \mathbf{X}_4(r) \Gamma^T \mathbf{B} \\
 &= \begin{bmatrix} 1 & r & r^2 & r^3 & r^4 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ -2 & 2 & 0 & 0 & 1 \\ 1 & -6 & 4 & 0 & 0 \\ 0 & 6 & -8 & 2 & 0 \\ 0 & -2 & 4 & -2 & 0 \end{bmatrix} \begin{bmatrix} 0 \\ -1/2 \\ -3/4 \\ -3/2 \\ 1 \end{bmatrix} \\
 &= r^4 = \omega(r).
 \end{aligned}$$

Example 3: Consider (SLODE) [15]

$$\omega^{(5)} + \frac{3}{r}\omega^{(4)} + 5!\omega = 4 + r^5, \quad (38)$$

with ICs.

$$\omega(0) = \omega'(0) = \omega''(0) = \omega'''(0) = \omega^{(4)}(0) = 0 \quad (39)$$

The exact solution is $\omega(r) = \frac{1}{5!} r^5$.

Applying the proposed method for $n = 5$ yields that

$$\mathbf{B} = [0 \quad 0 \quad -1/480 \quad 0 \quad -1/80 \quad 1/120]^T$$

so we obtain

$$\begin{aligned}
 \omega_5(r) &\cong N(r)\mathbf{B} \\
 &= \mathbf{X}_5(r) \Gamma^T \mathbf{B} \\
 &= \begin{bmatrix} 1 & r & r^2 & r^3 & r^4 & r^5 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ -2 & 2 & 0 & 0 & 0 & 0 \\ 1 & -6 & 4 & 0 & 0 & 1 \\ 0 & 6 & -12 & 4 & 2 & 0 \\ 0 & -2 & 12 & -8 & -2 & 0 \\ 0 & 0 & -4 & 4 & 0 & 0 \end{bmatrix} \begin{bmatrix} 0 \\ 0 \\ -1/480 \\ 0 \\ -1/80 \\ 1/120 \end{bmatrix} \\
 &= \frac{1}{5!} r^5 = \omega(r).
 \end{aligned}$$

Example 4: Consider (SLODE) [15]

$$\omega^{(6)} + \frac{1}{r}\omega^{(5)} + 6!\omega = -r^6 + 2 \quad (40)$$

With ICs.

$$\omega(0) = \omega'(0) = \omega''(0) = \omega'''(0) = \omega^{(4)}(0) = \omega^{(5)}(0) = 0 \quad (41)$$

The exact solution is $\omega(r) = \frac{1}{6!} r^6$.

In the same manner, we applied the proposed method when $n = 6$ and obtained

$$\mathbf{B} = [0 \quad 0 \quad 1/690 \quad 1/1080 \quad 1/720 \quad -1/360 \quad 1/720]^T$$

after that we substituted these values of \mathbf{B} in Eq. (15) thus getting the exact solution

$$\omega(r) = \omega_6(r) = \frac{1}{6!} r^6.$$

5. Conclusion

The singular linear ordinary differential equations (SLODEs) of various high orders were solved by using the collocation method based on NB1 Polynomials. The proposed method reduced the given problem,

which included the equation and appropriate initial conditions, into a linear algebraic equations system. The resulting system was solved by using the inverse matrix method. Thus we obtained the exact solutions to each of the studied problems, showing the potency, reliability and accuracy of the presented method.

References:

1. Aphirukmatakun C. and Dejdumrong N., "Monomial forms for curves in CAGD with their applications, Sixth International Conference on Computer Graphics", Imaging and Visualization, IEEE, (2009), 211-216.
2. Assabaai M. A. and Khred A., "Numerical Solutions of Singular Nonlinear Ordinary Differential Equations using Said-Ball Polynomials", Emirates Journal for Engineering Research, 27 (4) December (2022).
3. Bataineh A. S., Noorani M. S. M. and Hashim I., "Homotopy analysis method for singular initial value problems of Emden-Fowler type", Commun Nonlinear Sci Numer Simulat, 14 (2009), 1121-1131.
4. Bhatti A. H. and Karim S. B., "Solving Ordinary Differential Equations (ODEs) Using Least Square Method Based on Wang Ball Curves", Mathematics and Statistics, 10 (1) (2022), 74-87.
5. Bhrawy A. H. and Alofi A. S., "A Jacobi- Gauss collocation method for solving nonlinear Lane-Emden type equations", Commun Nonlinear Sci Numer Simulat, ,17 (2012), 62-70.
6. Chandrasekhar S., "Introduction to the Study of Stellar Structure", Dover, New York, (1967).
7. Chowdhury M. S. H. and Hashim I., "Solutions of Emden-Fowler Equations by homotopy- perturbation Method", Nonlinear Analysis: Real World Applications, 10 (2009), 104-115.
8. Dan Yu & Xinmeng Chen, "ANOTHER TYPE OF GENERALIZED BALL CURVES AND SURFACES", Acta Mathematica Scientia, 27B (4) (2007), 897-907.
9. Davis H. T., "Introduction to Nonlinear Differential and Integral Equations", Dover, New York, (1962).
10. Dehghan M. and Shakeri F., "Approximate solution of a differential equation arising in astrophysics using the variational iteration method", New Astronomy, 13 (2008), 53-59.
11. Demir H. and Sungu I. C., "Numerical solution of a class of nonlinear Emden-Fowler equations by using differential transformation method", Journal of Arts and Science, 12 (2009), 75-81.
12. Fayek S., El-Gamel M. and El-Baghdady G. I., "Solving Linear and Nonlinear Singular Differential Equations Using Bessel Matrix Method", International Journal of Innovative Science and Research Technology, 6 (3) (2021), 423-427.
13. Fowler R. H., "Further studies of Emdens and similar differential equations", Quarterly Journal of Mathematics, 2 (1) (1931), 259-288.
14. Gümğüm S., "Taylor wavelet solution of linear and nonlinear Lane-Emden Equations", Applied Numerical Mathematics, 158 (2020), 44-53.
15. Hasan Y. Q. and Mutaish A. M. S., "Solving Higher Order Singular Initial Value Problems by Adjusted Adomian Decomposition Strategy", IIARD International Institute of Academic Research and Development, 4(3) (2018), 1-9.
16. Hasan Y. Q., Osilagun J. A. and Adegbindin A. O., "Approximate Analytical Solution of a Class of Singular Differential Equations with Dirichlet Conditions by the Adomian Decomposition Method", IJMAO International of Mathematical Analysis and Optimizations: Theory and Applications, 1 (2019), 433-442.
17. He J. H., "Variational iteration method for autonomous ordinary differential systems", Appl. math. Comput., 114(2/3) (2000), 115-123.
18. Kashem B. E., "Modified Hermite Operational Matrix Method for Nonlinear Lane-Emden Problem", Al-Qadisiyah Journal of Pure Science, 25 (3) (2020), 71-79.
19. Kaur H., Mittal R. C., and Mishra V., "Modified Hermite Operational Matrix Method for Nonlinear Lane-Emden Problem", Computer Physics Communications, 184 (2013), 2169-2177.

20. Khalique C.M., Mahomed F. M. and Muatjetjeja B., "Lagrangian formulation of a generalized Lane–Emden equation and double reduction", *J. Nonlinear Math. Phys.*, 15 (2008), 152–161.
21. Khred A., Bamsaoud S. F., Bazighifan O. and Assabaai, M. A., "Improved Operational Matrices of DP-Ball Polynomials for Solving Singular Second Order Linear Dirichlet-type Boundary Value Problems", *Hadhramout University Journal of Natural & Applied Sciences*, 19(1) (2022), 39-50.
22. Khred A., Omar Z., Saaban A. and Adeyeye O., "Solving first and second order delay differential equations using new operational matrices of Said-Ball polynomials", *Journal of Interdisciplinary Mathematics*, 24(4) (2021), 921–930.
23. Khred A., Saaban A., Adyee and Fadhel, I. E. I., "Wang-Ball Polynomials for the Numerical Solution of Singular Ordinary Differential Equations", *Iraqi Journal of Science*, 62 (2021), 941-949.
24. Lane I. J. H., "On the theoretical temperature of the sun under the hypothesis of a gaseous mass maintaining its volume by its internal heat and depending on the laws of gases known to terrestrial experiment", *American Journal of Science and Arts*, 4 (1870), 57-74.
25. Momoniat E. and Harley C., "Approximate implicit solution of a Lane–Emden equation", *New Astronomy*, 11 (2006), 520–526.
26. Muatjetjeja B. and Khalique C.M., "Exact solutions of the generalized Lane–Emden equations of the first and second kind", *Pramana*, 77 (2001), 545–554.
27. Mutaish A. M. S. and Hasan Y. Q., "Adjusted Adomian Decomposition Method for Solving Emden-Fowler Equations of Various Order", *MAYFEB Journal of Mathematics*, 3 (2018), 1-10.
28. Parand K., Dehghan M., Rezaeia A. R. and Ghaderi S. M., "An approximation algorithm for the solution of the nonlinear Lane–Emden type equations arising in astrophysics using Hermite functions collocation method", *Comput. Phys. Commun.*, 181 (2010), 1096–1108.
29. Parand K., Shahini, M. and Dehghan M., "Rational Legendre pseudospectral approach for solving nonlinear differential equations of Lane–Emden type", *J. Comput. Phys.*, 228 (2009), 8830–8840.
30. Rajaraman R. and Hariharan G., "An efficient wavelet based spectral method to singular boundary value problems", *J Math Chem*, 53 (2015), 2095–2113.
31. Richardson O.U., "The Emission of Electricity from Hot Bodies", Longmans Green and Company, London, (1921).
32. Singh O. P., Pandey R. K. and Singh V. K., "Analytical algorithm of Lane-Emden type equation arising in astrophysics using modified homotopy analysis method", *Computer Physics Communications*, 180 (2009), 1116-1124.
33. Singh. R., Garg H. and Guleria V., "Haar wavelet collocation method for Lane-Emden equations with Dirichlet, Neumann and Neumann-Robin boundary conditions", *Journal of Computational and Applied Mathematics*, <https://doi.org/10.1016/j.cam.2018.07.004>.
34. Singh R., Guleria V. and Singh M., "Haar wavelet quasilinearization method for numerical solution of Emden–Fowler type equations", *Mathematics and Computers in Simulation*, 174 (2020), 123–133.
35. Singh R. and Kumar J., "An efficient numerical technique for the solution of nonlinear singular boundary value problems", *Computer Physics Communications*, 185 (2014), 1282–1289.
36. Shang X., Wu P. and Shao X., "An efficient method for solving Emden–Fowler equations", *Journal of the Franklin Institute*, 346 (2009), 889–897.
37. Verma A. K. and Tiwari D., "Higher resolution methods based on Quasilinearization and Haar Wavelets on Lane-Emden Equations", *international Journal of Wavelets, Multiresolution and Information Processing*, (2018), <https://doi.org/10.1142/S021969131950005X>.
38. Wang T., Liu Z. and Kong, Y., "The series expansion and Chebyshev collocation method for nonlinear singular two-point boundary value problems", *J. Eng. Math.*, 126 (5) (2021), 1-28.
39. Wazwaz A. M., "Adomian decomposition method for a reliable treatment of the Emden–Fowler equation", *Appl. Math. Comput.*, 161 (2005), 543–560.
40. Wazwaz A. M., "A new algorithm for solving differential equations of Lane–Emden type", *Appl. Math. Comput.*, 118 (2001), 287–310.

41. Wazwaz A. M., "A new method for solving singular initial value problems in the second-order ordinary differential equations", *Appl. Math. Comput.*, 128 (2002), 47–57.
42. Wazwaz A. M., Rach R., Bougoffa L. and Duan J. S., "Solving the Lane–Emden–Fowler Type Equations of Higher Orders by the Adomian Decomposition Method", *CMES*, 100 (2014), 507-529.
43. Wazwaz A. M., "Solving the Non-isothermal reaction Diffusion Model Equations in a Spherical Catalyst by the Variational Iteration Method", *Chemical Physics Letters*, 679 (2017), 132-136.
44. Wazwaz A. M., "Solving Two Emden–Fowler Type Equations of Third Order by the Variational Iteration Method", *Applied Mathematics & Information Sciences*, 5 (2015), 2429-2436.
45. Wazwaz A. M., "The variational iteration method for solving nonlinear singular boundary value problems arising in various physical models", *Commun. Nonlinear Sci. Numer. Simulat.*, 16 (2011), 3881–3886.
46. Yildirim A. and Ozis T., "Solutions of singular IVPs of Lane–Emden type by the variational iteration method, *Nonlinear Analysis*", 70 (2009), 2480–2484.
47. Zomot N. H. and Ababneh O. Y., "Solution of Differential Equations of Lane-Emden Type by Combining Integral Transform and Variational Iteration Method", *Nonlinear Analysis and Differential Equations*, 4(3) (2016), 143 – 150.