

Original Research Article

Effect of different IBA concentrations on air layering in Olive (*Olea europaea*L.)

Comment [A1]: It is recommended to change the title since it is the same as another work presenting, only that it changes the culture.

ABSTRACT

The present investigation entitled "Effect of different IBA concentrations on air layering in olive (*Olea europaea* L.)" was carried out in the Department of Horticulture, Sam Higginbottom University of Agriculture Technology and Sciences, Prayagraj in August 2022 to December 2022. The experiment was conducted to evaluate the best IBA concentrations among different concentrations of IBA for producing best layering and survival percentage in Olive. The experiment was laid out in Randomized Block Design having 8 treatments and replicated 3 times. The treatment T₄ in which IBA 4000 ppm was applied for air layering was found to be best in terms of shoot and root parameters observed, i.e., Number of days to root formation (46.11), Number of roots per layer (7.33), Root length (6.73 cm), Root thickness (1.66 mm), Number of new sprouts (2.67-1MAP, 3.89-2MAP & 5.67-3MAP), length of new shoot (2.78cm-1MAP, 3.73cm-2MAP & 5.44cm-3MAP), Number of new leaves (7.22-1MAP, 10.78-2MAP & 16.56-3MAP) and survival percentage (50.89%) in air layered olive plants, whereas minimum values were recorded in T₈ (control). It can be inferred from this study that, the use of IBA 4000 ppm was effective to produce better root and shoot parameters and successful establishment of air layers in olive.

Comment [A2]: Deletar

Comment [A3]: What is the objective of the research?

Keywords: *Air layering, Olive, IBA, Shoot Parameters and Root Parameters.*

Comment [A4]: Keywords should not be present in the title.

1. INTRODUCTION

Olive (*Olea europaea* L.) is a hardy, dicotyledonous, attractive evergreen subtropical fruit tree with grey foliage and is native of Mediterranean region. It belongs to family Oleaceae, which is also known as Elaeocarpaceae, having 30 genera and 600 species. It is known as the 'tree of liquid gold' because of great importance of its oil and green olives are considered as 'Green gold' in Greece. Olive tree is considered as Symbol of prosperity and peace.

Comment [A5]: Which authors claim this.

In India olive is a new crop, which was introduced in 1800 A.D. by British Missionaries, and its cultivation is still in its infancy stage in India. Olive is one of the important and oldest fruit trees in the world (Standish, 1960) and is successfully grown in the

areas having mean temperatures 15-20°C, with a minimum of 4°C and maximum of 40°C(Therios, 2009). The cultivated olive tree can reach heights ranging from just a few meters to 20 m. The tree starts bearing in 4 to 8 years, but commercial production is not reached for 15 or 20 years. The root system is generally shallow, spreading to 0.9 - 1.2 m even in deep soils. The above ground portion of the olive tree is recognizable by the dense assembly of limbs, short internodes, and compact nature of the foliage. Leaves are opposite, thick, leathery, growing over two years period.

Olive production in the country began in 2007 when olive saplings were imported from Israel and planted in the Thar Desert. The first olive yield in India occurred in 2012, and commercial olive oil production began in September 2013. The first Indian-made olive oil brand called Raj Oil was launched on 9 November 2016. India produced 150 tons of Olives in 2020. Rajasthan and Gujarat are major trading states in India. The economic part is fruit, which is a small drupe, 1-2.5 cm long when ripe. Olive cultivars may be used primarily for oil, eating, or both. Olives cultivated for consumption are generally referred to as “table olives”. About 92 percent olives are turned in to oil, while about 10 percent are used as table olives.

Comment [A6]: The source from which the information was obtained must be specified.

Ripe olive contains proteins (2%), oil (14%), vit. A, B, C and (1%) Fibre and 650 Calories per pound of fresh weight. One tablespoon oil contains 120 calories, (1.8%) protein, (21%) fat, (2.6%) Carbohydrates and (1.5%) fibre. Oil is rich source of polysaturated fatty acids and is free from cholesterol. Oil is also rich in minerals like Fe, Ca, P and Vitamin E, and it contains oleic acid (70-80%) and linolenic acid (7-12%). Oil has many industrial as well as culinary uses. It is edible and possesses therapeutic value and it is clinically advocated for hypertension, coronary heart disease and muscular pains. Oil consumption helps to reduce LDL and to raise HDL forms for cholesterol in blood. Oil is also used in textile lubricants, perfumed soaps, cosmetics, perfumery industry and burning of lamps.

Comment [A7]: The source from which the information was obtained must be specified.

Olive can be propagated through sexual and asexual methods but the sexual method is usually not recommended, because seedlings are not true to type and the juvenile period is too long to bear fruit. Asexual propagation include hardwood, semi hardwood cuttings and air-layering. Air layering is a method of producing plants from aerial branches which remain in position while rooting. In plants which are difficult to root, air layering has been proven as an effective method of increasing their rooting percentage.

Physiological basis of layering involves accumulating organic substances like carbohydrates, hormones, and growth factors in a stem where rooting is desired by interrupting their downward translocation from the shoot tips and leaves. The rooting ability of layering can be improved by preconditioning treatments such as etiolation, girdling, periderm stripping, and application of growth regulators like auxins. The accumulated substances stimulate the formation of adventitious roots in the layered stem, which then grow into the soil, enabling the stem to be cut from the parent plant.

Comment [A8]: The source from which the information was obtained must be specified.

Air layering is an easy method of propagation of horticultural crops and it is widely used for commercial multiplication. The growth, establishment and survival of branches from air layering plants depend on the quality of rooting media. There are many commercial rooting media used for layering, but many are expensive and locally unavailable. Cocopeat is a multipurpose growing medium made out of coconut husk. The fibrous coconut husk is pre washed, machine dried, sieved and made free from sand and other contaminations such as animal and plant residue. Increasing demand and mounting costs for peat, sphagnum moss etc. as a growing media in horticulture have led to the search for high quality and low-cost substrates as an alternative. Cocopeat is considered as a good growing media component containing good source of nutrients with acceptable pH, (Abad *et al.* 2002) it can also be mixed easily with other substrates.

The latest advance in the knowledge of growth regulators in plant propagation has further improved the scope of their use in vegetative propagation of various fruit crops. Application of root promoting substances during layering helps to get profuse roots within a short time. The use of plant growth regulators to increase the efficiency of propagation in cuttings and layering are now common. Growth substances accelerate the rooting, produce large root system and increase the percentage of survival. Growth regulators like IAA, NAA and IBA have been used to stimulate plant growth and specially root formation in layering. In air layering experiments among auxins, IBA has been found to be most effective in producing maximum number of roots with better vigour. IBA is widely used as a root-initiation promoter in horticultural crops. IBA is a suitable rooting hormone for this type of experiment because it shows a large amount of flexibility when dealing with the range of concentration that can be used.

Comment [A9]: What is the objective of the research?

2. MATERIALS AND METHODS

The experiment was conducted from August 2022 to February 2023, at the Horticulture Research Farm, Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences (SHUATS), Prayagraj (U.P.) which is located at 25° 39' 42" N latitude, 81° 67' 56" E longitude and 98 m altitude above the mean sea level.

The present investigation was carried out on semi-hardwood stems of Olive. Plant growth regulator, Indole -3-Butyric Acid was used as a root induction chemical at seven different concentrations and combinations replicated thrice in randomized block design. Cocopeat was used as the root growing media in layering and soil, sand and vermicompost in equal proportion were used as growing media in polybags at the time of transplantation of rooted air layered plants. Treatments were listed of T₁ (IBA 1000 ppm), T₂ (IBA 2000 ppm), T₃ (IBA 3000 ppm), T₄ (IBA 4000 ppm), T₅ (IBA 5000 ppm), T₆ (IBA 1000 ppm), T₇ (IBA 7000 ppm) and T₈ (Control).

To conduct air layering on Olive trees, healthy branches of one-year, old uniform vigour, and pencil thickness were selected. Only disease-free branches were chosen, and a careful girdling of the bark, about 2-2.5 cm, was done just below the bud. Two circular cuts were made around 45-60 cm below the top end of the selected shoot, being careful not to harm the underlying wood. Pre-prepared IBA at different concentrations were applied evenly on all sides of the upper cut of the ringed portion using a brush. The cut portion was then covered with coco peat as rooting media and transparent polythene wrappers were used to wrap the air layered portion. Observations were made after 1, 2 and 3 month(s) from the day of operation, recording various root parameters such as the number of days for root formation (initial rooting), number of roots, root length and root thickness. After detaching the air layers by making a cut just below the lowest end of the ringed surface with sharp secateurs, they were transplanted into the fruit nursery of the Department of Horticulture at SHUATS. Shoot parameters and survival percentage were recorded after 1, 2 and 3 months after planting into polybags.

3. RESULTS AND DISCUSSION

3.1 Root parameter

In olive air layers the minimum number of days taken for initial root formation 46.11 days was observed in T₄ (IBA 4000 ppm) which was on par with treatment T₅ (IBA 5000 ppm) which

Comment [A10]: Is there research to support these doses?

Comment [A11]: The design used is not specified, as well as the statistical analysis carried out.

Comment [A12]: No table is presented to better see the results. Not discussed

took 62.11 days for root formation. It was followed by treatment T₁ (IBA 1000 ppm) 68.88 days which took for root initiation. Maximum number of roots (7.33) were recorded in Treatment T₄ (IBA 4000 ppm) which was significantly superior to all other treatments. It was followed by treatment T₅ (IBA 5000 ppm (6.44), which was on par with treatment T₆ (IBA 6000ppm) which had 6.00 roots. Treatment T₄ (IBA 4000 ppm) recorded the longest root length (6.73 cm) which was statistically on par with Treatment T₅ (IBA 5000 ppm) which measured root length of 6.20 cm. It was followed by treatment T₂ (IBA 2000ppm) with (5.19 cm) root length, it was on par with treatment T₆ (5.15 cm), T₁ (4.82 cm) and T₇ (4.68 cm). Maximum root thickness (1.66 mm) was recorded in Treatment T₄ (IBA 4000 ppm.) which was on par with treatment T₅ (1.60 mm), T₃ (1.53 mm) and treatment T₆ (1.52mm).

3.2 Shoot parameters

Statistical analysis has revealed significant variance among the different treatments with respect to all shoot parameters recorded. Maximum number of new sprouts in transplanted olive air layers at 1, 2 and 3 month(s) after transplanting (2.67), (3.89) and (5.67) was observed in T₄ (IBA@4000 ppm). Maximum shoot length in transplanted olive air layers at 1, 2 and 3 month(s) after transplanting was 2.78 cm, 3.73 cm and 5.44 cm respectively it was recorded in T₄ (IBA@4000 ppm). Maximum number of new leaves in transplanted olive air layers at 1, 2 and 3 month(s) after transplanting 7.22, 10.78 and 16.56 was recorded in treatment, T₄ in which IBA 4000 ppm was applied

Comment [A13]: What kind of analysis was performed?
Not discussed

3.3 Survival percentage

Maximum survival rate in transplanted olive air layers was 57.56%, 54.44% and 50.89% at one month after transplanting, two months after transplanting and three months after transplanting was recorded in treatment T₄ (IBA@4000 ppm).

Comment [A14]: No table is presented to better see the results.
Not discussed

4. Conclusion

The study revealed that different concentrations of IBA had significant effect on shoot, root parameters and survival percentage in olive air layering. From the study, it is concluded that treatment T₄ (IBA 4000 ppm) was found to be best in terms of shoot and root Parameters observed. The least number of days to root formation (46.11 days), number of roots per layer (7.33), root length (6.73 cm), root thickness (1.66 mm), number of new sprouts (5.67), length of new shoot (5.44 cm), and number of new leaves (16.56), survival percentage (50.89%) was recorded in treatment T₄, where, 4000 ppm IBA was used in air layering. So, IBA 4000 ppm can be suggested for air layering based on this study as successful vegetative propagation method in olive.

Comment [A15]: It is recommended to adjust the conclusion once the statistical data have been presented. The values they present are non-conclusive results.

5. References

- Abad, M., Noguera, P., Puchades, R., Maquieira, A. and Noguera, V. (2002).** Physio-chemical and chemical properties of some coconut dusts for use as peat substitute for containerized ornamental plants. *Biores. Tech.*, 8(3): 241-245
- Alam, R. and Sajid, M. (2017).** Effect of timing intervals on the rooting response and performance of olive cultivars through air layerage. *Sarhad J. Agric.* 34(1): 31-39.
- Alam, R., Sajid, M., and Shahwar, D. (2022).** Influence of IBA Concentrations on the rooting response of olive Cultivars in Air-Layering. *Sarhad J. Agric.* 38(1): 41-47.
- Afzal, F., Adnan, M., Rahman, I. U., Noor, M., Khan, A., Iqbal, S., and Subhan, F. (2017).** Growth response of olive cultivars to air layering. *Pure Applied Biol.* 6(4): 1403-1409.
- Ameen Ayesha, Jalil Ahmad and Shahid Raza (2017)** Effect of different concentrations of IBA on production of coratina olive plant. *Sci. Res. J.* 3(11): 125-128.
- Baghel, M., Raut, U.A. and Ramteke, V. (2016).** Effect of IBA concentrations and time of air-layering in guava cv. L-49. *Res. J. Agri. Sci.*, 7(1):117-120.
- Bose T.K., Mitra S.K. & Sadhu M.K. (1986).** Guava In: Propagation of tropical and sub-tropical horticultural crops. *Naya Prakash, Calcutta*, pp. 291–301

- Bhumika Verma and GD Sahu (2022)** Studies on the effect of different rooting media on survival and success of air layering in acid lime (*Citrus aurantifolia* Swingle) under Chhattisgarh plain. *Pharma Inn. J.*; 11(12): 225-228
- Chinnapa, K.C. (1962).** Effect of plant growth regulators on the rootage of air layers in Litchi. Lalbagh, 7: 28-31.
- Hartmann, H.T. and Kester, D.E. (1975).** *Plant Propagation Principles and Practices*. 3rd Edition. Printice Hall, New Jersey.
- Ibrahim, Omer. (2020).** Developing air layering practices for propagation of *Dracaena marginata* Lam. Utilizing phloroglucinol and seaweed extract as IBA-synergists or alternatives. *Sci. J. of Flowers and Ornamental Plants*. 7. 185-197.
- Kumar, V., Singh M.K., Kumar, M., Prakash, S., Kumar, A., Rao, S. and Malik, S. (2015).** Effect of different doses and rooting media on rooting of stem cutting of lemon (*Citrus lemon* Burm) cv. Pant Lemon-1. *J. of Plant Develop. Sci.*, 7(7): 587-591.
- Lalramhluna, P. and Prasad, M.V. (2016).** Effect of different levels of Indole-3- butyric acid on growth, development, survival and establishment of air layering lemon (*Citrus lemon* L.) cv. Assam Lemon under Allahabad agro-climatic conditions. *Int. J. Life Sciences Scientific Res.*, 2(5): 599-600.
- Naithani, C.D., Nautiyal, R.A., Rana, K.D. and Mewar, D. (2018).** Effect of time of air layering, IBA concentrations, growing ratio and their interaction on the rooting behaviour of Pant Prabhat guava (*Psidium guajava* L.) under sub-tropical condition of Garwahl Himalayas. *Int. J. of Pure and App. Biosci.*, 6(3): 169-180.
- Rymbai, H. and Reddy, S.G. (2010)** Effect of IBA, time of layering and rooting media on air layering and plantlets survival under different growing nursery conditions in guava. *Indian J. of Hort.*, 67: 99-104.
- Sanchezuedaneta, A.B., Suarez, E., Gonzalez, R.M., Amaya, Y., Colmenares C. B., and Ortega, J. (2009).** Effect of IBA on the air layer rooting of guava (*Psidium guajava* L.) in the Barat Country, Venezuela. *Pre. Evaluat. Revista UDO Agricola*, 9(1):113-120.
- Standish, R. (1960).** *The First Trees. The Story of the Olive.* Phoenix House, London

Syed Abdul Qadir Gilani, Kamran Shah, Imran Ahmed, Abdul Basit, Muhammad Sajid, Aneela Shah Bano, Gulshan Ara and Umaira Shahid (2019). Influence of indole butyric acid (IBA) concentrations on air layerage in guava (*Psidium guajava* L.) cv. Safeda. *Pure & App. Bio.* Vol. 8, Issue 1, pp355-362.

Therios. and Ioannis Nikolaos (2009). [Olives: Volume 18 of Crop Production Science in Horticulture \(History of Olive Growing, page 1\)](#). CABI. ISBN 978-1-84593-620-4

Tomar, K.S. (1979). Studied on rootage and growth and Karonda (*Carissa carandus* L.) air layers with and without growth regulators. M.Sc. (Ag.) Thesis JNKVV, Jabalpur.

Tyagi, S.K. and R.M. Patel (2004). Effect of growth regulators on rooting of air layering of guava (*Psidium guajava* L.) cv. Sardar Guava. *Orissa. J. Hort.* 32(1): 58-62.

UNDER PEER REVIEW

Table 1.0 Root parameters as effected by different IBA concentrationson air layering in olive.

Treatment	Treatment Combination	Days to root formation	No. of Roots	Root length (cm)	Root Thickness (mm)
T1	IBA@ 1000 ppm	68.88	4.33	4.82	1.23
T2	IBA@ 2000 ppm	87.88	4.88	5.19	1.26
T3	IBA@ 3000 ppm	83.33	5.11	4.56	1.53
T4	IBA@ 4000 ppm	46.11	7.33	6.73	1.66
T5	IBA@ 5000 ppm	62.11	6.44	6.20	1.60
T6	IBA@ 6000 ppm	85.78	6.00	5.15	1.52
T7	IBA@ 7000 ppm	82.89	4.66	4.68	1.53
T8	Control	97.00	3.44	3.19	1.13
	S.E. (m) (±)	7.28	0.22	0.18	0.10
	CD (5%)	22.09	0.68	0.56	0.31

Table 2.0 Shoot parameters and survival percentage as effected by different IBA concentrations on air layering in olive.

Treatment	NUMBER OF NEW SROUTS			LENGTH OF NEW SHOOTS			NUMBER OF NEW LEAVES			SURVIVAL PERCENTAGE		
	1MAP	2MAP	3MAP	1MAP	2MAP	3MAP	1MAP	2MAP	3MAP	1MAP	2MAP	3MAP
T1	1.78	2.33	3.33	1.49	2.09	4.40	5.44	5.44	11.67	41.56	38.44	34.67
T2	1.67	3.33	4.22	1.53	2.23	4.77	6.33	6.78	12.11	41.33	39.33	36.89
T3	1.78	3.22	4.22	1.50	3.27	4.62	6.11	7.11	14.44	45.11	44.11	38.67
T4	2.67	3.89	5.67	2.78	3.73	5.44	7.22	10.78	16.56	57.56	54.44	50.89
T5	2.00	3.44	4.33	1.76	3.41	4.99	6.56	8.44	16.00	53.67	47.11	43.00
T6	1.89	3.11	4.11	1.51	3.34	4.91	6.22	8.11	14.78	44.89	43.22	39.67
T7	1.89	2.44	4.00	1.50	3.32	4.77	5.67	7.56	14.56	43.78	42.00	38.44
T8	1.44	2.00	2.78	1.40	2.06	3.73	5.00	5.89	10.56	35.00	33.78	33.33
S.E. (m) (±)	0.21	0.12	0.16	0.11	0.05	0.05	0.30	0.36	0.30	0.58	0.67	0.41
CD (5%)	0.64	0.37	0.50	0.33	0.14	0.16	0.91	1.08	0.90	0.82	0.95	0.58