

## **Influence of Inorganic Fertilizers in conjunction with Organic Manures on Growth, Yield and Quality of Broccoli (*Brassica oleracea* var. *Italica*)**

---

### **ABSTRACT**

A research work was conducted in the research farm of Nalanda College of Horticulture, Noorsarai, Nalanda (Bihar-803113) during Rabi season of academic year 2021-22 to study the Influence of Inorganic Fertilizers in conjunction with organic manures on growth, yield and Quality of Broccoli (*Brassica oleracea* var. *Italica*) in a randomized block design with 11 treatments. The Treatment, T<sub>5</sub> (50% RDN + 50% N as PM) recorded maximum values for plant height (37.41cm and 56.13cm), number of leaves per plant (16.53 and 20.11), leaf area (749.51cm<sup>2</sup> and 949.57 cm<sup>2</sup>) at 45 DAT and 60DAT respectively. The treatment, T<sub>5</sub> also recorded least days for head initiation (44.60 DAT) and highest value for diameter of head (15.93cm), length of head (14.97cm), wt. of head/plant (407.16gm), wt. of head/plot (16.29Kg.), wt. of head/ha (150.65q.), vitamin C (92.43mg/g), dry matter (11.57%), gross returns (Rs. 4,14,287.50/ha) and net returns (Rs. 3,19,448.00/ha) with B:C ratio of 3.36. This was followed by treatment T<sub>4</sub> (50% RDN + 50% N as VC). Among all the treatments, the Treatment T<sub>5</sub> (50% RDN + 50% N as PM) was found to be the best for obtaining better growth and optimum yield. Regarding qualitative characters and better soil health, treatment T<sub>4</sub> (50% RDN + 50% N as VC) performed well.

**Keywords:** Broccoli, FYM, VC, PM, Yield, Plant growth, quality, gross returns.

### **Introduction**

Broccoli (*Brassica oleracea* L. var. *Italica*) is a winter/cool season vegetable and belongs to family Cruciferae and commonly known as harigobi in hindi because. It is either consumed raw as a salad, cooked as a single or mixed vegetable with potato, and used in curries, soups, and pickles. There are two types of broccoli: heading and sprouting, sprouting broccoli is more widely consumed in India. Broccoli has anti-carcinogenic properties and reduce the risk of prostate cancer by up to 45 percent. It has about 130 times more Vitamin A contents than cauliflower and 22 times more than cabbage. Broccoli is also known as the “Crown of Jewel Nutrition” due to a rich source of vitamins, minerals, proteins etc. (Thamburaj and Singh, 2010). In the global market approximately 40 percent is marketed as fresh and remaining 60 percent

marketed as frozen. The exact data about the cultivable area and production of broccoli is currently unavailable in India. It was a rare cole crop in India, but it is now becoming more common in metropolitan's cities, hotels, and restaurants (Thamburaj and Singh, 2010).

Many chemical fertilizers and just a little amount of organic fertilizers are used by various workers to enhance the production, which eventually has an impact on both human and soil health (Meena *et al.*, 2017). Integrated nutrient management is an essential need of the modern era since excessive use of chemical fertilizer has resulted in a pattern of declining soil fertility (Attigah *et al.*, 2013). The basic goal of integrated nutrient management is to cultivate a piece of land so that the soil can support optimum crop quality production while remaining sustainable (Mishra *et al.*, 2014).

Organic manure plays a direct role in plant growth, as a supply of all essential macro and micronutrients in forms that are available during mineralization and increasing physical and chemical qualities of soils (Chatterjee *et al.*, 2015). As a result, using locally produced manures in vegetable production operations may boost crop yield while requiring less chemical fertilizer. Consumers now expect veggies to be of higher safety and superior quality.

Hence, keeping in view of aforementioned facts, a research work entitled "Influence of Inorganic Fertilizers in conjunction with Organic Manures on Growth, Yield and Quality of Broccoli (*Brassica oleracea* var. *Italica*)" was conducted to determine the best inorganic fertilizer and organic manure combination for maximum growth, yield, quality and net profit in broccoli and for better soil health.

### **Materials and methods**

Present research work was conducted during Rabi season (2021-22) at research farm of Nalanda College of Horticulture, Noorsarai (Nalanda) with following objectives: (i) To find out the effect of inorganic fertilizers and organic manures on growth, yield and quality of broccoli, (ii) To find out nitrogen content and nitrogen use efficiency of broccoli, (iii) To find out effect of manures and fertilizers on properties of soil and (iv) To find out the economics. The soil condition of the research field was low in available nitrogen, medium in available phosphorus, potash and organic carbon while, good in pH and electrical conductivity. A randomized block

design with three replications was used to apply eleven various treatments, including the use of organic manures (FYM, Vermicompost, and poultry manure) and inorganic fertilizers (Urea, SSP, and MOP). The treatments included T<sub>1</sub> (Control), T<sub>2</sub> (RDN= 150 Kg/Ha), T<sub>3</sub> (50% RDN + 50% N as FYM), T<sub>4</sub> (50% RDN + 50% N as VC), T<sub>5</sub> (50% RDN + 50% N as PM), T<sub>6</sub> (25% RDN + 75% N as FYM), T<sub>7</sub> (25% RDN + 75% N as VC), T<sub>8</sub> (25% RDN + 75% N as PM), T<sub>9</sub> (25% RDN + 25% N as VC + 50% N as FYM), T<sub>10</sub> (25% RDN + 25% N as VC + 50% N as PM), T<sub>11</sub> (25% RDN + 25% N as VC + 25% N as PM + 25% as FYM). The crop was raised with a spacing of 60 cm × 45 cm and plot size of 4.5 m × 2.4 m. All of the experimental plots were maintained according to the standard cultural guidelines for broccoli. One month prior to transplanting (or during field preparation), organic manures (FYM, Vermicompost and Poultry Manure) were applied as per the requirement of treatments and mixed well with a spade. Inorganic fertilizer i.e., nitrogenous fertilizer was applied as per the selected treatments in which half dose of nitrogen through urea and common dosage of P<sub>2</sub>O<sub>5</sub> and K<sub>2</sub>O @ 80 kg/ha each through SSP and MOP respectively during transplanting. The remaining half dose of nitrogen was given 30 DAT as per requirement of treatments. The observations like various growth parameters yield and yield attributing characteristics, qualitative characteristics and nutrient status of post harvested soil were noted. On the basis of current market pricing for inputs and outputs, the economics of the experiment were calculated.

## Results and Discussion

The purpose of the present research was to understand how integrated nutrient management affected broccoli. Observations on the characteristics of the plants were made to explore the impact of chemical fertilizers and various organic manures on growth parameters, yield parameters, qualitative features, soil characteristics and economics. Inorganic fertilizers alone or in combination with different organic manures were found to have a significant effect on growth parameters as compared to control (Table 1). The data was recorded on plant height (cm), number of leaves plant<sup>-1</sup>, leaf area per plant (cm<sup>2</sup>) of broccoli at 45 DAT and 60 DAT. The application of treatment T<sub>5</sub> (50% RDN + 50% N as PM) showed maximum plant height (37.41 cm and 56.13 cm, respectively), number of leaves plant<sup>-1</sup> (916.53 and 20.11, respectively) and leaf area (749.51cm<sup>2</sup> and 949.57 cm<sup>2</sup>, respectively). The minimum values of growth characters were recorded in treatment T<sub>1</sub> (control). This variation might be caused by nutrient availability,

the fact that organic manure improves soil aggregation, aeration, and water holding capacity while inorganic fertilizers provide more readily available forms of nutrients. As a result, mixing organic and inorganic nutrients gives superior results in better nutrient absorption and utilization by plants leading to better plant growth. Wani and Ahmed (2011), Chetri *et al.* (2012) and Mohanta *et al.* (2018) all provided confirmation of the findings.

Nutrient application of inorganic fertilizers and organic manures alone or in combination significantly increased yield and yield attributing characters of broccoli as compared to control (Table 2). The application of treatment T<sub>5</sub> (50% RDN + 50% N as PM) was recorded minimum days for head initiation (44.60) and maximum values for diameter of head (15.93 cm), Length of head (14.97cm), Wt. of head/plant (407.16 gm), Wt. of head/Plot (16.29 Kg.) and Wt. of head/ha (150.65 q.). This might be as a result of the beneficial role of organic manures in improving physical, chemical and biological properties of soil, which result in better nutrient absorption by plants, which might account for increased carbohydrate accumulation due to increased photosynthesis and resulted in higher values for yield contributing characters. These results conform with the finding of Kumar *et al.* (2011) in cauliflower, Chatterjee *et al.* (2014) in cabbage, and Atal *et al.* (2019) in broccoli. Maximum head weight in various combinations of organic and inorganic fertilizers observed by Choudhary *et al.* (2018) in broccoli

**Table- 1. Growth parameters of broccoli**

Treatments	Plant Height at 45 DAT (cm)	Plant Height at 60 DAT (cm)	No. of leaves at 45 DAT	No. of leaves at 60 DAT	Leaf area at 45 DAT (Cm <sup>2</sup> )	Leaf area at 60 DAT (Cm <sup>2</sup> )
T <sub>1</sub> (Control)	22.63	32.70	9.33	14.30	388.72	597.10
T <sub>2</sub> (RDN = 150 Kg/Ha)	30.23	45.37	14.28	17.97	634.44	867.33
T <sub>3</sub> (50% RDN + 50% N as FYM)	35.34	49.90	14.43	18.73	691.10	907.70
T <sub>4</sub> (50% RDN + 50% N as VC)	35.80	52.37	14.77	19.37	725.40	919.83
T <sub>5</sub> (50% RDN + 50% N as PM)	37.41	56.13	16.53	20.11	749.51	949.57
T <sub>6</sub> (25% RDN + 75% N as FYM)	30.29	45.37	13.66	17.57	579.60	808.07
T <sub>7</sub> (25% RDN + 75% N as VC)	32.19	46.60	14.30	18.32	605.57	765.07
T <sub>8</sub> (25% RDN + 75% N as PM)	32.83	47.40	13.20	17.03	523.74	739.83

<b>T<sub>9</sub></b> (25% RDN + 25% N as VC + 50% N as FYM)	28.73	44.90	12.97	17.01	517.47	733.62
<b>T<sub>10</sub></b> (25% RDN + 25% N as VC + 50% N as PM)	26.40	43.60	12.47	16.17	474.47	714.17
<b>T<sub>11</sub></b> (25% RDN + 25% N as VC + 25% N as PM + 25% as FYM)	27.95	44.03	12.63	16.33	487.57	735.97
<b>Grand Mean</b>	<b>30.89</b>	<b>46.22</b>	<b>13.51</b>	<b>17.54</b>	<b>558.84</b>	<b>794.39</b>
<b>S.Em±</b>	<b>1.69</b>	<b>1.99</b>	<b>0.41</b>	<b>0.78</b>	<b>70.64</b>	<b>19.85</b>
<b>CD (5%)</b>	<b>4.87</b>	<b>5.74</b>	<b>1.18</b>	<b>2.23</b>	<b>203.48</b>	<b>57.19</b>

**Table- 2. Yield parameters of broccoli**

<b>Treatments</b>	<b>Days taken to head initiation (DAT)</b>	<b>Diameter of Head (cm)</b>	<b>Length of head (cm)</b>	<b>Wt. of Head/Plant (gm)</b>	<b>Wt. of Head/Plot (Kg.)</b>	<b>Wt. of Head/Ha (q.)</b>
<b>T<sub>1</sub></b> (Control)	48.70	8.70	8.07	167.03	6.68	61.80
<b>T<sub>2</sub></b> (RDN = 150 Kg/Ha)	47.00	14.10	13.33	353.58	14.14	130.82
<b>T<sub>3</sub></b> (50% RDN + 50% N as FYM)	45.83	14.83	13.60	375.83	15.03	139.06
<b>T<sub>4</sub></b> (50% RDN + 50% N as VC)	45.37	15.60	14.23	394.57	15.78	145.99
<b>T<sub>5</sub></b> (50% RDN + 50% N as PM)	44.60	15.93	14.97	407.16	16.29	150.65
<b>T<sub>6</sub></b> (25% RDN + 75% N as FYM)	46.73	14.17	12.57	337.72	13.51	124.96
<b>T<sub>7</sub></b> (25% RDN + 75% N as VC)	46.93	14.67	13.30	328.99	13.16	121.73
<b>T<sub>8</sub></b> (25% RDN + 75% N as PM)	46.40	14.78	12.43	313.17	12.53	115.87
<b>T<sub>9</sub></b> (25% RDN + 25% N as VC + 50% N as FYM)	47.17	13.87	12.10	305.08	12.20	112.88
<b>T<sub>10</sub></b> (25% RDN + 25% N as VC + 50% N as PM)	47.43	13.27	10.33	277.58	11.10	102.70
<b>T<sub>11</sub></b> (25% RDN + 25% N as VC + 25% N as PM + 25% as FYM)	47.30	13.83	10.90	288.57	11.54	106.77
<b>Grand Mean</b>	<b>46.68</b>	<b>13.98</b>	<b>13.26</b>	<b>322.66</b>	<b>12.91</b>	<b>119.38</b>
<b>S.Em±</b>	<b>0.33</b>	<b>0.23</b>	<b>0.42</b>	<b>6.68</b>	<b>0.27</b>	<b>2.47</b>
<b>CD (5%)</b>	<b>0.96</b>	<b>0.68</b>	<b>1.21</b>	<b>19.25</b>	<b>0.27</b>	<b>7.12</b>

The results obtained during the investigation in respect to quality parameters viz., Dry matter percentage, T.S.S vitamin C content and nitrogen percentage in head present in Table 3. Different treatment organic manure and inorganic fertilizer was found to influence significantly. Treatments T<sub>5</sub> (50% RDN + 50% N as PM) recorded highest value (11.56<sup>o</sup>B) for dry matter percentage followed by treatment T<sub>4</sub> (50% RDN + 50% N as VC). While, the maximum T.S.S.

(12.1<sup>o</sup>B) is found in treatment T<sub>4</sub> (50% RDN + 50% N as VC) and which is followed by treatments T<sub>5</sub> (50% RDN + 50% N as PM) with (11.96<sup>o</sup>B).

**Table- 3. Quality parameters of broccoli**

Treatments	Dry matter % of head	TSS ( <sup>o</sup> B )	Vit. C content (mg/100 gm)	Nitrogen % in head
T <sub>1</sub> (Control)	7.43	8.36	65.31	3.08
T <sub>2</sub> (RDN = 150 Kg/Ha)	9.78	11.16	81.25	3.44
T <sub>3</sub> (50% RDN + 50% N as FYM)	10.26	11.3	86.61	3.74
T <sub>4</sub> (50% RDN + 50% N as VC)	11.07	12.1	92.42	3.82
T <sub>5</sub> (50% RDN + 50% N as PM)	11.56	11.96	89.21	3.75
T <sub>6</sub> (25% RDN + 75% N as FYM)	10.6	10.93	85.53	3.59
T <sub>7</sub> (25% RDN + 75% N as VC)	8.46	11.53	85.13	3.72
T <sub>8</sub> (25% RDN + 75% N as PM)	7.63	11.03	81.06	3.66
T <sub>9</sub> (25% RDN + 25% N as VC + 50% N as FYM)	9.35	10.7	80.35	3.60
T <sub>10</sub> (25% RDN + 25% N as VC + 50% N as PM)	8.16	10.4	73.53	3.57
T <sub>11</sub> (25% RDN + 25% N as VC + 25% N as PM + 25% as FYM)	8.17	10.63	79.76	3.41
<b>Grand mean</b>	<b>9.32</b>	<b>10.92</b>	<b>81.84</b>	<b>3.58</b>
<b>S.Em±</b>	<b>0.13</b>	<b>0.12</b>	<b>0.77</b>	<b>0.05</b>
<b>CD (5%)</b>	<b>0.36</b>	<b>0.35</b>	<b>2.23</b>	<b>0.16</b>

The maximum vitamin-C content (92.42 mg/100g) was recorded under treatment T<sub>4</sub> (50% RDN + 50% N as VC) followed by T<sub>5</sub> (50% RDN + 50% N as PM) with 89.21 mg/100g. Meaning that the combined application of various fertilizer and manure played a major contribution to the higher total soluble solids in the broccoli heads. Similar observations were also noted by Upadhyay *et al.* (2012) in cabbage, Kumar *et al.* (2013) in broccoli and Chatterjee *et al.* (2015) in broccoli. The maximum nitrogen percentage (3.82%) in head was recorded in treatment T<sub>4</sub> (50% RDN + 50% N as VC) followed by T<sub>5</sub> (50% RDN + 50% N as PM) (3.75%). However minimum value for these qualitative parameters was recorded in treatment T<sub>1</sub> (Control). Similar findings were made by Ouda and Mahadeen (2008), who found that the control treatment resulted in the lowest leaf N content (3.18%), whereas the maximum leaf N content (3.87%) was obtained when both inorganic fertilizer and organic manure was applied in appropriate dosages. Findings of Shalini *et al.* (2002) in knol-khol, Sharma *et al.* (2008) in sprouting broccoli and Mohanta *et al.* (2018) also came to same conclusion. Effect on soil property after harvest

presented in Table- 4. Treatment T<sub>5</sub> (50% RDN + 50% N as PM) recorded the highest amount of available nitrogen (220.4 kg/ha) in soil after harvest, which was significantly higher than treatment T<sub>1</sub> (Control). Sharma *et al.* (2008) claimed that the reason for the increase in nitrogen concentration in soil is that organic manures and chemical fertilizers increase the direct supply of nitrogen and promote the growth of microorganisms, which may be related to the conversion of organically bound nitrogen to inorganic and usable forms. Kumar *et al.* (2017) observed that availability of NPK in soil was increased with **integrated application of organic manures and synthetic fertilizers. They also noted that the addition** of organic manure reduced the C:N ratio, which further sped up mineralization and led to an immediate release of nitrogen into the soil. Sreenivasa *et al.* (2010) and Devakumar *et al.* (2011) came to same conclusion.

Maximum phosphorus (18.25 kg ha<sup>-1</sup>) and potash (280.36 kg ha<sup>-1</sup>) availability in soil was found in treatment T<sub>4</sub> (50% RDN + 50% N as VC) whereas minimum available phosphorus and potash was noted in treatment T<sub>1</sub> (Control). Pawar *et al.* (2018) noted that the causes of the increased availability of phosphorus in soil might be due to the decomposition of organic matter and the solubilization of phosphorus from **the native soil pool. Similar observations were recorded by Narayanamma *et al.* (2005) and Sarangthem *et al.* (2011). The probable reason for high potash availability is due to the improvement of** potassium (K) status in soil accounts to reduced fixation rate and release of K as a result of the interaction of organic sources with clay particles. The above outcomes are corresponding with the findings of Kumar P *et al.* (2017) and Das *et al.* (2019) in sprouting broccoli.

**Table- 4. Effect on soil property after harvesting of crop**

Treatments	N (kg/ha)	P (kg/ha)	K (kg/ha)	pH	EC (dS m <sup>-1</sup> )	OC (%)
T <sub>1</sub> (Control)	180.35	13.12	232.45	6.83	0.35	0.64
T <sub>2</sub> (RDN = 150 Kg/Ha)	211.6	16.84	272.62	6.8	0.37	0.66
T <sub>3</sub> (50% RDN + 50% N as FYM)	218.04	17.35	274.23	7.05	0.38	0.74
T <sub>4</sub> (50% RDN + 50% N as VC)	218.13	18.25	280.36	6.84	0.4	0.75
T <sub>5</sub> (50% RDN + 50% N as PM)	220.4	18.15	277.22	7.05	0.38	0.78
T <sub>6</sub> (25% RDN + 75% N as FYM)	215.75	17.56	274.82	7.08	0.37	0.69
T <sub>7</sub> (25% RDN + 75% N as VC)	217.8667	18.05	276.65	7.06	0.39	0.7
T <sub>8</sub> (25% RDN + 75% N as PM)	214.4833	17.92	275.72	6.97	0.36	0.72

<b>T<sub>9</sub></b> (25% RDN + 25% N as VC + 50% N as FYM)	214.66	17.22	273.66	7.11	0.35	0.67
<b>T<sub>10</sub></b> (25% RDN + 25% N as VC + 50% N as PM)	212.88	16.62	274.47	6.87	0.37	0.71
<b>T<sub>11</sub></b> (25% RDN + 25% N as VC + 25% N as PM + 25% as FYM)	213.55	16.51	275.41	7.09	0.38	0.68
<b>Grand Mean</b>	<b>212.52</b>	<b>17.05</b>	<b>271.60</b>	<b>6.97</b>	<b>0.37</b>	<b>0.70</b>
<b>S.Em±</b>	<b>0.52</b>	<b>0.17</b>	<b>1.34</b>	<b>0.04</b>	<b>0.01</b>	<b>0.02</b>
<b>CD (5%)</b>	<b>1.50</b>	<b>0.51</b>	<b>3.86</b>	<b>0.12</b>	<b>0.04</b>	<b>0.06</b>

The data pertaining to pH, electrical conductivity and organic carbon in Table-4 clearly indicates that the difference between the treatments was found to be non-significant. After termination of research work, highest pH (7.11) was recorded in T<sub>9</sub> (25% RDN + 25% N as VC + 50% N as FYM) whereas, minimum (6.83) was recorded in T<sub>1</sub> (Control). Similar result was observed by Ouda and Mahadeen (2008). They also pointed that application of organic, inorganic manures and their interaction had no significant effect on soil pH. There was minute increase EC value in treatment T<sub>4</sub> (50% RDN + 50% N as VC) over rest of the treatments and it might be due to large amounts of soluble salts and HCO<sup>3-</sup> contained in organic manures may be the cause of the small rise in EC value. Singh *et al.* (2017) noticed that the use of organic manures may enhance the amount of organic carbon and organic matter in vegetable crops because less fossil fuel energy is consumed, which results in lower emissions of carbon dioxide and nitrous dioxide, which raise soil carbon concentration.

**Table- 5. Economics of treatments**

<b>Treatments</b>	<b>Treatment costs Rs./ha</b>	<b>Fixed input Rs./ha</b>	<b>Total input costs (Rs./ha)</b>	<b>Yield (q/ha)</b>	<b>Gross income (Rs.)</b>	<b>Net income (Rs.)</b>	<b>B:C</b>
<b>T<sub>1</sub></b> (Control)	0	68,210	68,210	61.79	1,69,922.5	1,01,713	1.49
<b>T<sub>2</sub></b> (RDN = 150 Kg/Ha)	10740	68,210	78,950	130.82	3,59,755	2,80,805	3.56
<b>T<sub>3</sub></b> (50% RDN + 50% N as FYM)	45245	68,210	1,13,455	139	3,82,250	2,68,795	2.37
<b>T<sub>4</sub></b> (50% RDN + 50% N as VC)	42345	68,210	1,10,555	145.9	4,01,225	2,90,670	2.63
<b>T<sub>5</sub></b> (50% RDN + 50% N as PM)	26630	68,210	94,840	150.65	4,14,287.5	3,19,448	3.36

T <sub>6</sub> (25% RDN + 75% N as FYM)	62497	68,210	1,30,707	124.95	3,43,612.5	2,12,906	1.62
T <sub>7</sub> (25% RDN + 75% N as VC)	58147	68,210	1,26,357	121.73	3,34,757.5	2,08,401	1.65
T <sub>8</sub> (25% RDN + 75% N as PM)	34575	68,210	1,02,785	115.87	3,18,642.5	2,15,858	2.10
T <sub>9</sub> (25% RDN + 25% N as VC + 50% N as FYM)	61047	68,210	1,29,257	112.88	3,10,420	1,81,163	1.40
T <sub>10</sub> (25% RDN + 25% N as VC + 50% N as PM)	42432	68,210	1,10,642	102.8	2,82,700	1,72,058	1.55
T <sub>11</sub> (25% RDN + 25% N as VC + 25% N as PM + 25% as FYM)	51740	68,210	1,19,950	110.7	3,04,425	1,84,475	1.53

The economics of different treatments viz., yield (t/ha), cost of cultivation, gross return and benefit cost ratio has been worked out and presented in Table-5. Treatment T<sub>5</sub> (50% RDN + 50% N as PM) recorded maximum gross return of Rs, 4,14,287 ha<sup>-1</sup> and maximum net return Rs. 3,19,448 ha<sup>-1</sup> followed by treatment T<sub>4</sub> (50% RDN + 50% N as VC) with Rs, 4,01,225 ha<sup>-1</sup> and Rs. 2,90,670 ha<sup>-1</sup>, respectively whereas the cost benefit ratio of treatment T<sub>2</sub> (RDN= 150 Kg/ha) i.e. (3.56) was found to be the best treatment combinations terms of economics returns.

## Conclusion

Based upon the results recorded during research, it could be concluded that the treatment T<sub>5</sub> (50% RDN + 50% N as PM) was shown to be the best treatment with regard to growth, yield, and yield attributing features in broccoli as compared to control and other treatments followed by treatment T<sub>4</sub> (50% RDN + 50% N as VC). Regarding qualitative characters and for better soil health, treatment T<sub>4</sub> (50% RDN + 50% N as VC) performed well. The noteworthy results of economic return with B:C ratio of 3.36 was also given with an application of treatment T<sub>5</sub> (50 percent RDN + 50 percent N as PM) followed by treatment T<sub>4</sub> (50% RDN + 50% N as VC).

## REFERENCES

Atal MK, Dwivedi DH, Narolia SL, Bharty N and Kumari R (2019) Influence of bio-fertilizer (*Rhizobium radiobacter*) in association with organic manures on growth and yield of broccoli

(*Brassica oleracea* L. var. *italica* Plenck) cv. Palam Samridhi under Lucknow conditions. *Journal of Pharmacognosy and Phytochemistry*, 1: 604-608.

Attigah AS, Asiedu EK, Agyarko K and Dapaah HK (2013) Growth and yield of okra (*Abelmoschus esculentus* L.) as affected by organic and inorganic fertilizers. *ARPJ: J. Agri. Biol. Sci.*, 8(12): 766-770.

Chatterjee R, Dipika Mal and Nimbalkar KH (2015) Effect of vermincompost and inorganic fertilizers on growth, yield and quality of sprouting broccoli (*Brassica oleracea* L. var. *italica* Plenck). *Int. J. of Bio. Res. and Stre. Mang*, 5(4): 507-512.

Chetri DA, Singh AK and Singh VB (2012) Effect of integrated nutrient management on yield, quality and nutrient uptake by capsicum (*Capsicum annum*) cv. California wonder. *Journal of Soils and Crops*, 22(1): 44-48.

Choudhary K, Dev P, Kumar J, Kumar V and Kumar T (2018) Effect of integrated nutrient management on yield parameters of broccoli (*Brassica oleracea* L. var. *italica*) cv. Pusa KTS-1. *International Journal of Agricultural Invention*, 3:223-26.

Dash SK, Sahu GS, Das S, Sarkar S and Pathak M (2019) Effect of integrated nutrient management on yield, yield attributes and economics of broccoli. *International Journal of Current Microbiology and Applied Sciences*, 8:3254-58.

Devakumar N, Rao GGE and Shuba S (2011) Evaluation of locally available media for the growth and development of nitrogen fixing micro-organisms. Proceedings of the 3rd scientific conference of ISOFAR Organic are life- knowledge for tomorrow. 28 September - 01 October, 2011, Korea. pp. 504-09.

Devi HJ, Maity and Paria NC (2003) Effect of different sources of nitrogen on yield and economics of cabbage. *Environ. Ecol.*, 21(4): 878-880.

Kumar M, Das B, Prasad KK and Kumar P (2013) Effect of integrated nutrient management on growth and yield of broccoli (*Brassica oleracea* var. *italic*) under Jharkhand condition. 40(1): 117-120.

- Kumar P, Bhardwaj ML, Kumar D, Kumar R, Tripathi D, Thakur KS, Bharat N, Gautam N, Kumar S and Dogra B (2017) Comparative performance of organic and inorganic fertilizers on plant growth, head yield, soil health and severity of black rot in sprouting broccoli cv. Green Head. *International Journal of Farm Sciences*, 7:69-76.
- Kumar S, Verma MK and Yadav Y C (2011) Studies on effect of biofertilizers with chemical fertilizers on growth and yield of cauliflower (*Brassica oleracea* var. *botrytis*) cv. Pusa Snowball K-1. *Annals of Horticulture*, 4(2): 202-205.
- Meena VK, Dubey AK, Jain VK, Tiwari A and Negi P (2017) Effect of plant growth promoters on flowering and fruiting attributes of okra [*Abelmoschus esculentus* (L.) Moench]. *Crop Res.*, 52(1, 2 & 3): 37-40.
- Mishra PP, Das AK and Mishra N (2014) Effect of integrated nutrient management on yield, quality and economics of knolkhol (*Brassica oleracea* L. cv. *gongylodes*). *Asian J. Horticulture*, 9(2): 382-385.
- Mohanta R, Nandi AK, Mishra SP, Pattnaik A, Hossain MM and Padhiary AK (2018) Effects of integrated nutrient management on growth, yield, quality and economics of sprouting broccoli (*Brassica oleracea* var. *italica*) cv. Shayali. *Journal of Pharmacognosy and Phytochemistry*, 7(1): 2229-2232.
- Narayanamma M, Chiranjeevi CH, Reddy IP and Ahmed SR (2005) Integrated nutrient management in cauliflower (*Brassica oleracea* var. *botrytis* L.). *Vegetable Science*, 32:62- 64.
- Ouda BA, and Mahadeen AY (2008) Effect of fertilizers on growth, yield, yield components, quality and certain nutrient contents in broccoli (*Brassica oleracea*). *Int. J. Agri. Biol.*, 10 (3): 627–32.
- Pawar RU, Barkule S, Kirti S and Rasal D (2018) Effect on soil health of cauliflower (*Brassica oleracea* var. *botrytis*) cultivation with integrated nutrient management. *Journal of Applied and Natural Science*, 10:1026-31.
- Sarangthem I, Misra ADD and Chakraborty Y (2011) Cabbage productivity, nutrient uptake and soil fertility as affected by organic and biosources. *Agricultural Science Digest*, 31:260.

- Shalini SB, Channal HT, Hebsur NS, Dharnatti PR and Sarangamath PA (2002) Effect of integrated nitrogen management on yield of knol-khol and population of *Azospirillum* in vertisols of Karnataka. *Journal of Agricultural Sciences*, 15(3):151-153.
- Sharma A, Parmar DK, Kumar P, Singh Y and Sharma RP (2008) Azotobacter soil amendment integrated with cow manures reduces need for NPK fertilizers in Sprouting Broccoli, *International Journal of Vegetable Science*, 14(3): 273- 285.
- Shiyam JO and Binang WB (2010) Effect of poultry manure and urea on flowering occurrence and leaf productivity of *Amaranthus cruentus*. *Journal of Applied Sciences and Environmental Management*, 15(1): 13-15.
- Singh V, Shah N and Rana DK (2017). Combined Effect of Organic Manures and Bio-fertilizers on growth and yield of broccoli under Garhwal Himalayan Region. *Hort Flora Research Spectrum*, 5(4): 345-347.
- Sreenivasa MN, Nagaraj MN and Bhat SN (2010) Beejamruth: A source for beneficial bacteria. *Karnataka Journal of Agricultural Sciences*, 17:72-77.
- Swarup, V. 2012. Vegetable science and technology in India. Kalayani Publishers, New Delhi, pp 370-371.
- Thamburaj S and Singh Narendra (2010) Textbook of Vegetables, Tubercrops and Spices. Indian Council Agricultural Research. 137p.
- Upadhyay RK, Patra DD, Tewari SK and Baksh H (2012) Effect of farm yard manure, vermicompost, chemical fertilizer and integrated weed management on yield, yield contributing characters and quality of mint (*Mentha arvensis* L.) *Crop Research Hisar*, 44(3): 344-348.
- Wani ABJ, Raj Narayan, Ahmed N, Singh AK, Chattoo MA and Sumati Narayan (2011) Influence of organic and inorganic sources of nutrients on growth, yield and quality of cauliflower (*Brassica oleracea* L. var. botrytis). *Environment and Ecology*, 29(4A): 1941-1947.