

Characterization and functional properties of lactic acid bacteria isolated from fermented maize dough used for the preparation of *doklu*

ABSTRACT

Lactic acid bacteria (LAB) have long been safely applied in the production of fermented foods and beverages. These bacteria are important for their contribution to food quality and also for their undeniable effects on health. In Africa, despite the abundance of isolated food sources and the diversity of identified microorganisms, the market for probiotics and functional foods based on selected local strains is almost non-existent. This paper aims to determine some technological properties of LAB isolates from fermented maize dough used for *doklu* production for potential application in food and probiotic products. A total of 25 LAB strains isolated from maize dough fermentation processes during *doklu* production were identified by 16S rRNA gene sequencing. Technological studies such as growth and acidification kinetics, diacetyl production, hydrogen peroxide production and proteolytic activity were carried out. They survival and growth under unfavorable conditions of temperature, salt concentration and pH were also analyzed. The isolates were identified as *Pediococcus acidilactici* (56%), *Lactobacillus fermentum* (24%), *Pediococcus pentosaceus* (12%) and *Lactobacillus plantarum* (8%). The technological properties of the LAB strains showed that 10 of them rapidly acidify the medium with pH variation (Δ pH) greater than 1 after 6 hours of fermentation. Good production of diacetyl and hydrogen peroxide as well as a good proteolytic activity were obtained for several strains. *Pediococcus acidilactici* S15, S55, S56; *Lactobacillus plantarum* S32, S121 and *Lactobacillus fermentum* S44 showed best technological properties. However, only 5 of them obtained good resistance following the various survival and growth under unfavorable conditions of temperature, salt concentration and pH. This work has therefore demonstrated that the lactic acid bacteria involved in the natural fermentation of maize dough have interesting properties for various applications whether in the processing of food products as well as in the development of probiotic products.

Keywords: *Doklu*, Lactic acid bacteria (LAB), maize dough, *Pediococcus acidilactici*, Technological properties

1. INTRODUCTION

Foods derived from cereals are popular worldwide and differ depending on the local culture and traditions [1]. They serve as important components of the daily diet, providing carbohydrates, proteins, dietary fibers and vitamins [2]. *Doklu*, one of these cereal based foods, is a product obtained from fermented corn dough consumed by people in southern and South-Eastern of Côte d'Ivoire. The people often appreciate *doklu* for its sour taste [3]. The manufacturing process of this product includes a fermentation step performed by a wide range of microorganisms, predominated by lactic acid bacteria (LAB). LAB involved in this spontaneous fermentation were mainly identified as *Lactobacillus fermentum*, *Pediococcus acidilactici*, *Lactobacillus plantarum*, *Pediococcus pentosaceus*, and *Weissella cibaria* [4].

Studies on the technological and safety properties of LAB from different origins evidence that such microbial group has appropriate characteristics for many applications [5]. LAB has long been safely applied in the production of fermented foods and beverages. They occupy a central role in traditional starter cultures and are known to enhance the shelf life,

and improve microbial safety, texture and sensory profile of the fermented product, mainly through the production of organic acids, aroma compounds, bacteriocins, exopolysaccharides and several enzymes [6]. LAB are also common components of probiotics, due to the fact that they are « generally regarded as safe » as they have long been used in the manufacture of foods and are desirable members of the intestinal microflora [7].

In Africa, despite the abundance of isolated food sources and the diversity of identified microorganisms, the market for probiotics and functional foods based on selected local strains is almost non-existent. Isolation of original LAB from traditional fermented foods is of crucial interest in view of there is a paucity of literature regarding novel and emerging uses of LAB as probiotics from most of developing countries. However, for application of these LAB strains as dietary adjuncts, in addition to screening for beneficial properties several factors which may influence the survival and colonization of these bacteria need to be addressed [8]. This paper aims to determine some technological properties of LAB isolates from fermented maize dough used for *doklu* production. We reported the isolation, characterization and identification of novel LAB strains. Also, the evaluation of some physiological and technological properties and the selection of some potential probiotic candidates.

2. MATERIAL AND METHODS

2.1 LAB isolation and identification

2.1.1 Sample collection and isolation

Fermented maize dough samples were collected just at the end of the fermentation in sterile containers from a traditional producer at Abidjan (Côte d'Ivoire). Immediately after collection, samples were transferred in icebox to the laboratory within 1 h for analyses. Lactic Acid Bacteria isolation was performed after dilution of 10 g of maize dough diluted with 90 mL of sterile buffered peptone water (BPW). Successive decimal dilutions were then prepared, and spread out on Man Rogosa and Sharpe (MRS) agar (Biokar, France). Plates were incubated under microaerobic condition at 37°C for 48 h. Subsequently, colonies were randomly picked from the countable plate from each sample, purified by successive streaking on the same medium and tested for Gram coloration and catalase production. Gram-positive and catalase-negative isolates were stored at -80°C in MRS medium containing glycerol (25%).

2.1.2 DNA extraction

Two ml of overnight bacterial culture were used for DNA extraction. Bacterial cells were pelleted by centrifugation for 10 min at 6000 rpm and protoplast by 1 h incubation at 37°C in 300 µL of 50 mM Tris·HCl (50 mM) EDTA (5mM) at pH 7.5 containing lysozyme (10 mg/mL), Mutanolysin (60 U/mL) and RNase (200µg/mL). The total DNA was then extracted as previously described [9].

2.1.3 PCR Conditions

The V1-V3 region of the 16S rRNA gene were amplified using extracted DNA from isolated strains. PCR amplifications were performed in a final volume of 50 µL, containing 10% of PCR buffer; 2.5 mM of dNTP, 10 pM of each primer (V1-AGTCAGTCAGCCGAGTTTGATCMTGGCTCAG and V3-TATGGTAATTCAATTACCGCGGCTGCTGG), 5 U/µL of Taq polymerase and 500 ng/µL of DNA extract. Samples were amplified in a thermal cycler (GeneAmp PCR System 9700) programmed as follows: initial denaturation of DNA for 1 min at 94°C; 30 cycles (each of 1 min at 94°C, 1 min at 65°C and 1 min at 72°C) and a final extension 10 min at 72°C. PCR products were then separated by electrophoresis on a 1 % (w/v) agarose gel.

2.1.4 Sequencing

The amplified fragments were sequenced by Eurofins Genomics (Ebersberg, Germany) and taxonomic analysis was performed using Blast NCBI database (https://blast.ncbi.nlm.nih.gov/Blast.cgi?PAGE_TYPE=BlastSearch).

2.1 Technological characterization

2.2.1 LAB strain preparation

For analysis, the strains were sub-cultured on MRS agar plate and incubated at 37°C for 48 h under microaerobic conditions. Overnight (12h, 37°C) bacterial cultures were harvested by centrifugation at 4000 rpm for 10 min and washed with sterile BPW. The obtained pellets were re-suspended in BPW at a final concentration of approximately 10⁹ CFU mL⁻¹ [10].

2.2.2 Preparation of maize extract (MFE) broth

Maize flour extract broth was prepared as follows: 500 g of maize grain was washed, steeped in 5 L of water for 12 h and germinated at 25°C for three days. After removal of sprouts, the grains were milled into powder. The maize flour obtained was suspended in distilled water (300 g in 1 L) and boiled for 15 min. The obtained liquid phase was centrifuged at 4000 rpm for 10 min at 4°C. The obtained solution was sterilized by autoclaving at 121°C for 15 min and used as culture broth in subsequent experiments.

2.2.3 Diacetyl production

Diacetyl production was determined by HCl titration of fermented MFE [11]. A volume of 7.5 mL of hydroxylamine solution (1 M) was added to 25 mL of 24 h fermented MFE and to 25 mL of unfermented MFE broth (control). Both flasks were titrated with 0.1N HCl to a greenish yellow end point using bromophenol blue as indicator. The concentration of diacetyl produced was calculated as follows:

$$Ak = \frac{(R - S)(100E)}{W}$$

Where Ak (mg) is the quantity of diacetyl, R the volume (mL) of 0.1N HCl consumed in residual titration, S the volume (mL) of 0.1 N HCl consumed in titration of sample, E the equivalence factor of HCl to diacetyl is 21.52 mg and W is the volume of sample (25 mL).

2.2.4 Hydrogen peroxide production

The hydrogen peroxide production in fermented MFE was also quantified by titration. To 25 mL of this mixture, 25 ml of diluted H₂SO₄ (10%) was added. The preparation was then titrated with 0.1N potassium permanganate (KMnO₄). The end point was the point at which the pale pink color persisted for 15s before decolorization. Each ml of 0.1N KmnO₄ is equivalent to 1.701 mg of H₂O₂ [11]. The volume of H₂O₂ produced was then calculated as follows:

$$H_2O_2 \text{ concentration de } H_2O_2 = \frac{V(KMnO_4) \times N(KMnO_4) \times E \times 100}{V(H_2SO_4) \times V}$$

V(KMnO₄) : volume of KMnO₄ (ml)

N(KMnO₄) : KMnO₄ normality

E : KMnO₄ equivalence factor

V(H₂SO₄) : volume of H₂SO₄

V : volume of sample

2.2.5 Proteolytic activity

The proteolytic activity of isolate was determined using skim milk agar [12]. Standard plate count agar supplemented with 10% sterile skim milk was inoculated by touch with a fresh bacteria culture previously grown on MRS agar. The inoculated plates were incubated at 37°C for 24h. Proteolytic activity was estimated by the halo diameter.

2.2.6 Temperature, pH and salt stress

Strains ability to grow in various stress conditions was evaluated using MRS broth inoculated with 1% of overnight culture (i) at incubation temperature of 4, 30 or 45°C, (ii) with 2, 4, or 6.5 % NaCl (w/v) and (iii) at pH adjusted to 4 and 9.6 with HCl and NaOH respectively. The strains thermoresistance was evaluated by their survival after for 30 min incubation at 62°C [13].

2.2.7 Statistical analysis

Statistical analysis was carried out with XLSTAT™ software (version 2016.1). One-way analysis of variance (ANOVA) and Tukey's test were used to to perform multiple comparisons between analyzed variables, with significance level at p < 0.05. Relationships among the technological characteristics of the isolates were determined by Principal Component Analysis (PCA) using R software (version 3.6.1). The discriminating variables were growth and acidification kinetics, diacetyl production, hydrogen peroxide production, proteolytic activity.

3. RESULTS

3.1 LAB species identified

A total of 25 News isolates were selected and characterized by 16S (V1-V3 region) primers. As shown in **Table 1**, the sequencing of 16S rDNA and phylogenetic analysis revealed four species which are *Pediococcus acidilactici* (56%), *Lactobacillus fermentum* (24%), *Pediococcus pentosaceus* (12%) and *Lactobacillus plantarum* (8%).

Table 1. Molecular identification of new LAB isolated from fermented maize dough

Strain ID	Closely related Type strain	LAB	16SrRNA similarity (%)	Accession number
S1	<i>Pediococcus acidilactici</i>		98,98	OK427311
S5	<i>Pediococcus acidilactici</i>		98,78	OK427312
S8	<i>Pediococcus pentosaceus</i>		99,39	OK427313
S15	<i>Pediococcus acidilactici</i>		99,18	OK427314
S16	<i>Pediococcus acidilactici</i>		98,98	OK427315
S32	<i>Lactobacillus plantarum</i>		100	OK415420
S33	<i>Lactobacillus fermentum</i>		99,8	OK427317
S41	<i>Pediococcus acidilactici</i>		98,98	OK427318
S44	<i>Lactobacillus fermentum</i>		99,8	OK427319
S45	<i>Lactobacillus fermentum</i>		99,8	OK427320
S46	<i>Lactobacillus fermentum</i>		99,39	OK427321
S47	<i>Lactobacillus fermentum</i>		99,39	OK427322
S55	<i>Pediococcus acidilactici</i>		100	OK415422
S56	<i>Pediococcus acidilactici</i>		100	OK415423
S60	<i>Pediococcus acidilactici</i>		99,39	OK427325
S61	<i>Pediococcus acidilactici</i>		99,39	OK427326
S77	<i>Pediococcus acidilactici</i>		99,39	OK427327
S78	<i>Pediococcus acidilactici</i>		99,18	OK427328
S79	<i>Pediococcus acidilactici</i>		99,39	OK427329
S80	<i>Pediococcus pentosaceus</i>		99,39	OK427330
S81	<i>Lactobacillus fermentum</i>		99,8	OK427331
S121	<i>Lactobacillus plantarum</i>		100	OK415421
S130	<i>Pediococcus acidilactici</i>		98,98	OK427333
S141	<i>Pediococcus pentosaceus</i>		99,39	OK427334
S146	<i>Pediococcus acidilactici</i>		98,98	OK427335

3.2 Technological properties

3.2.1 Growth kinetics

The results of the growth kinetics in MFE of the 25 LAB strains from Fermented maize dough are shown in **Figure 1A**. Based on the measured optical density (OD) during fermentation, all strains displayed good abilities of growth in maize-based broth. Some strains such *Lactobacillus fermentum* S46 and *Pediococcus acidilactici* S141, contrarily to the others showed rapid growth in the earlier steps of fermentation, with an OD about 0,9 after only 6h. But at the end of fermentation, most of them displayed a good final OD value.

3.2.2 Acidification kinetics

The acidifying capacity of the 25 tested LAB strains was evaluated by determining their pH evolution during fermentation in MFE (Figure 1B). All isolates were able to lower pH during fermentation. Ten (10) LAB strains (*Pediococcus acidilactici* S55, S16, S77, S79, *Lactobacillus fermentum* S46, S47, S44; *Lactobacillus plantarum* S121, S32, and *Pediococcus pentosaceus* S141) showed abilities to rapidly acidify the MFE broth, with ΔpH values greater than 1 after only 6h. The other isolates had a slow onset of acidification, but at the end of fermentation displayed high ΔpH values.




Fig. 1. Growth Kinetics (A) and acidification abilities (B) of LAB strains during fermentation in MFE broth

3.2.3 Diacetyl production

Diacetyl production in MFE broth was carried by HCl titration. Most of tested LAB strains produced more or less diacetyl. Seven out of 25 isolates (*Lactobacillus fermentum* S46, S47, S81, *Pediococcus acidilactici* S60, S56, *Lactobacillus plantarum* S121 and *Pediococcus pentosaceus* S141) show best production with more than 80 mg (Figure 2A). For the other isolates, the production was average although some were very weak.

3.2.4 Hydrogen peroxide production

Potassium permanganate titration was used to determine hydrogen peroxide production by LAB strains. The concentrations of hydrogen peroxide produced are shown in Figure 2B. Ten strains obtained a production greater than 2 mM with the highest quantity (2,99 mM) produced by *Pediococcus acidilactici* S15.




Fig. 2. Diacetyl (A) and hydrogen peroxide (B) production of LAB strains during fermentation in MFE broth

3.2.5 Proteolytic activity

Protease activity of LAB strains was assessed based on the ability to produce clear zone on skim milk agar (Figure 3). Most strains achieved positive responses (Figure 4). Five strains revealed a high level of hydrolysis of milk proteins with lysis diameters greater than 8 mm. Those are *Lactobacillus plantarum* S121, S32, *Lactobacillus fermentum* S44 and *Pediococcus acidilactici* S55, S56. All other strains had an average activity between 3 to 7 mm.



Fig. 3. Proteolytic activity of LAB strains on agar medium




Fig. 4. Proteolytic activity of LAB strains

3.2.6 Salt, temperature and pH stress

The results of the tested LAB strains growth under different conditions are presented in **Table 2**. It was observed that all the strains grew at 30 and 45 °C. However, at 4 °C none of the tested strains was able to grow **except** *Lactobacillus fermentum* S45 and *Pediococcus pentosaceus* S80 which showed poor growth. At the concentration of 6.5% NaCl, *Lactobacillus fermentum* S45, *Pediococcus acidilactici* S56 and *Pediococcus pentosaceus* S80 displayed good growth, contrarily to the other strains. More or less growth was observed with most strains at NaCl concentrations of 2 and 4%. Moreover, all strains displayed growth at pH 4, however this growth is low for some isolates. Except *Pediococcus acidilactici* S15 and *Lactobacillus fermentum* S45, reasonable growth was observed with all strains at pH 9.6. After 30 min of incubation at 62°C, all strains were able to grow, however *Pediococcus acidilactici* S1 and S130 showed poor growth

Table 2. Growth characteristics of the strains at different pH, NaCl concentration and temperature.

Strains	Temperatures			pH		NaCl			Thermo-resistance
	4°C	30°C	45°C	4	9.6	2%	4%	6.5%	
<i>Pediococcus acidilactici</i> S1	-	+++	+++	++	++	++	++	+	+
<i>Pediococcus acidilactici</i> S5	-	+++	+++	++	+++	+++	+++	++	+++
<i>Pediococcus pentosaceus</i> S8	-	+++	++	+++	+++	+++	+++	++	+++
<i>Pediococcus acidilactici</i> S15	-	+++	++	++	+	+++	++	+	++
<i>Pediococcus acidilactici</i> S16	-	+++	+++	++	+++	+++	+++	+	+++
<i>Lactobacillus plantarum</i> S32	-	+++	+++	+++	+++	+++	+++	++	+++
<i>Lactobacillus fermentum</i> S33	-	+++	+++	++	++	++	++	-	+++
<i>Pediococcus acidilactici</i> S41	-	+++	+++	++	+++	+++	+++	++	+++
<i>Lactobacillus fermentum</i> S44	-	+++	+++	++	++	++	++	-	+++
<i>Lactobacillus fermentum</i> S45	+	+++	+++	+++	+++	+++	+++	+++	+++
<i>Lactobacillus fermentum</i> S46	-	+++	++	++	++	++	++	++	+++
<i>Lactobacillus fermentum</i> S47	-	+++	+++	+++	+	+++	++	+	+++
<i>Pediococcus acidilactici</i> S55	-	+++	+++	+++	+++	+++	+++	++	+++
<i>Pediococcus acidilactici</i> S56	-	+++	+++	+++	+++	+++	+++	+++	+++
<i>Pediococcus acidilactici</i> S60	-	+++	+++	+++	++	+++	+++	+	+++
<i>Pediococcus acidilactici</i> S61	-	+++	+++	+++	++	+++	++	-	+++
<i>Pediococcus acidilactici</i> S77	-	+++	+++	+++	+++	+++	+++	++	+++
<i>Pediococcus acidilactici</i> S78	-	+++	+++	++	++	+++	+++	++	+++
<i>Pediococcus acidilactici</i> S79	-	+++	++	+++	++	+++	++	+	++
<i>Pediococcus pentosaceus</i> S80	+	+++	+++	+++	++	+++	+++	+++	+++
<i>Lactobacillus fermentum</i> S81	-	+++	+++	+++	+++	+++	+++	+	+++
<i>Lactobacillus plantarum</i> S121	-	+++	+++	+++	+++	+++	+++	++	+++
<i>Pediococcus acidilactici</i> S130	-	+++	+++	+++	+++	+++	+++	++	+
<i>Pediococcus pentosaceus</i> S141	-	+++	++	++	+++	+++	+++	+	++
<i>Pediococcus acidilactici</i> S146	-	+++	+++	+++	+++	+++	++	+	+++

« +++ »: good growth; « ++ »: growth; « + »: poor growth; « - »: no growth

3.2.7 PCA analysis

The score plot (**Figure 5**) shows the relationship among the technological parameters used in this study. Apart from Growth and Acidification kinetics, no strict relationship found between the other parameters. Six strains form a distinct group in the positive direction for all variables. Those are *Pediococcus acidilactici* S15, S55, S56 ; *Lactobacillus plantarum* S32, S121 and *Lactobacillus fermentum* S44. They represent the bacteria with the highest activities for all variables combined.




Fig. 5. Principal Component Analysis (PCA) of the technological characteristics of LAB isolates

4. DISCUSSION

LAB constitute one of the most important bacterial groups used for the processing of fermented dairy, meat, vegetable, and cereal products [14]. In addition to their technological importance, many species of LAB act as probiotics [15]. When probiotics are added to foods, they must be able to survive in the product and become active when they enter the gastrointestinal tract of consumers. Therefore, in addition to assessing their probiotic potential, these cultures must also be tested for their functional and physiological properties [16]. In this study, 25 LAB strains isolated from fermented maize dough and identified by sequencing were subjected to different tests in order to select the most efficient ones for various biotechnological uses. Sequencing of the V1-V3 region of the 16S rRNA gene revealed the presence of four different species that are *Pediococcus acidilactici*, *Lactobacillus fermentum*, *Pediococcus pentosaceus* and *Lactobacillus plantarum*. As already mentioned by other authors, these species of lactic acid bacteria are those commonly found in fermented maize dough [4,17]. These bacteria play an important role in the fermentation of grain-based foods. [18] related that, *Lactobacillus plantarum* and *Lactobacillus fermentum* are important species in fermented maize dough. The importance of *Lactobacillus fermentum* in maize fermentation has been confirmed by previous studies in Ghana, Benin and Mexican products [19,20,21,22]. The high amylolytic activities found in different strains suggest that *Lactobacillus fermentum* could be a key organism for maize fermentation, making the large amounts of starch available to the overall community. In addition, the fermentation products (lactate, formate, and ethanol) may also serve as carbon sources for organisms, such as yeasts [17]. Interestingly, [4] reported a high prevalence of bacteriocin-producing *Lactobacillus fermentum* strains, and their detection in different stages of *Doklu* production indicates a high potential of these strains to grow and dominate the microbial population in the fermented maize dough.

Evaluation of acidification potential, proteolytic activity, diacetyl production and other tests have contributed to the technological and physiological characterization of the isolate LAB. Strains cultured for 48 hours in MFE were evaluated for their ability to grow correctly and to acidify the fermentation medium. The results obtained showed that all strains grow well in the broth. Some strains stood out for their rapid growth during the earlier stages of fermentation. After 24 h of fermentation, most strains reached an OD greater than 1 and this growth was independent of species. Knowledge on these growth parameters is necessary for more efficient industrial applications, considering the known nutritional requirement of this type of bacteria reported by some authors [23,24]. Acidifying activity is the most sought-after metabolic property of LAB used in the food industry. It is considered as an essential criterion for the selection of strains of technological interest. In this work, 75% of the strains showed a good acidification capacity with pH variation greater than 1.5 after 24 hours of fermentation. This good acidifying capacity of LAB has already been demonstrated by numerous studies [25,26,27]. The acidification capacity of these bacteria is due to the production of different types of organic acids that can contribute to make the environment unfavorable to the growth of certain microorganisms that could be undesirable from a hygienic point of view. Rapid acidification is also a priority in the development of lactic starter cultures.

After only 6 hours of fermentation, 10 strains showed pH variation greater than 1, among which are represented the four identified species. [28] had previously worked on the rapid acidification of LAB strains isolated from *Nunu* in Ghana, and these authors had also demonstrated the rapid acidification properties of *Lactobacillus plantarum* and *Lactobacillus fermentum* strains. The rapid acidification capacity of *Lactobacillus plantarum* has also been demonstrated by other authors [29,30].

The bacterial strains studied were also tested for the production of some metabolites of interest, namely diacetyl and hydrogen peroxide. Diacetyl is an essential aromatic compound. It is of industrial interest because it contributes to the flavor of many fermented foods. This aromatic compound is produced by lactic bacteria by citrate fermentation. Several bacteria of the genera *Lactococcus*, *Leuconostoc*, *Lactobacillus* and *Pediococcus*, can synthesize it [31]. In our study, the best production rate was obtained by seven strains including three *Lactobacillus fermentum*, two *Pediococcus acidilactici* and two *Lactobacillus plantarum*. In addition to contributing to the flavor of food products, diacetyl is characterized by broad antimicrobial activity [32]. It is able to inhibit Gram-negative bacteria and yeasts [33]. Hydrogen peroxide has also an antimicrobial effect. Its production was detected with all studied strains with concentrations higher than 2 mM for 10 of them. These results are in agreement with those obtained by [34] who tested 63 LAB strains and they were all good hydrogen peroxide producers. The hydrogen peroxide action can manifest itself both on undesirable germs and on those essential to the fermentation smooth running. Inhibition is achieved by oxidation of membrane lipids of target strains or by destruction of cellular protein structures [35].

Proteolytic activity of LAB strains was expressed on agar medium by lysis zones. According to [36], a strain is said to be correctly proteolytic if it presents a lysis zone diameter between 5 and 15 mm. In this study, 22 of the 25 strains tested (76%) obtained a proteolysis diameter greater than 5mm. LAB proteolytic systems play an important role in the ripening processes that give foods their rheological properties and organoleptic characteristics [37]. The use of these bacteria as food additives may be an ideal factor to modulate the proteolytic activity of a food product [38]. It should be noted, however, that excessive proteolysis can lead to the uncontrolled production of bitter peptides and other undesirable compounds [39].

LAB survival and growth under unfavorable conditions of temperature, salt concentration and pH were determined. No remarkable growth was observed when cultivating the strains at 4°C. At 45°C, very good growths were recorded for all strains showing their ability to grow at temperatures above normal LAB growth temperatures. According to [40], most lactic strains grow in a temperature range between 15°C and 42°C. Strains that can remain viable at higher temperatures are called "thermoreistant". The 6.5% NaCl level allowed a good growth of only 3 strains out of the 25 studied strains. These are *Lactobacillus fermentum* S45, *Pediococcus acidilactici* S56 and *Pediococcus pentosaceus* S80. The ability of these strains to grow properly at 6.5% NaCl level could be distinctive since the literature generally presents an intolerance of lactic strains to this concentration [41]. Salt-resistant strains can be of technological interest especially in fermentations of salted products (salted cheeses, olives, cucumbers, meats, etc.) [42]. Regarding pH, the optimal values required for the growth and survival of bacteria are generally between 4 and 8. Under abnormal pH conditions, normal cellular components are not synthesized, which affects cell division and prevents growth [43]. Most of the strains studied were able to grow at pH 4, but at different intensities. Not surprisingly, pH 9.6 was the most critical. However, good growth was obtained for 13 of the strains. Despite treatment at 62°C for 30 min, 20 LAB strains showed good growth after incubation. It should be noted that during the drying of probiotics, the microorganisms are subjected to treatments involving extreme temperatures that can be very high (up to 200°C) during spray drying or very low (down to -196°C) during freeze drying and storage [44]. In general, resistance to these stresses is higher when the cells are previously exposed and adapted to a previous treatment with a homologous stress. For example, better heat tolerance was obtained when *Lactobacillus rhamnosus* GG was pre-exposed to 60°C for 20 min [45].

4. CONCLUSION

Based on partial 16S rRNA gene sequencing, 25 LAB isolated from fermented maize dough used for *doklu* production, were identified as *Pediococcus acidilactici*, *Lactobacillus fermentum*, *Lactobacillus plantarum* and *Pediococcus pentosaceus*. The obtained results show that certain isolates have promising technological properties for possible industrial applications. These strains have generally shown their capacity for good growth, acidification and the production of certain interesting metabolites such as diacetyl and hydrogen peroxide in a medium prepared from maize flour. They also demonstrated their ability to survive and growth under unfavorable conditions.

REFERENCES

1. De Vuyst L, Neysens P. The sourdough microflora: biodiversity and metabolic interactions. Trends in Food Science and Technology. 2005 ; 16, 43-56.

2. Nadeem M, Anjum FM, Amir RM, Khan MR, Hussain S, Javed MS. An overview of antinutritional factors in cereal grains with special reference to wheat – A review. *Pakistan Journal of Food Sciences*. 2010 ; 20: 1-4.
3. Assouhoun NMC, Djéni NT, N'Guessan KF, Koussémon M. Preliminary study on antimicrobial properties of lactic acid bacteria involved in the fermentation of corn dough during *Doklu* processing in Côte d'Ivoire. *Global Science Books*. 2012 ; 6 : 65-70.
4. Assouhoun-Djeni NMC, Djéni NT, Messaoudi S, Lhomme E, Koussémon-Camara M, Ouassa T, Chobert JM, Onno B, Dousset X. Biodiversity, dynamics and antimicrobial activity of lactic acid bacteria involved in the fermentation of maize flour for *doklu* production in Côte d'Ivoire. *Food Control*, 2016 ; 62 : 397-404.
5. Berbegal C, Peña N, Russo P, Grieco F, Pardo I, Ferrer S, Spano G, Capozzi V. Technological properties of *Lactobacillus plantarum* strains isolated from grape must fermentation. *Food Microbiology*. 2016 ; 27:187–194.
6. Das AJ, Das MJ, Miyaji T, Deka SC. Growth and metabolic characterization of four lactic acid bacteria species isolated from rice beer prepared in Assam, India. *Access Microbiology*, 2019 ; 1 : 1-14.
7. Tannock GW. Medical Importance of the Normal Microflora. Boston. Springer. 1999 ; 515 p.
8. Sharma K, Attri S, Goel G. Selection and Evaluation of Probiotic and Functional Characteristics of Autochthonous Lactic Acid Bacteria Isolated from Fermented Wheat Flour Dough Babroo. *Probiotics and Antimicrobial Proteins*. 2018 ; 11 : 774-784.
9. Biswas L, Gruss A, Ehrlich SD, Maguin E. High efficiency gene inactivation and replacement system for gram positive bacteria. *Journal of Bacteriology*. 1993 ; 175 : 3628-3635.
10. Alfonzo A, Giusi V, Onofrio C, Rosalia DG, Raimondo G, Nicola F, Giancarlo M, Luca S. Diversity and Technological Potential of Lactic Acid Bacteria of Wheat Flours. *Food Microbiology*. 2013 ; 36(2) : 343-54.
11. Edema M O, Sanni AI. Functional properties of selected starter cultures for sour maize bread. *Food Microbiology*. 2008 ; 25: 616-625.
12. Thapa N, Pal j, Tamang JP. Phenotypic identification and technological properties of lactic acid bacteria isolated from traditionally fish products of the eastern himalayas. *International Journal of Food Microbiol*. 2006; 107 : 8-33.
13. Hassaine O. Caractéristiques d'intérêt technologiques de souches de bactéries lactiques isolées de lait camélien du Sud Algérien. Thèse de doctorat, Université d'Oran Es-sénia, Oran. 2013 ; 180 p.
14. Hurtado A, Reguant C, Bordons A, Rozès N. Lactic acid bacteria from fermented table olives. *Food Microbiology*, 2012 ; 31 : 1-8.
15. Souza JV, Dias FS. Protective, technological, and functional properties of select autochthonous lactic acid bacteria from goat dairy products. *Current Opinion In Food Science*, 2017 ; 13 : 1-9.
16. Bujnakova D, Strakova E. Safety, probiotic and technological properties of *Lactobacilli* isolated from unpasteurised ovine and caprine cheeses. *Annual Microbiology*, 2017 ; 67 : 813-826.
17. Ben-Omar N, Ampe F. Microbial community dynamics during production of the Mexican fermented maize dough Pozol. *Applied and Environmental Microbiology*, 2000 ; 66 : 3664-3673.
18. Chaves-Lopez C, Serio A, Delgado-Ospina J, Rossi C, Grande-Tovar CD, Paparella A. Exploring the Bacterial Microbiota of Colombian Fermented Maize Dough "Masa Agría" (Maiz Añejo). *Frontiers Microbiology*, 2016 ; 7 : 1168.
19. Obinna-Echem P. C., Kuri V. & Beal J. (2014). Evaluation of the microbial community, acidity and proximate composition of akamu, a fermented maize food. *Journal of the Science of Food and Agriculture*. 94 : 331-340.
20. Oguntoyinbo FA, Turlomousis P, Gasson MJ, Narbada A. Analysis of bacterial communities of traditional fermented West African cereal foods using culture independent methods. *International Journal of Food Microbiology*. 2011 ;145 : 205-210.
21. Ampe F, Ben-Omar N, Moizan C, Wachter C, Guyot JP. Polyphasic study of the spatial distribution of microorganisms in Mexican pozol, a fermented maize dough, demonstrates the need for cultivation- independent methods to investigate traditional fermentations. *Applied and Environmental Microbiology*. 1999 ; 65 : 5464-5473.
22. Hayford AE, Petersen A, Vogensen FK, Jakobsen M. Use of conserved randomly amplified polymorphic DNA (RAPD) fragments and RAPD pattern for characterization of *Lactobacillus fermentum* in Ghanaian fermented maize dough. *Applied and Environmental Microbiology*. 1999 ; 65 : 3213-3221.
23. Loubière P, Novak L, Coccain-Bousquet M, Lindley N. Besoins nutritionnels des bactéries lactiques : interactions entre flux de carbone et d'azote. INRA Editions. 1996 ; 76 : 5-12.
24. Jensen PR, Hammer K. Minimal requirements for exponential growth of *Lactococcus tectis*. *Applied and Environmental Microbiology*. 1993 ; 59 : 4363-4366.
25. Maghnia D. Etude de potentiel technologique des bactéries lactiques isolées des aliments fermentés traditionnels algériens. *Mémoire de magister en Microbiologie Alimentaire*. Université d'Oran-Es-Senia. 2011 ; 126 p.
26. Idoui T, Boudjerda J, Leghouchi E, Karam NE. Lactic acid bacteria from "Sheep's Dhan", a traditional butter from sheep's milk: Isolation, identification and major technological traits. *Grasas Y Aceites*. 2009 ; 60 (2) : 177-183.
27. Alonso-Calleja C, Carballo J, Capita R, Bernardo A, Garcia-Lopez ML. Changes in the microflora of Valdeteja raw goats milk cheese throughout manufacturing and ripening. *Lebensmittel-Wissenschaft Und- Technologie*. 2002 ; 35 : 222-232.

28. Akabanda F, Owusu-Kwarteng J, Tano-Debrah K, Parkouda C, Jespersen L. The Use of Lactic Acid Bacteria Starter Culture in the Production of Nunu, a Spontaneously Fermented Milk Product in Ghana. *International Journal of Food Science*, 2014. 1-11.
29. Idoui T, Karam NE. Lactic acid bacteria from *Jijel's* traditional butter: isolation, identification and major technological traits. *Grasas y Aceites*. 2008 ; 59 (4) : 361-367.
30. Haddadin JSY. Kinetic studies and sensorial analysis of Lactic Acid Bacteria isolated from white cheese made from sheep raw mil. *Pakistan Journal of Nutrition*. 2005 ; 4 : 78-84.
31. Leveau JY, Boiux M, De Roissart HB. La flore lactique : technique d'analyse et de contrôle dans les industries agroalimentaires. 2e Ed., Tec & Doc, Lavoisier. 1991 ; 3: 240 p.
32. Lanciotti R, Patrignani F, Bagnolini F, Guerzoni ME, Gardini F. Evaluation of diacetyl antimicrobial activity against *Escherichia coli*, *Listeria monocytogenes* and *Staphylococcus aureus*. *Food Microbiology*. 2003 ; 20 : 537-543.
33. Ammor S, Tauveron G, Dufour E, Chevallier I. Antibacterial activity of lactic acid bacteria against spoilage and pathogenic bacteria isolated from the same meat small-scale facility: 1-Screening and characterization of the antibacterial compounds. *Food Control*. 2006 ; 17 : 454-461.
34. Enitan A, Adeyemo J, Ogunbanwo ST. Influence of growth conditions and nutritional requirements on the production of hydrogen peroxide by lactic acid bacteria. *African Journal of Microbiology Research*. 2011 ; 5 : 2059-2066.
35. Zalan Z, Barath A, Halasz A. Influence of growth medium on hydrogen peroxide and bacteriocin production of *Lactobacillus* strains. *Food Technology and Biotechnology* 2005 ; 43 : 219-225.
36. Vuilleumard JC. Microbiologie des aliments. Evolution de l'activité protéolytique des bactéries lactiques. Paris. *Tec & Doc*, Lavoisier. 1986 ; 65 p.
37. Law BA, Kolstad J. Proteolytic systems in lactic acid bacteria. *Antonie van Leeuwenhoek*. 1983 ; 49: 225-245.
38. Sanders ME, Merenstein D, Merrifield CA, Hutkins R. Probiotics for human use. *Food and Nutrition Bulletin*. 2018 ; 43 : 212-225.
39. Buffa M, Morais J, Jimenez-Belenguer A, Hernandez-Gimenez E, Guamis B. Technological characterization of lactic acid bacteria isolated from raw ewes' milk for cheese making. *Milchwissenschaft*. 2005 ; 61 : 404-407.
40. Tailliez P. Les lactobacilles : propriétés, habitats, rôle physiologique et intérêt en santé humaine. *Actualités microbiologiques*. 2004 ; 6 (1) : 35-41.
41. Guiraud JP. Microbiologie alimentaire. Dunod. 1998 ; 615p.
42. Karam-Zadi H, Karam N. Bactéries lactiques du lait de chamelle d'Algérie : mise en évidence de souches de *Lactococcus* résistantes au sel. *Tropicultura*. 2006 ; 24 (3) : 153-156.
43. Das AJ, Das MJ, Miyaji T, Deka SC. Growth and metabolic characterization of four lactic acid bacteria species isolated from rice beer prepared in Assam, India. *Access Microbiology*. 2019 ; 1 : 1-14.
44. Mills S, Stanton C, Fitzgerald GF, Ross RP. Enhancing the stress responses of probiotics for a lifestyle from gut to product and back again. *Microbial Cell Factories*. 2011 ; 10 : 1-19.
45. Corcoran BM, Stanton C, Fitzgerald G, Ross RP. Life under stress : the probiotic stress response and how it may be manipulated. *Current Pharmaceutical Design*. 2008 ; 14 : 1382-1399.