

Differential changes in some antioxidant enzymes and biomolecules in leaves of tomato infected by *Phytophthora infestans* (Mont.) de Bary.

Abstract

In Cameroon, tomato yields remain low due to attack by pathogens and insects. *Phytophthora infestans* is a phytopathogenic fungus responsible for the downy mildew of tomato, a fungal disease responsible for enormous economic losses. To contribute to the search for means of control, the stimulatory effect of the tomato defence system by extracts of some plants was evaluated. Tomato plants were treated with the aqueous macerates of *Callistemon citrinus* (*C. citrinus*) and *Cymbopogon citratus* (*C. citratus*) at 10% (W/V). After 4 days of spraying with the macerates, plants were inoculated with a virulent strain of PI. Tomato leaves collected at 2-day intervals for 10 days were used to determine the contents of chlorophyll a, proteins, carbohydrates, lipids, amino acids and proline. The activities of the main antioxidant enzymes were also assessed: ascorbate peroxidase (APX), phenylalanine ammonia lyase (PAL), peroxidase (POX) and superoxide dismutase (SOD). The results showed that treatment of tomato plants with aqueous macerates of *C. citrinus* and *C. citratus* and their inoculation with PI induced an increase in chlorophyll a, protein, carbohydrate, lipid, amino acid and proline contents in tomato leaves; an increase in APX, PAL, POX, and SOD activities. Our results suggest that increased metabolite synthesis on the one hand, and increased enzymatic activities, on the other hand, could mitigate the oxidative damage that occurs during pathogen expansion. Aqueous macerates of *C. citrinus* and *C. citratus* could be used as natural products to stimulate the defence system in tomato against *Phytophthora infestans*.

Keywords: Tomato, Fungal diseases, Plant extracts, Antifungal, Induced resistance

1. Introduction

Tomato (*Lycopersicon esculentum* Mill.) is a vegetable cultivated in almost all countries of the world. Tomato has a high biological value due to its content in carotenes, vitamins C and E and lycopene and a balanced content of mineral elements (Keatinge, 2012).

Its global production in 2017 was estimated at 182.30 million tons with an average yield of 188 tons/ha. In Cameroon, tomato remains the most important vegetable in terms of production. In 2017, its production was estimated at 1.27 million tons with a yield of 12.14 tons/ha (FAOSTAT, 2019). This yield is far from the world yield, this gap is mainly justified by many constraints faced by tomato cultivation. Among these, the most striking in Cameroon are adverse climatic conditions and pests, susceptibility to diseases (alternariosis, fusariosis, mildew) (Mekam *et al.*, 2019). Indeed, tomatoes are the target of many bacterial and fungal diseases. Phytosanitary analysis of crops and sampling carried out in tomato growing sites in Cameroon in the Centre (Yaounde) and Western (Dschang) regions revealed a recurrence of fungal diseases caused by several fungi including *Fusarium oxysporum* f.sp. *lycopersici* that causes leaf yellowing, wilting and damping-off. This plant pathogenic fungus causes field losses ranging from 46 to 100% (Dakole *et al.*, 2016).

Nowadays, the control of most biotic plant diseases is mainly based on the very intensive use of synthetic chemical bactericides and fungicides (Konje *et al.*, 2019). However, despite the acceptable results, chemical control has several limitations, namely environmental pollution, high cost of synthetic chemicals, the risk of pathogen resistance to chemical substances, and human intoxication, especially since analyses of vegetables and fruits in Cameroon have shown pesticide residues. These limitations motivate and justify the search for effective and eco-friendly alternatives to control pathogenic microorganisms and crop pests. Thus, the use of natural substances (essential oils and solvent extracts) as plant disease control agents is receiving increasing attention. In Cameroon, local plant extracts have been reported as having properties *in vitro* and *in vivo* against various pathogens (Mekam *et al.*, 2019). Botanicals with antifungal compounds have been identified and can be exploited for the management of plant diseases because they have low mammalian toxicity, target specificity, biodegradability and contain many active ingredients (Kagale *et al.*, 2004). *Callistemon citrinus*, *Cymbopogon citratus* and *Oxalis barelii* because of their antifungal activity notably against *Phytophthora infestans* and *Fusarium oxysporum* of tomato are used as a biocontrol agent (Dakole *et al.*, 2016). Plant extracts contain compounds that not only have a direct antimicrobial effect on the pathogen, but also stimulate the plant's natural defences, making them one of the most promising alternatives among crop protection strategies (Abdelkhalek *et al.*, 2021). Studies have shown that spraying rice plants with *Datura metel* leaf extracts can induce systemic resistance of this plant against *Rhizoctonia solani* and *Xanthomonas oryzae* pv *oryzae* through the accumulation of pathogenicity-related proteins (Samir *et al.*, 2019). Treatment of plants with aqueous extracts has enabled plants to resist

pathogenic microorganisms and control several fungal diseases through metabolic changes (accumulation of phenolic compounds and antioxidant defence enzymes) (Fangue *et al.*, 2021).

The objective of this work was to evaluate the stimulating effect of the defence system of aqueous macerates of *Callistemon citrinus* and *Cymbopogon citratus* in tomato (*Lycopersicon esculentum* Mill.) infected by *Phytophthora infestans* Mont. de Bary.

2. Materials and methods

2.1. Plant and soil material

Seeds of the Rio Grande variety of tomato purchased at the Mfoundi market (Yaounde, Cameroon).

The soil used was obtained from the soil of Nkolbisson (Yaounde, Cameroon).

The leaves and stems of *Callistemon citrinus* and *Cymbopogon citratus* (DC) STAFW were harvested in Mbandjock (Yaoundé, Cameroon).

Salicylic acid, a natural stimulator of the plant defence system, was used as a positive control.

2.2 Fungal material

A strain of *Phytophthora infestans* (Mont.) de Bary was isolated from tomato fruits and leaves harvested in the field and showing downy mildew symptoms. Its isolation and identification were done in the Phytopathology laboratories of the *Institut de Recherche Agricole pour le Développement (IRAD)* of Nkolbisson (Yaounde, Cameroon) and the *Institut du Développement Rural (IDR)* of the *Université Polytechnique de Bobo-Dioulasso* (Burkina Faso). Identification was done according to Agarwal *et al.*, (1989) and Mathur and Kongsdal (2003) and confirmed by the Phytopathology Laboratory of the Danish Seed Health Center (DSHC), Copenhagen, Denmark. The pathogenicity of these strains was proven on 20-day-old tomato plants from which the same pathogens were re-isolated. The strain of *Phytophthora infestans* (Mont.) de Bary was maintained on V8 agar medium in 90 mm diameter Petri dishes at 20 ± 2 °C in the dark. The strain was cultured for 7-10 days to allow sporulation of the fungi, subcultured from time to time to keep the growth active and kept at 5°C in a V8 agar medium. Fourteen-day cultures were used for the inoculation of tomato plants.

2.3. Preparation of aqueous macerates and salicylic acid solutions

The plant material (*Callistemon citrinus* and *Cymbopogon citratus*) was carefully harvested and shade-dried in the greenhouse for 14 days and then crushed with a mill. A 10% (W/V) macerate was prepared. The mixture was kept for 48 hours in the greenhouse and at

room temperature in the laboratory. The aqueous macerates were obtained by pressing and filtering through a sieve and a fine cloth. The extracts obtained were used for spraying tomato plants.

Salicylic acid was used as a positive control and prepared at 0.01% (W/V) as described by **Mandal *et al.*, (2013)**.

2.4 Obtaining tomato plants

The seeds were weighed and sterilized with 1% (v/v) sodium hypochlorite solution. Then they were rinsed twice with sterile distilled water for 02 minutes while shaking vigorously each time.

The seeds obtained were sown in a nursery tray containing sterile soil consisting of a sand/soil mixture in the proportions 1:3 (v/v). The plants were left to grow with regular watering for 14 days and then transplanted into 10 kg pots (1 plant per pot) in a greenhouse. 10 days after transplanting, the tomato plants were divided into 4 lots:

- The first batch was made up of control plants sprayed with distilled water (S);
- The 2nd batch consisted of plants treated with 10% macerate *C. citratus* (Tr Cymb);
- The 3rd batch consisted of plants treated with 10% macerated *C. citrinus* (Tr Call);
- The 4th batch consisted of plants treated with salicylic acid (Tr AS).

After 3 days of treatment (spraying), each batch was divided into 2 subgroups: subgroup 1 consisting of tomato plants of each of the above treatments but not inoculated: S, Tr Cymb, Tr Call, Tr AS; subgroup 2 consisting of tomato plants of each of the above treatments inoculated with PI: I (S having been inoculated), (Tr Cymb + I), (Tr Call + I), (Tr AS +I).

For each treatment, five replicates were conducted independently and the pots were arranged in the greenhouse in a completely randomised design.

The experiment was conducted in a completely randomised block design.

2.5. Pathogen Inoculation

From a 14-day PI culture on agar V8 medium, the fungal suspension was prepared and adjusted to 5×10^5 conidia/ml. Five (2) ml of the conidia suspension were homogenized and inoculated onto the tomato leaves. The inoculated plants (subgroup 2) were covered with black polyethylene bags for 48 h to create adequate moisture for **fungal** growth and development. Non-inoculated tomato plants (subgroup 1) were used as controls. For biochemical assays, leaves were collected on the day of inoculation and at 2-day intervals during 10 days. These samples were washed with tap water, drained, weighed and frozen for further analysis.

2.6. Determination of Chlorophyll a

The extraction of chlorophyll was done in 80% acetone at 4°C. For this purpose, 5 g of leaves were cut into slices of 2 cm diameter and put into test tubes containing 3 ml of 80% acetone. The tubes were incubated in the dark for 24 hours to allow solubilisation of the photosynthetic pigments. The determination of chlorophyll a was done by determining the optical densities at 663 and 646 and the calculation was done according to the formula of **Lichtenthaler and Welburn (1983)**.

$$\text{Chlorophyll a (mg.g}^{-1}\text{FW)} = 12.21 \text{ OD}_{663} - 2.81 \text{ OD}_{646}$$

2.7. Extraction and determination of total soluble proteins

The extraction of the total soluble proteins was carried out according to the method described by **Mbouobda et al., (2007)**. Two (2) grams of leaves were ground cold (4° C) in 4 ml of TAME buffer (Tris 0.5 M; 0.3M ascorbic acid; 2% permethylethanol; EDTA 0.01 M; Triton x100, 2%, pH 7) supplemented with the addition of polyvinyl pyrrolidone (PVP) and sterile fine sand. The grind was incubated for 30 min at 4° C and then centrifuged at 10,000 g for 20 min. The recovered supernatant constituting the total soluble protein extract was stored at -20° C. for subsequent analysis. The amount of protein in the crude extract was determined using the **Bradford method (1976)** with bovine serum albumin (BSA) as standard. The protein content was expressed in mg.g⁻¹ of fresh matter.

2.8. Determination of carbohydrates

The plant samples were boiled in 80% ethanol for extraction. Test tubes containing 1 ml of ethanolic extract each were placed into the water bath for evaporation. The residue and 1 ml of distilled water were incubated at 49° C for 30 min. The solution was then neutralised with 1N NaOH using methyl red as a colour indicator. 1 ml of Nelson's reagent was added to each tube. The tubes were heated for 20 min in a boiling water bath, were cooled and 1 ml of the arsenomolybdate was added. The solution was mixed well and diluted to obtain 25 ml and the OD was measured at 495 nm using a spectrophotometer. The concentration of reducing sugars was calculated from the glucose standard and expressed in µg.g⁻¹ of fresh matter (**Nelson, 1944**).

2.9 Determination of total lipids

The tube containing 350 μl of supernatant (with 80% ethanol extract) was evaporated to complete desiccation in the heating blocks at 95° C. 37.5 μl of 96% sulphuric acid were added and the mixture was heated for five minutes. After cooling, 1000 μl of Vanillin was added and then after 10 to 15 min to read the OD at 525 nm after development of a characteristic red coloration. A calibration curve obtained from commercial vegetable oil was used to calculate the lipid concentration (**Foleh et al., 1957**). The lipid content was expressed in $\mu\text{g.g}^{-1}$ of fresh material.

2.3.7 Determination of total free amino acids

In test tubes, 1 ml of ethanol extract from the leaves was taken up in 25 ml of distilled water and neutralised with 0.1N NaOH using methyl red as a colour indicator (**Moore and Stein, 1948**). 1 ml of ninhydrin was added and the mixture was heated in a boiling water bath for 20 min, then 5 ml of a dilution solution (distilled water/n-propanol in equal volume) was added. This was followed by cooling and dilution to 25 ml. The absorbance was measured at 570 nm with a spectrophotometer. The standard line uses a leucine solution. The total amino acid content was expressed in mg.g^{-1} of fresh matter.

2.3.8. Determination of proline

The Proline Assay was carried out on a sample of 100 mg of fresh material taken and placed in a test tube to which 2 ml of 40% methanol was added. The sample was heated for 1 hour in a water bath at 85°C. After cooling, 1 ml of the extraction solution was added to 1 ml of the mixture (distilled water/acetic acid/ortho-phosphoric acid) (6/15/4) and 2 ml of ninhydrin. The mixture was brought to a boil at 100°C for 30 min in a water bath (**Bates, 1973**). After cooling, 5 ml of toluene was added after vortexing. The OD reading was taken at 528 nm. The proline content was calculated from the proline standard and expressed as $\mu\text{g.g}^{-1}$ of fresh material.

2.3.9. Assessment of Ascorbate peroxidase (APX) activity

APX activity (EC 1.11.1) was assessed according to the method of **Nakano and Asada (1981)**. Enzyme activity was determined by following the decrease in absorbance of ascorbate at 290 nm. The reaction medium consisted of: 50 mM phosphate buffer, pH 7; 0.5 mM ascorbic acid; H_2O_2 0.1 mM (5 / 2 / 1) (V/V/V) in a final volume of 5 ml. To each test tube containing 5 ml of the reaction medium, the enzyme extract (10 μl) was added. The optical density was read at 290 nm against a blank in which the enzyme extract is replaced by the extraction buffer. The molar extinction coefficient $2.8 \text{ mM}^{-1}.\text{cm}^{-1}$ was used to calculate the ascorbate peroxidase activity expressed as moles of H_2O_2 reduced. $\text{min}^{-1}.\text{g}^{-1}$ of fresh

material. One Unit of Ascorbate Activity is required for the reduction of 1 mole of H_2O_2 per minute at 25°C.

9. Evaluation of phenylalanine ammonia lyase (PAL) activity

PAL activity (EC.4.3.1.5) was assessed based on the rate of cinnamic acid production as described by **Saunders and McClure (1975)**. Thus, 1 ml of 50 mM Tris-HCl buffer pH 8.8 containing β -mercaptoethanol, 0.5 ml of 10 mM L-phenylalanine, 0.4 ml of distilled water and 0.1 ml of enzyme extract were incubated at 37° C for 1 h. The reaction was stopped by adding 0.5 ml of 6M HCl and the product was extracted with 15 ml of ethyl acetate followed by evaporation of the extraction solvent using a rotary evaporator. The solid residue was taken up in 3 ml of 0.05 M NaOH and the concentration of cinnamic acid was quantified by reading the optical density at 290 nm. One PAL activity Unit is equal to 1 μ mol of cinnamic acid produced per minute per g of fresh plant material.

.3.10. Evaluation of peroxidase activity (POX)

POX activity (EC.1.11.1.7) was assessed by the method described by **Hammerschmidt et al., (1982)**. The reaction medium (3 ml) consisted of 10 mM sodium phosphate buffer pH 6 containing 10 mM H_2O_2 at 0.25% (V/V). A volume of 100 μ l of enzyme extract was added to initiate the reaction. The enzyme activity was determined by calculating the ratio of optical Density at 470 nm per minute per 0.01 which was defined as the Activity Unit of peroxidase activity expressed in terms of Unit per g of fresh plant material.

2.3.11. Assessment of superoxide dismutase (SOD) activity

SOD activity (EC.1.15.1.1) was assessed using the method described by **Beauchamp and Fridovich (1971)**. It measures the ability of the enzyme to inhibit the photochemical reduction of nitro blue tetrazolium chloride (NBT). The reaction medium (3 ml) consists of: 50 mM phosphate buffer pH 7.8; 13mM L-methionine; NBT 75 μ M; EDTA 1 mM; 0.05M sodium carbonate; 20 μ l of enzyme extract. Riboflavin 2 μ M was added after all reagents and the reaction was initiated by placing the test tube 30 cm below a 15W ampoule for 10 min at 25°C. The tube was then transferred into the dark for 10 min and the optical density read at 560 nm. One SOD Activity Unit was defined as the amount of enzyme required to inhibit the reduction of NBT by 50%. SOD activity was expressed as Units per minute per g of fresh plant material.

2.5. Statistical analyzes

The results obtained were subjected to statistical analysis for the calculation of means, standard deviations and the search for significant differences, using the SPSS 22.1 software.

The one-factor ANOVA test coupled with the Student-Newman-Keuls test was used to evaluate the Smallest Significant Difference (PPDS) at $P < 0.05$

3. Results

3.1. Effect of plant extracts on chlorophyll a content

In healthy plants, the Chl a content on day 0 was $12.50 \pm 0.63 \text{ mg. g}^{-1}$ and did not vary significantly with time until day 8. When the plants were inoculated (I), there was a significant decrease in Chl a on day 2 and on day 10, the percentage was 37%. In treated plants (Tr), the Chl a content increased significantly ($P < 0.05$) for all treatments. On day 10, the content increased by 178.03% and 186.58% with *C. citrinus* and *C. citratus* respectively. In treated and inoculated plants (Tr + I), Chl a content decreased significantly on days 2 and 4 in *C. citratus* treated plants and on days 2, 4 and 6 in *C. citrinus* treated plants. A gradual and significant increase was then noted for all treatments. On day 10, the increase was 173.26% and 271.13% for *C. citrinus* and *C. citratus* respectively. The Chl a content increased significantly from day 2 for all treatments (**Table 1**).

Table 1: chlorophyll a content (expressed in $\text{mg. g}^{-1}\text{FW}$) in tomato leaves on a time course after treatment with plants extracts and inoculation with PI and in the control

DAI	<i>C. citrinus</i>				<i>C. citratus</i>		Salicylic acid	
	S	I	Tr	Tr+I	Tr	Tr+I	Tr	Tr+I
0	12.50 ^b	12.10 ^b	11.70 ^a	11.56 ^b	11.70 ^a	11.57 ^b	11.70 ^a	11.57 ^a
	± 0.63	± 0.99	± 0.08	± 0.21	± 0.18	± 0.21	± 0.18	± 0.21
2	11.70 ^{ab}	7.16 ^a	12.50 ^b	6.50 ^a	13.10 ^{ab}	5.10 ^a	44.43 ^b	22.80 ^c
	± 0.30	± 0.20	± 0.52	± 0.10	± 0.20	± 0.12	± 1.97	± 1.50
4	11.67 ^{ab}	6.83 ^a	13.50 ^c	4.33 ^a	15.50 ^b	4.83 ^a	44.10 ^b	19.90 ^b
	± 0.40	± 0.53	± 0.30	± 0.15	± 0.30	± 0.13	± 1.20	± 1.20
6	11.83 ^{ab}	5.16 ^a	13.16 ^c	5.00 ^a	16.58 ^c	20.66 ^c	56.74 ^c	22.83 ^c
	± 0.15	± 0.13	± 0.15	± 0.10	± 0.30	± 1.53	± 1.97	± 1.20
8	11.40 ^{ab}	5.16 ^a	20.83 ^d	21.80 ^c	22.00 ^d	27.00 ^d	66.10 ^d	87.40 ^d
	± 0.20	± 0.12	± 1.53	± 1.10	± 2.00	± 2.00	± 5.20	± 5.20
10	11.33 ^a	4.50 ^a	20.83 ^d	20.03 ^c	21.83 ^d	31.37 ^e	69.67 ^e	101.00 ^e
	± 0.45	± 0.20	± 1.25	± 1.53	± 1.15	± 1.53	± 1.89	± 2.10

Data are means \pm SD of five replications. Different letters after the numbers in each column indicate a significant difference ($p < 0.05$) between values, Duncan test.

DAI: day after inoculation; S: control plants; I: plants inoculated with PI; Tr: plants treated with extract or salicylic acid; Tr + I: plants treated and inoculated with PI.

3.2. Effect of plant extracts on protein content

In healthy plants, the protein content was 38.77 ± 0.76 mg. g⁻¹FW on day 0. It increased significantly ($P < 0.05$) from day 2 and reached the peak on day 6. When the plants are inoculated (I), the protein content of 40.43 ± 1.04 mg. g⁻¹ FW at day 0 decreased at day 4 and then there was a progressive increase until day 10. In plants treated with plant extracts, the protein content increased progressively and significantly over time by more than 100% at day 10 for all treatments. Inoculation after treatment with plant extracts and salicylic acid (Tr + I) resulted in a significant and progressive increase in protein content over time. Protein contents were 2 to 5 times higher on day 10 for all treatments (Figure 1).




Figure 1: Proteins content (expressed in mg.g⁻¹ FW) in leaves of tomato plants on a time course after treatment with plants extracts and inoculation with PI and in the control. Data presented are the means \pm SD of five replicates. Different letters indicate a significant difference ($p < 0.05$) between values, Duncan test. A: *C. citrinus*; B: *C. citratus*; C: Salicylic acid. S: control plants; I: plants inoculated with PI; Tr *C. citrinus*, Tr *C.*

citrus, Tr Salicylic acid: plants treated respectively with *Callistemon citrinus*, *Cymbopogon citratus* and salicylic acid. Tr + I *C. citrinus*, Tr + I *C. citratus*, Tr + I salicylic acid: Plants treated with *Callistemon citrinus*, *Cymbopogon citratus* and salicylic acid and inoculated with PI respectively.

2.3. Effect of plant extracts on carbohydrates content

In healthy plants (S), the carbohydrate content was $77.33 \pm 1.53 \text{ mg g}^{-1}$ FP at day 0, which increased significantly from day 2 to day 8. In the inoculated plants (I), the carbohydrate content increased gradually and significantly ($P < 0.05$) with a peak on day 6, an increase of 183.46%. After treatment with plant extracts, this content increased by more than 100% from day 2 for all treatments and reached a maximum value on day 6. In the inoculated plants after treatment (Tr + I), there was a significant ($P < 0.05$) increase in carbohydrate content from day 2. The percentage increase was 205.12% and 223.50% with *C. citrinus* and *C. citratus* respectively. This content increased gradually over time. In general, the increase of this content is higher than in the treated plants (Tr) (Figure 2).




Figure 2: Sugars content (expressed in $\text{mg}\cdot\text{g}^{-1}$ FW) in leaves of tomato plants on a time course after treatment with plants extracts and inoculation with PI and in the control. Data presented are the means \pm SD of five replicates. Different letters indicate a significant difference ($p < 0.05$) between values, Duncan test. **A:** *C. citrinus*; **B:** *C. citratus*; **C:** Salicylic acid. S: control plants; I: plants inoculated with PI; Tr *C. citrinus*, Tr *C. citratus*, Tr Salicylic acid: plants treated respectively with *Callistemon citrinus*, *Cymbopogon citratus* and salicylic acid. Tr + I *C. citrinus*, Tr + I *C. citratus*, Tr + I salicylic acid: Plants treated with *Callistemon citrinus*, *Cymbopogon citratus* and salicylic acid and inoculated with PI respectively.

3.4. Effect of aqueous plant macerates on lipid content

In healthy plants (S), the lipid content on day 0 was $27.5 \pm 2.00 \text{ mg g}^{-1}$ FW. When the plants were inoculated (I), there was a significant decrease in lipid content ($P < 0.05$) on days 2 and 4. The lipid content then increases on day 8 and decreases on day 10. The decrease in lipid content on day 2 was 50.19%. Treatment of the plants with plant extracts resulted in a gradual and significant increase from day 2 with a peak on day 8. The increase at day 8 was more than 100% for all treatments. In the post-treatment inoculated plants (Tr + I), lipid content increased gradually and significantly ($P < 0.05$) by more than 100% from day 2 and peaked on days 10 (*C. citrinus*) and 8 (*C. citratus*). In general, the increase in lipid content was greater in treated and inoculated plants (Tr + I) than in treated plants (Tr) over time (**Figure 3**).




Figure 3: Lipids content (expressed in mg.g^{-1} FW) in leaves of tomato plants on a time course after treatment with plants extracts and inoculation with PI and in the control. Data presented are the means \pm SD of five replicates. Different letters indicate a significant difference ($p < 0.05$) between values, Duncan test. **A:** *C. citrinus*; **B:** *C.citratus*; **C:** Salicylic acid. S: control plants; I: plants inoculated with PI; Tr *C. citrinus*, Tr *C. citrinus*, Tr Salicylic acid: plants treated respectively with *Callistemon citrinus*, *Cymbopogon citratus* and salicylic acid. Tr + I *C. citrinus*, Tr + I *C. citrinus*, Tr + I salicylic acid: Plants treated with *Callistemon citrinus*, *Cymbopogon citratus* and salicylic acid and inoculated with PI respectively.

3.5. Effect of plant extracts on amino acids content

In healthy plants (S), the amino acid content was $8.57 \pm 0.20 \text{ mg.g}^{-1}$ FW on day 0. When the plants were inoculated (I), the amino acid content increased gradually and significantly ($P < 0.05$) with a maximum content at day 10 with increase of 54.18%. After treatment with plant extracts, there was a significant and gradual increase in the amino acid content. From day 2, the percentage increase was 71.63% and 625.00% respectively with *C.*

citrinus and *C. citratus* and it continued until day 10. In plants inoculated after treatment (Tr + I), there was a significant ($P < 0.05$) and gradual increase in amino acid content from day 2. The percentages of increase on day 2 were 71.81% and 55.46% respectively with *C. citrinus* and *C. citratus* treatments. In general, the amino acid content in treated and inoculated plants (Tr + I) was higher than that of treated plants (Tr) (**Figure 4**).




Figure 4: Amino acids content (expressed in mg.g^{-1} FW) in leaves of tomato plants on a time course after treatment with plants extracts and inoculation with PI and in the control. Data presented are the means \pm SD of five replicates. Different letters indicate a significant difference ($p < 0.05$) between values, Duncan test. **A:** *C. citrinus*; **B:** *C. citratus*; **C:** Salicylic acid. S: control plants; I: plants inoculated with PI; Tr *C. citrinus*, Tr *C. citratus*, Tr Salicylic acid: plants treated respectively with *Callistemon citrinus*, *Cymbopogon citratus* and salicylic acid. Tr + I *C. citrinus*, Tr + I *C. citratus*, Tr + I salicylic acid: Plants treated with *Callistemon citrinus*, *Cymbopogon citratus* and salicylic acid and inoculated with PI respectively.

3.6. Effect of plant extracts on proline content

In healthy plants (S), the proline content on day 0 was $1.60 \pm 0.10 \mu\text{g.g}^{-1}$ FW. When the plants were inoculated (I), there was a significant increase in proline content ($P < 0.05$) from day 2 of 145.62%. The treatment of the plants with the plant extracts resulted in a gradual and significant increase in proline content over time with a maximum value at day 10. In the inoculated plants after treatment (Tr + I), the proline content increased gradually and significantly ($P < 0.05$) in the same way as in the treated plants (Tr). In general, the proline content was higher in treated and inoculated (Tr + I) plants than in treated (Tr) plants over time (Figure 5).




Figure 5: Proline content (expressed in $\mu\text{g.g}^{-1}$ FW) in leaves of tomato plants on a time course after treatment with plants extracts and inoculation with PI and in the control. Data presented are the means \pm SD of five replicates. Different letters indicate a significant difference ($p < 0.05$) between values, Duncan test. **A:** *C. citrinus*; **B:** *C. citratus*; **C:** Salicylic acid. S: control plants; I: plants inoculated with PI; Tr *C. citrinus*, Tr *C.*

*citratu*s, Tr Salicylic acid: plants treated respectively with *Callistemon citrinus*, *Cymbopogon citratus* and salicylic acid. Tr + I *C. citrinus*, Tr + I *C. citratus*, Tr + I salicylic acid: Plants treated with *Callistemon citrinus*, *Cymbopogon citratus* and salicylic acid and inoculated with PI respectively.

3.7. Effet des extraits de plantes sur la variation des activités des enzymes antioxydants des feuilles de *L. esculentum*.

III.1.3.1. Ascorbate peroxidase (APX) activity

In healthy plants (S), ascorbate peroxidase activity was $20.7 \pm 1.00 \text{ U.g}^{-1}\text{FW}$ from day 0 and did not vary significantly until day 10. When plants are inoculated (I), APX activity increases significantly ($P < 0.05$) with a peak of activity obtained at day 2. APX activity then progressively decreases until day 10 with values that remain higher than in healthy plants. Treatment of the plants with the plant extracts resulted in a significant ($P < 0.05$) increase in APX activity. With values of $25.00 \pm 2.00 \text{ U.g}^{-1} \text{ FW}$ (*C. citrinus*) and $19.9 \pm 2.00 \text{ U.g}^{-1} \text{ FW}$ (*C. citratus*) at day 0, it increased with a peak at day 2 with 3 and 8 times higher values. It then decreased progressively until day 10 but with higher values than healthy (S) and inoculated (I) plants. When the plants were inoculated after the treatment with the plant extracts (Tr + I), the APX activity increased in the same way as in the treated plants but with higher values (**Figure 6**).




Figure 6: APX activity (expressed in U.g⁻¹ FW) in leaves of tomato plants on a time course after treatment with plants extracts and inoculation with PI and in the control. Data presented are the means \pm SD of five replicates. Different letters indicate a significant difference ($p < 0.05$) between values, Duncan test. **A:** *C. citrinus*; **B:** *C. citrinus*; **C:** Salicylic acid. S: control plants; I: plants inoculated with PI; Tr *C. citrinus*, Tr *C. citrinus*, Tr Salicylic acid: plants treated respectively with *Callistemon citrinus*, *Cymbopogon citratus* and salicylic acid. Tr + I *C. citrinus*, Tr + I *C. citrinus*, Tr + I salicylic acid: Plants treated with *Callistemon citrinus*, *Cymbopogon citratus* and salicylic acid and inoculated with PI respectively.

3.8. Phenylalanine ammonia lyase (PAL) activity

Phenylalanine ammonia lyase (PAL) activity varied from $68.67 \pm 1.53 \mu\text{mol} \cdot \text{min}^{-1} \cdot \text{g}^{-1}$ FW to $73.67 \pm 1.53 \mu\text{mol} \cdot \text{min}^{-1} \cdot \text{g}^{-1}$ FW at day 0 as a function of treatment in healthy plants. When plants were inoculated (I), PAL activity increased significantly ($P < 0.05$) from day 2 to 71.95%. After treatment of the plants (Tr), there was a significant ($P < 0.05$) increase in PAL activity from day 2 of 62.00%, and 61.80% respectively in *C. citrinus* and *C. citratus*. It then

increases gradually and peaks in activity are obtained on day 6. When plants are inoculated after treatment (Tr + I), PAL activity increases in the same way as in treated plants but with higher values. Peaks in activity are obtained at day 6 and this activity remains high and higher than in healthy (S), inoculated (I) and treated (Tr) plants (**Figure 7**).




Figure 7: PAL activity (expressed in $\mu\text{g}^{-1}\text{FW}$) in leaves of tomato plants on a time course after treatment with plants extracts and inoculation with PI and in the control. Data presented are the means \pm SD of five replicates. Different letters indicate a significant difference ($p < 0.05$) between values, Duncan test. **A:** *C. citrinus*; **B:** *C. citratus*; **C:** Salicylic acid. S: control plants; I: plants inoculated with PI; Tr *C. citrinus*, Tr *C. citratus*, Tr Salicylic acid: plants treated respectively with *Callistemon citrinus*, *Cymbopogon citratus* and salicylic acid. Tr + I *C. citrinus*, Tr + I *C. citratus*, Tr + I salicylic acid: Plants treated with *Callistemon citrinus*, *Cymbopogon citratus* and salicylic acid and inoculated with PI respectively.

3.9. Peroxidase (POX) activity

In the leaves of healthy plants (S), the peroxidase activity (POX) varied from 2.87 ± 0.25 U.g⁻¹ FW to 4.33 ± 0.15 U.g⁻¹ FW. When the plants were inoculated (I), it increased on day 2 and reached a peak on day 4 (9.00 ± 1.88 U.g⁻¹ FW). After the treatment of the plants, there was a significant increase ($P < 0.05$) in POX activity. From a value of 2.93 ± 0.21 U.g⁻¹ FP on day 0, it was 2 to 7 times higher from day 2. PAL activity then increased gradually with a peak in activity on day 4. When plants are inoculated after treatment (Tr + I), POX activity increases in the same way as in treated plants but with higher values. A peak in activity is obtained on day 4 with the different treatments. (Figure 8).




Figure 8: POX activity (expressed in U.g⁻¹ FW) in leaves of tomato plants on a time course after treatment with plants extracts and inoculation with PI and in the control. Data presented are the means \pm SD of five replicates. Different letters indicate a significant difference ($p < 0.05$) between values, Duncan test. **A:** *C. citrinus*; **B:** *C. citratus*; **C:** Salicylic acid. S: control plants; I: plants inoculated with PI; Tr *C. citrinus*, Tr *C. citratus*, Tr Salicylic acid: plants treated respectively with *Callistemon citrinus*, *Cymbopogon citratus* and salicylic acid. Tr

+ I *C. citrinus*, Tr + I *C. citratus*, Tr + I salicylic acid: Plants treated with *Callistemon citrinus*, *Cymbopogon citratus* and salicylic acid and inoculated with PI respectively.

3.10. Superoxide dismutase (SOD) activity

In the leaves of healthy plants (S), the SOD activity of $40.10 \pm 0.20 \text{ U} \cdot \text{min}^{-1} \cdot \text{g}^{-1} \text{ FW}$ at day 0 did not vary significantly until day 10. When plants were inoculated (I), SOD activity increased significantly ($P < 0.05$) over time with a maximum value at day 4. After treatment of the plants (Tr), there was a significant ($P < 0.05$) increase in activity with 2-3 times higher values on day 2 depending on the treatment. When plants are inoculated after treatment (Tr + I), SOD activity increases in the same way as in treated plants but with higher values. A peak in activity is obtained on day 4 with the different treatments (Figure 9).




Figure 9: SOD activity (expressed in $U\ min^{-1}\ g^{-1}\ FW$) in leaves of tomato plants on a time course after treatment with plants extracts and inoculation with PI and in the control. Data presented are the means \pm SD of five replicates. Different letters indicate a significant difference ($p < 0.05$) between values, Duncan test. **A:** *C. citrinus*; **B:** *C. citratus*; **C:** Salicylic acid. S: control plants; I: plants inoculated with PI; Tr *C. citrinus*, Tr *C. citratus*, Tr Salicylic acid: plants treated respectively with *Callistemon citrinus*, *Cymbopogon citratus* and salicylic acid. Tr + I *C. citrinus*, Tr + I *C. citratus*, Tr + I salicylic acid: Plants treated with *Callistemon citrinus*, *Cymbopogon citratus* and salicylic acid and inoculated with PI respectively.

4. Discussion

The effect of plant extracts on the tomato defence system against *Phytophthora infestans* (PI) was evaluated through the determination of the content of biomolecules (chlorophyll a, proteins, carbohydrates, lipids, amino acids, proline) in tomato leaves and the evaluation of the activities of antioxidant enzymes (ascorbate peroxidase (APX), phenylalanine ammonia lyase (PAL), peroxidase (POX), Superoxyde dismutase (SOD)).

Our study reveals that when tomato plants are inoculated with PI, the chlorophyll a content has decreased. This decrease is believed to be a consequence of downy mildew, the pathogen releases toxins that trigger the release of ROS. It has been proven that the toxins produced by PI inhibit photosynthesis. The decrease in Chlorophyll a content is also believed to be due to the high level of lipid peroxidation causing cell death in tomato leaf tissues (El-Khallal, 2007). When plants are treated with plant extracts, an increase in Chl a content is noted. This increase could be justified by the stimulating effect of the extracts on the enzymatic activity of ribulose 1,5 bisphosphate carboxylase, a key a key enzyme of photosynthesis (Khodary, 2004).

Protein content decreased in roots under inoculated conditions, but when tomato plants were treated with plant extracts, there was an increase in protein content. This result is in agreement with that obtained by Samir et al., (2019) who showed that tomato plants inoculated with *Fusarium oxysporum* accumulate proteins. According to this author, the increase in protein content could be explained by an increased synthesis of the enzymes involved in detoxification reactions.

When tomato plants were inoculated or treated with plant extracts, there was an increase in total free sugar content. Indeed, plant extracts are at the origin of the physiological notification of tomatoes due to photosynthesis. The accelerating effect of the extracts on photosynthesis will allow an increased synthesis of carbohydrates. Furthermore, it has been shown that cell wall glycoproteins rich in glycine, hydroxyproline, and proline are involved in the response to plant stress (Cassab, 1998). Couée et al., (2006) have shown that in some

plant species, increased synthesis of sugar such as mannitol is linked to increased resistance to oxidative stress.

The different actions carry out on tomatoes, namely PI inoculation, the treatment with plant extracts and the treatment followed by inoculation, resulted in a significant increase in the MDA content, which is a product of the membrane lipids peroxidation. It is a marker of oxidative damage and its concentration indicates that induction of oxidative stress is effective (**Abdelaziz et al., 2022**). The increase in MDA concentrations in plants could be explained by a higher peroxidation of lipids in root cells.

When tomato plants were inoculated or treated with plant extracts, there was an increase in the content of total amino acids and proline particularly. This result is in agreement with the one obtained by **Omokolo and Boudjeko (2005)** in the *Xanthosoma sagittifolium/Pythium myriotylum* interaction. Indeed, the increase in amino acids content after treatment with plant extracts could serve in the synthesis of molecules such as proteins, phenolic compounds and phytoalexins which are molecules involved in defence mechanisms (**Graham et al., 1999**). The particular increase in proline content suggests the preferential orientation of the metabolic pathways of amino acid synthesis pathways towards proline synthesis. **Akladious et al., (2015)** showed that the treatment of tomato plants with salicylic acid under oxidative stress conditions was responsible for the accumulation of proline. Several other authors have shown that plants under various stresses accumulate proline and this accumulation is positively correlated with the tolerance of oxidative stress (**Anjum et al., 2012; Li et al., 2013**).

When plants were inoculated with PI, APX activity increased. Treatment of plants with extracts resulted in a significant increase in APX activity. When plants were inoculated after treatment with plant extracts, APX activity increased in the same way as in treated plants but with higher values. This result is in agreement with the one obtained by **Mbouobda (2013)** in the roots of *Xanthosoma sagittifolium* treated with BTH and CTH and inoculated with *Xanthosoma sagittifolium*. Localized in peroxisomes and chloroplasts in plants, APX uses ascorbate as an electron donor and prevents H₂O₂ from leaving peroxisomes. In addition to being involved in various physiological processes such as photosynthesis, differentiation and growth, it is thought to be involved in stabilising plants under various biotic or abiotic stresses.

After inoculation of the tomato plants, the significant increase in PAL activity in the roots indicates the activation of this crossroads enzyme which would trigger the different defence mechanisms in the plant. When plants were treated with plant extracts and then

inoculated with PI a significant increase in PAL activity is noted. **Shadle et al., (2003)** have shown that PAL activity in tobacco increases after infection, which reduce the incidence of disease by stimulating defence reactions. This confirms the key role of PAL in the synthesis of phenylpropanoids which are molecules involved in the synthesis of secondary metabolites, mainly phytoalexins and salicylic acid which act in reducing the incidence of plant diseases through direct antifungal activity and stimulation of plant defence mechanisms (**Aziz et al., 2006**).

POX activity was higher in plants treated with extracts than those not treated. This would be related to oxidative stress with the involvement of peroxidase activity as an antioxidant response against H₂O₂. For **Samir et al., (2019)**, this stress response involves a homeostatic response and control of the degradation process to maintain the integrity of vital cell functions. **Mandal and Mitra, (2008)** have shown that treatment of *Lycopersicon esculentum* with chitosan induces a significant increase in POX activity, and an increase in the content of phenolic compounds, which protects them against parasitic attacks.

Effective destruction of H₂O₂ requires SOD action. Our results indicate an increase in SOD activity in the roots of tomato plants treated with plant extracts, treated and inoculated. This increase in activity maintains a positive ROS balance in favour of H₂O₂ accumulation. This result is in agreement with the one obtained by **Li et al., (2008)** suggesting an involvement of SOD in the apple defence mechanism at all maturity stages.

CONCLUSION

The treatment of tomato with the aqueous extracts of *Callistemon citrinus* L., *Cymbopogon citratus* (DC) STAFW and *Oxalis barrelieri* L. stimulates its antioxidant defence system against *Phytophthora infestans*. This stimulation is manifested in tomato leaves by an increase in the content of chlorophyll a, proteins, carbohydrates, lipids, amino acids and proline; an increase in ascorbate peroxidase, phenylalanine ammonia lyase, peroxidase and superoxide dismutase activities. The increase in metabolite synthesis on the one hand and the increase in enzyme activities on the other hand could mitigate the oxidative damage that takes place during the expansion of the pathogen. Aqueous macerates of *C. citrinus* and *C. citratus* could be used as natural products to stimulate the tomato defence system against *Phytophthora infestans*.

References

- Abdelaziz A.M., Kalaba,M.H., Hashem,A.H., Sharaf O.H. and Attia M.S. (2022).** Biostimulation of tomato growth and biocontrol of Fusarium wilt disease using certain endophytic fungi. *Botanical studies*, 63, 1-14.
- Abdelkhalek, A., Al-Askar, A. A., Alsubaie, M. M., and Behiry, S. I. (2021).** First report of protective activity of *Paronychia argentea* extract against Tobacco Mosaic Virus infection. *Plant Theory*, 10, 2435.
- Agarwal, P.C., Carmen Mortensen, N. and Mathur, S.B. (1989).** Seed-borne diseases and seed health testing of rice. Technical Bulletin N° 3 and Phytopathological Papers N° 30. Danish Government Institute of Seed Pathology for Developing Countries, Copenhagen, Denmark and CAB International Mycological Institute, Kew, Surrey, United Kingdom, 106p.
- Akladious A., Isaac, G.S. and Abu-Tahon, M.A. (2015).** Induction and resistance against *Fusarium* wilt disease of tomato by using sweet basil (*Ocimum basilicum* L.) extract. *Canadian. Journal of Plant Science*, 95, 689-701.
- Anjum, S.A., Farooq, M., Xie, X.Y., Lie, X.J. and Ijaz, M.F. (2012).** Antioxidant defence system and proline accumulation enable hot pepper to perform better under drought. *Science Horticulture*, 140, 66-73.
- Aziz, T.P., Couderchet, M., Vernet, G. and Aziz, A. (2006).** Chitosan stimulates defence reactions in grapevine leaves and inhibits the development of *Botrytis cinerea*. *European Journal of Plant Pathology*, 114, 405-413.
- Bates, L.S., Waldren, R.P. and Teare, I.D. (1973).** Rapid determination of the free proline in water stress studies. *Plant soil*, 38, 205-208.
- Beauchamp, C. and Fridovich, I. (1971).** Superoxide dismutase: improved assays and an assay applicable to acrylamide gels. *Annal of Biochemistry*, 44, 276-286.
- Bradford, M.M. (1976).** A rapid and sensitive method for the quantification of micrograms of quantities of protein utilizing the principle of protein-dye binding. *Annal of Biochemistry*, 72, 248-254.
- Cassab, G.I (1998).** Plant cell wall proteins. *Annual Review of Plant Physiology and Plant Molecular Biology*, 49, 281-309.
- Couée, I., Sulmon, C., Gouesbet, G. and El Amrani, A. (2006).** Involvement of soluble sugars in reactive oxygen species balance and responses to oxidative stress in plants. *Journal of Experimental Botany*, 57, 449-459.
- Dakole, D.C., Nguetack, J., Dongmo, L.B.J., Galani, Y.J.H., Azah, U.R., Somda, I and Amvam Z.P.H. (2016).** Antifungal potential of essential oils, aqueous and ethanol extracts of

thirteen plants against *Fusarium oxysporum* f. sp *lycopersici* and *Phytophthora infestans* (Mont.) de Bary as major tomato pathogens in Cameroon. *International Journal of Current Research*, 19(2), 128-145.

El- Khallal, S.M. (2007). Induction and Modulation of Resistance in Tomato Plants Against Fusarium Wilt Disease by Bioagent Fungi (Arbuscular Mycorrhiza) And/or Hormonal Elicitors (Jasmonic Acid& Salicylic Acid): Changes in Growth, Some Metabolic Activities and Endogenous Hormones Related to Defence Mechanism. *Australian Journal of Basic and Applied Sciences*, 1(4), 691-705.

Fangue, G.Y., Mouafo R. A., Fomekong K M., Effa O M. P and Djocgoue, P. F. (2021). Allelopathic Effect of Three Wild Plants (*Azadirachta indica*, *Tithonia diversifolia* and *Thevetia peruviana*) on Tomato (*Lycopersicum esculentum* Mill.) Growth and Stimulation of Metabolites Involved in Plant Resistance. *American Journal of Plant Sciences*, 12, 285-299.

FAOSTAT. (2019). Food and Agriculture Organization of the United Nations. <https://faostat3.fao.org/download/Q/QC/E>. Consulted on 10-9-2022.

Foleh, J., Less M., Solune and Stanley, G. (1957). A simple method for the isolation and purification of total lipids from animal tissues. *Journal of Biology and Chemistry*, 226, 497-509.

Graham, T.L., Kim, J.E. and Graham, M.Y. (1990). Role of constitutive isoflavone conjugates in the accumulation of glyceolin in soybean infected with *Phytophthora megakaria*. *Molecular Plant-Microbe interaction*, 6, 157-166.

Hammerschmidt, R., Nuckles, E. and Kuc, J. (1982). Association of enhanced peroxidase activity with induced systemic resistance of cucumber to *Colletotrichum lagenarium*, *Physiology and Plant Pathology*, 20, 73-82.

Kagale, S., Marimuthu, T., Thayumanavan, B., Nandakumar,R. and Samiyappan, R. (2004). Antimicrobial activity and induction of systemic resistance in rice by leaf extract of *Datura metel* against *Rhizoctonia solani* and Xoo. *Physiology Molecular Plant Pathology*, 65, 91-100.

Keatinge, D. (2012). Vegetables: Less visible, but vital for human health-Why nutrient-dense indigenous vegetables must be on the plate for economic development, food security, and health. *AVRDC News Brief*, 31 May 2012.

Khodary, S. (2004). Effect of salicylic acid on the growth, photosynthesis and carbohydrate metabolism in salt stressed maize plants. *International Journal of Agricultural Biology*, 6 (1), 5-8.

- Konje, C.N., Abdulai, A.N., Tange, D.A., Nsobinyui, D., Tarla, D.N. and Tita, M.A. (2019).** Identification and management of pests and diseases of garden crops in Santa Cameroon. *Journal of Agriculture and Ecology Research International*, 18, 1-9
- Li, B., Wei, J., Wei, X., Tang, K., Liang, Y., Shu, K. and Wang, B. (2008).** Effect of sound wave stress on antioxidant enzyme activities and lipid peroxidation of *Dendrobium candidum*. *Colloids and Surfaces. Biointerfaces*, 63, 269-275.
- Li, Z.G., Ding, X.J. and Du, P.F. (2013).** Hydrogen sulfide donor sodium hydrosulfide-improved heat tolerance in maize and involvement of proline. *Journal of Plant Physiology*, 170, 741-747.
- Lichtenthaler, H.K. and Wellburn, A.R. (1983).** Determinations of total carotenoids and chlorophylls a and b of leaf extracts in different solvents. *Biochemical Society Transactions*, 11, 591 - 592.
- Mandal, S. and Mitra, A. (2008).** Accumulation of cell wall-bound phenolic metabolites and their upliftment in hairy root cultures of tomato (*Lycopersicon esculentum* Mill.). *Biotechnology Letters*, 30, 1253–1258.
- Mandal, S., Kar, I., Mukherjee, A.K. and Acharya, P. (2013).** Elicitor-Induced Defence Responses in *Solanum lycopersicum* against *Ralstonia solanacearum*. *The Scientific World Journal*, 1-9.
- Mathur, S.B. and Kongsdal, O. (2003).** Common laboratory seed health testing methods for detecting fungi. (1st Edition). Bassersdorf: International Testing Association, 425p.
- Mbouobda, H.D. (2013).** Stimulation of defence mechanism system by chitosan (CHT) and Benzo-(1,2,3)-thiadiazole-7-carbothionic acid-S-méthyl ester acid (BTH) in *Xanthosoma sagittifolium* (macabo) / *Pythium myriotylum* interaction. PhD thesis. University of Yaounde 1, 142p.
- Mekam, P.N., Martini, S., Nguefack, J., Tagliazucchi, D., Mangoumou, G.N. and Stefani, E. (2019).** Activity of extracts from three tropical plants towards fungi pathogenic to tomato (*Solanum lycopersicum*). *Phytopathologia Mediterranea*, 58 (3), 573-586.
- Moore, S. and Stein, W.H. (1948).** Photometric method for use in the chromatography of amino acids. *Journal of Biology*, 176, 367-388.
- Nakano, Y., and Asada, K. (1981).** Hydrogen peroxide is scavenged by ascorbate-specific peroxidase in spinach chloroplasts. *Plant Cell Physiology*, 22, 867-880.
- Nelson, N. (1944).** A photometric adaptation of Somogyi's method for the determination of reducing sugar. *Analytic Chemistry*, 31, 426-428.

Omokolo, N.D. and Boudjeko, T. (2005). Comparative analyses of alterations in carbohydrates, amino acids, phenols and lignin in roots of five cultivars of *Xanthosoma sagittifolium* infected by *Pythium myriotylum*. *South African Journal of Botany*, 71 (3-4), 432-440.

Samir K B., HG Prakesh, H.G., Ram, P. and Javed B. (2019). Induced synthesis of defence molecules in tomato (*Solanum lycopersicum* L.) against *Fusarium* wilt through plant extracts *Bangladesh Journal of Botany*, 48, 169-175.

Saunders, J.A. and McClure, J.W. (1975). Phytochrome controlled phenylalanine ammonia lyase activity in *Hordeum vulgare* plastids. *Phytochemistry*, 14, 1285-1289.

Shadle, G.L., Wesley, V., Korth, K.L., Chen, F., Lamb, C. and Dixon R.A. (2003). Phenylpropanoid compounds and disease resistance in transgenic tobacco with altered expression of L-phenylalanine ammonia-lyase. *Phytochemistry*, 64, 153-161.

UNDER PEER REVIEW