

## **Original Research Article**

### **EFFECT OF SULPHUR AND ZINC ON GROWTH AND YIELD OF LENTIL (*Lens culinaris* L.)**

#### **ABSTRACT**

A field experiment was conducted during *Rabi* season of 2022 at the Crop Research Farm, Department of Agronomy, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj (U.P.) India. To study the Response of Sulphur and Zinc on growth and yield of Lentil. The treatments consist of Sulphur 20, 30, 40 kg/ha and Zinc 5.0, 6.0, 7.0 kg/ha. The soil of experimental plot was sandy loamy in texture, nearly neutral in soil reaction (pH 7.8), low in organic carbon (0.35%). Results revealed that the higher plant height (37.68 cm), number of nodules (19.25), plant dry weight (18.83 g/plant), crop growth rate (15.7 g/m<sup>2</sup>/day, number of pods/plant (148.77), number of seeds/pod (1.80), 1000 seed weight (25.27 gm), seed yield (1.78 t/ha) and stover yield (3.05 t/ha) were significantly influenced with application of Sulphur 40 kg/ha + Zinc 7.0 kg/ha.

**Keywords:** *Lentil, Sulphur, zinc, growth parameters, and yield attributes.*

**Comment [H1]:** Write it in materials section.

## INTRODUCTION

The second-most significant cold season legume crop in India is lentil (*Lens culinaris*L.) (Ram and Punia, 2018). It has a 1.32 million hectares surface area and a 1.18 million tonnes of production were produced, and each hectare produced 894 kg of output (Directorate of Economics and Statistics, 2020). Uttar Pradesh (38.47%), Madhya Pradesh (29.95%), Bihar (10.26%), and West Bengal (13.88%) are the biggest producing states. A crucial component of the diets of developing nations is lentil. About twice as much protein than cereal is found in lentil. Additionally, it provides every important amino acid that cereal grains typically lack (Elias *et al.*, 1986; Thavarajahet *al.*, 2011). After soybeans, lentils offer the second-highest protein-to-calorie ratio of any type of legume. A person may get a number of critical nutrients from lentil crops, which are abundant in protein (20–30%), minerals (2–5%), and vitamin B9. Especially for iron (Fe), zinc (Zn), and selenium (Se), lentil is generally rich in micronutrients and has the ability to offer appropriate dietary quantities.

Sulphur is a crucial secondary plant nutrient that is essential for several physiological processes in plants, including the production of amino acids (methionine,

cysteine, and cysteine), the synthesis of proteins, and the production of chlorophyll. It engages in the metabolic processes of vitamins (such as biotin and thiamine) and a portion of coenzyme A and pyrophosphate. It also stimulates enzymes. Lack of Sulphur may be the cause of poor fruiting, poor blooming, and stunted development. According to reports, when it comes to Sulphur requirements, pulses come in second place behind oil seeds (Parashar 2022). Increased Sulphur deficit was caused by the use of Sulphur-free fertilizers and the cultivation of high yielding cultivars under intensive cropping systems. Since the lentil crop is farmed with insufficient irrigation, Sulphur treatment may be necessary to improve the water usage efficiency(Aditya 2021).

Intense agricultural activities due to the burgeoning human population have resulted into micronutrient deficiencies in soil. Zinc deficiency is prevalent worldwide, especially in calcareous soils. In India, 51.2 and 19.2% of soils were found to be deficient in Zn and Fe, respectively (Shukla *et al.*, 2021). Micronutrient deficiencies in soils and crops have led to severe consequences, including reduced yield and low micronutrient concentration in crops, thus resulting in micronutrient malnutrition in humans and animals. Worldwide, Zn and Fe deficiencies have

affected one-fifth and one-third of the population, respectively (Chaudhari *et al.*, 2021). Zinc deficiency has been observed mainly in developing countries (Bailey *et al.*, 2012). It is essential for optimum growth and functioning of immune system, enzyme catalyzed biochemical reactions, neurobehavioral development, protein and DNA synthesis (Allen 2006) and its deficiency results in adverse health conditions such as diarrhea, stunted physical and mental growth, loss of appetite, etc. Iron deficiency results into anemia, which globally has affected 40.0 and 42.0% of pregnant women and children, respectively, and even resulted to 20.0% of maternal deaths. Inadequate Fe intake causes the increased mortality of pregnant women and newborns and compromises the immune system.

Keeping these points in view, the present study entitled “**Effect of Sulphur and Zinc on Growth and Yield of Lentil (*Lens culinaris*L.)**” was conducted at Crop Research Farm, Department of Agronomy, Naini Agriculture Institute, Sam Higginbottom University of Agriculture Technology and Sciences, Prayagraj, Uttar Pradesh during *rabi* season of 2022-23. [To study the Response of Sulphur and Zinc on growth and yield of Lentil.](#)

## MATERIALS AND METHODS

The experiment was conducted during *Rabi* 2022, Crop Research Farm, Department of Agronomy, Naini Agriculture Institute, Sam Higginbottom University of Agriculture Technology and Sciences, Prayagraj, Uttar Pradesh. Which is located at 25.24'42''N latitude, 81°50'56'' E longitude and 98m altitude above the mean sea level (MSL). The experiment was conducted in ~~Randomized~~ ~~randomized~~ ~~Block~~ ~~block~~ ~~Design~~ ~~design~~ with 10 treatments each replicated thrice. The plot size of each treatment was 3m x 3m. Factors are three levels of Sulphur (20, 30 and 40 kg/ha) and three levels of Zinc (5, 6 and 7 kg/ha). The S and Zn were supplied, both are applied as basal at the time of sowing. The lentil variety PL-406 was sown on 26 November 2022 by maintaining a spacing of 30cm × 10cm. Harvesting was done taking 1m<sup>2</sup> area from each plot. And from it three plants were randomly selected for recording growth and yield parameters. The treatment details are as follows, T<sub>1</sub> - (Sulphur 20 kg/ha + Zinc 5 kg/ha), T<sub>2</sub> - (Sulphur 20 kg/ha + Zinc 6 kg/ha), T<sub>3</sub> - (Sulphur 20 kg/ha + Zinc 7 kg/ha), T<sub>4</sub> - (Sulphur 30 kg/ha + Zinc 5 kg/ha), T<sub>5</sub> - (Sulphur 30 kg/ha + Zinc 6 kg/ha), T<sub>6</sub> - (Sulphur 30 kg/ha + Zinc 7 kg/ha), T<sub>7</sub> - (Sulphur 40 kg/ha + Zinc 5 kg/ha), T<sub>8</sub> - (Sulphur 40 kg/ha + Zinc 6 kg/ha), T<sub>9</sub> - (Sulphur 40 kg/ha + Zinc kg/ha), T<sub>10</sub>. (N 20

Kg/ha + P 40 kg/ha + K 20 kg/ha) Control. The observations were recorded for plant height, nodules/plant, dry weight, grain yield and stover yield. The data were subjected to statistical analysis by analysis of variance method (Gomez and Gomez, 1976).

## RESULTS AND DISCUSSION

### Growth parameters

#### Plant height (cm):

The significantly higher plant height (37.68 cm) was observed in treatment-9 (Sulphur 40 kg/ha + Zinc 7 kg/ha). However, treatment-8 (Sulphur 40 kg/ha + Zinc 6 kg/ha) was statistically at par with treatment-9 (Sulphur 40 kg/ha + Zinc 7 kg/ha). Significant and higher plant height may be owing due to Sulphur application in the soil might be due to increase availability and uptake of soil nutrients by the crop contributed by Sulphur fertilization. Availability of Sulphur to plants has helped in better development and thickening of xylem and collenchyma tissue which might have resulted into increased plant height. Similar results are reported with Teja et al., (2021). Further, increase in plant height might be due to with the application of zinc activation of several enzymes which are zinc dependent viz. carbonic anhydrase, ribulose biphosphate

carboxylase, aldolase fructose 1 to 6 bisphosphatase, starch synthetase and sucrose synthetase. Zinc is known to produce the growth hormones and precursor of auxin i.e., tryptophan (Pramanik and Bera A.K. (2012) and Debnath et al. (2018).

#### Nodules/plant:

The significantly higher number of nodules per plant (7.93 cm) was observed in treatment-9 (Sulphur 40 kg/ha + Zinc 7 kg/ha). However, treatment-8 (Sulphur 40 kg/ha + Zinc 6 kg/ha) was statistically at par with treatment-9 (Sulphur 40 kg/ha + Zinc 7 kg/ha). The increase of number of root nodules with increasing level of Zn might be because Zn help to improve more nodulation and leg hemoglobin formation. Proper nutrition of plants with S increases the amount of glucose flowering to the roots and ATP biosynthesis. These are in conformity with the present findings of Valenciano et al. (2011), Singh et al. (2012).

#### Plant dry weight (g):

The significantly higher plant dry weight (18.83 gm/plant) was observed in treatment-9 (Sulphur 40 kg/ha + Zinc 7 kg/ha). However, treatment-8 (Sulphur 40 kg/ha +

Zinc 6  
kg/ha) was statistically at par with treatment-  
9 (Sulphur 40 kg/ha + Zinc 7 kg/ha).

The significantly higher plant dry weight (18.83 gm/plant) was observed with the application of Sulphur. Sulphur plays a major role in photosynthesis which has a direct bearing on plant growth and development, increasing levels of Sulphur application resulted in an increment in dry weight of lentil, the results were found to be similar with **Singh et al. (2000)**. Further, increase in dry matter might be due to with the application of Zinc. Zinc is an important element for the synthesis of tryptophan, which is the precursor for the synthesis of IAA (Indole acetic acid), a growth hormone, involved in stem elongation (**Patel et al., 2007; Shahram and Gholamreza 2012**). It is also considered to be a precursor for auxin synthesis, involved in nitrogen metabolism and several oxidation-reduction reactions, stability of RNA and starch formation. Thus, its suitable supply effects in higher dry matter production, ultimately growth and development of plants. Parallel results were found by **Meena et al., (2012)**.

#### **Crop Growth Rate (g/m<sup>2</sup>/day):**

The significantly higher crop growth rate (15.7 g/m<sup>2</sup>/day) was observed in treatment-

9 (Sulphur 40 kg/ha + Zinc 7 kg/ha). However, treatment-8 (Sulphur 40 kg/ha + Zinc 6 kg/ha) was statistically at par with treatment-9 (Sulphur 40 kg/ha + Zinc 7 kg/ha). Zinc plays a major role in photosynthesis, enzymes activation, fertilization and translocation of assimilates which are responsible for the increase in seed yield. The better crop growth and development might be due to the combined application of Zinc. The results of the present investigation are in close conformity with the findings of **Tiwari et al. (2018)**.

#### **YIELD AND YIELD ATTRIBUTES**

##### **Number of Pods/plant:**

The significant and higher number of pods/plant (148.77) were observed in treatment-9 with (Sulphur 40 kg/ha + Zinc 7 kg/ha), which was significantly superior over the rest of the treatments. However, treatment-8 (Sulphur 40 kg/ha + Zinc 6 kg/ha) was found to be statistically at par with treatment-9 (Sulphur 40 kg/ha + Zinc 7 kg/ha). The significant and higher number of pods/plant (148.77) were recorded with the application of Sulphur and zinc. Improved availability of Sulphur and favorable nutritional environment might have helped the plants at the peak growth period and flowering stages which ultimately increased the number of

Pods per plant (**Mourya et al., 2021**). Further, increase in pods per plant might be due to application of Zinc. Zinc has a greater role in the production of auxin and indole acetic acid, which helps in increased plant growth which resulted in more pods per plant. Similar results were reported by **Upadhyay (2016)**.

#### **Number of seeds/pod:**

The significant and higher number of seeds/pod (1.80) were observed in treatment-9 with (Sulphur 40 kg/ha + Zinc 7 kg/ha), which was significantly superior over the rest of the treatments. However, treatment-8 (Sulphur 40 kg/ha + Zinc 6 kg/ha), was found to be statistically at par with treatment-9 (Sulphur 40 kg/ha + Zinc 7 kg/ha). Number of seeds per pod had increased with application of Sulphur, which leads to transfer of photosynthates and its accumulating from growing parts of plant to seeds which make them plump and bold and also affects the seed size and weight. These results were close with **Choubey et al., 2013**, and also with the application of zinc has also significant improvement was observed might be due to Zinc improved translocation of photosynthates towards reproductive system and thereby enhancing the yield of the crop. Better photosynthetic activity also may have resulted in better translocation of photosynthates from source

to sink due to less crop competition between the plants which might have led to higher yield attributes. These findings are similar to those reported by **Reddy and Ahlawat (1996)**.

#### **Test Weight (g):**

The significant and higher Test weight (25.27 gm) was observed in treatment-9 with (Sulphur 40 kg/ha + Zinc 7 kg/ha), which was significantly superior over the rest of the treatments. However, treatment-8 (Sulphur 40 kg/ha + Zinc 6 kg/ha), was found to be statistically at par with treatment-9 (Sulphur 40 kg/ha + Zinc 7 kg/ha).

The Sulphur application increased the test weight significantly up to 40 kg S ha<sup>-1</sup> over control. This improvement in the growth and yield attributing characters might be due to lowering soil pH with elemental Sulphur addition which was slightly on higher side and improving physical condition of the soil (**Choudhary and Das 1996**). The results so obtained get support with those of **Deo and Khaldelwal (2009)**.

#### **Seed Yield (t/ha):**

The significant and higher seed yield (1.78 t/ha) was observed in treatment-9 with (Sulphur 40 kg/ha + Zinc 7

kg/ha), which was significantly superior over the rest of the treatments. However, treatment-8 (Sulphur 40 kg/ha + Zinc 6 kg/ha), was found to be statistically at par with treatment-9 (Sulphur 40 kg/ha + Zinc 7 kg/ha). The synergistic effect of Sulphur and Zinc may be due to utilization of large quantities of nutrients through their well-developed root system and nodules which might have resulted in better plant development and ultimate yield at low initial status of available Sulphur and Zinc content in the experimental soil. The seed yield of lentil is a function of the product of number of pods per plant, number of seeds per pod and test weight. The increase in the number of pods per plant and number of seeds per pod increased the seed yield of lentil. Similar results are in conformity with **Teja et al. (2021)**. Increase in these attributes due to the involvement of the zinc in enzyme activation, membrane integrity, chlorophyll formation, stomatal balance and starch utilization at early stages which enhanced accumulation of assimilate in the grains resulting in heavier grains. These results are in agreement with the findings of **Krishna et al., (2022)**.

#### **Stover Yield (t/ha):**

The significant and higher stover yield (3.05 t/ha) was observed in treatment-9 with (Sulphur 40 kg/ha + Zinc 7

kg/ha), which was significantly superior over the rest of the treatments. However, treatment-8 (Sulphur 40 kg/ha + Zinc 6 kg/ha), was found to be statistically at par with treatment-9 (Sulphur 40 kg/ha + Zinc 7 kg/ha). The increase in straw yield due to Sulphur application might be attributed to its low availability in experimental soils. The synergistic effect of S may be due to utilization of large quantities of nutrients through well-developed root system which might have resulted in better plant development and ultimate yield at low initial status of available Sulphur in the experimental soil. Similar results reported by **Sahay et al., (2015)**.

#### **Harvest Index (%):**

The significant and higher Harvest Index (37.17 %) was observed in treatment-8 with (Sulphur 40 kg/ha + Zinc 6 kg/ha), which was significantly superior over the rest of the treatments. However, treatment-9 (Sulphur 40 kg/ha + Zinc 7 kg/ha), was found to be statistically at par with treatment-8 (Sulphur 40 kg/ha + Zinc 6 kg/ha). The significant and higher Harvest Index (37.17 %) was observed with the application of Sulphur 40 kg/ha. Increase in harvest index due to better translocation of photosynthesis from growing parts to storage parts which increases the economical yield of the plant. These results are supported by **Chaubey et**

al., 2019 and Shukla et al., 2014.

## CONCLUSION:

It is concluded that with the application of Sulphur 40 kg/ha along with Zinc 7.0 kg/ha (Treatment-9), has performed positively and improves growth and yield parameters.

## REFERENCES:

- Aditya Delu, BS Brar (2021). Effect of different level of Sulphur on growth and yield of lentil crop. *International Journal of Chemical Studies*;9(1):193-197.
- Amit Kumar Tiwari, Ved Prakash, Atik Ahmad and Singh RP (2018). Effect of Biofertilizers and Micronutrients on Nutrient Uptake, Growth, Yield and Yield Attributes of Lentil (*Lens culinaris* L.). *International Journal of Current Microbiology and Applied Sciences*;7(02): 3269-3275.
- Allen, L.; de Benoist, B.; Dary, O. (2006); Hurrell, R. Guidelines on Food Fortification with Micro-nutrients. Available online: <https://www.who.int/nutrition>.
- BagadiMouryaTeja, Vikram Singh and Shruti G George (2021). Effect of Sulphur and zinc on growth and yield of lentil (*Lens culinaris* M.). *The Pharma Innovation Journal*; 10(11): 370-372.
- BanothMurali Krishna, H Sai Kumar, Guguloth Priyanka, Malavath Vinod Naik and Umesha C. (2022). Influence of boron and zinc on growth and yield of green gram (*Vignaradiata* L.). *The Pharma Innovation Journal*; 11(3): 1674-1678.
- Bailey, R.L.; West, K.P.; Black, R.E (2015). The epidemiology of global micronutrient deficiencies. *Annals of Nutrition and Metabolism*.,66:22–33.
- Choubey, S.K., Dwivedi, V.P., & Srivastava, N.K., (2013). Effect of different levels of phosphorus and Sulphur on growth and yield and quality of lentil (*Lens culinaris* M). *Indian Journal of Scientific Research*, 4(2), 149-150.
- Choudhary, H.P. and Das, S.K. (1996). Effect of P, S and Mo application on yield of rainfed black gram and their residual effect on safflower and soil and water conservation in eroded soil *Journal of Indian Society of Soil Science*., 44: 741-745.
- Debnath P, Patta naik SK, Sah D, Chandra

**Comment [H2]:** Please, Unify the format (writing the references)

G, Pandey AK (2018). Effect of boron and zinc fertilization on growth and yield of cowpea (*Vigna unguiculata* Walp.) in Inceptions of Arunachal Pradesh. *Journal of the Indian Society of Soil Science.*;66(2):229-234.

Deo, Chandra and Khaldewal, R.B. (2009). Effect of P and S nutrition on yield and quality of chickpea (*Cicer arietinum* L.). *Journal of Indian Society of Soil Science.*;57: 352-356.

Directorate of Economics and Statistics (2020). Agricultural Statistics at a Glance 2019-20 Government of India Ministry of Agriculture and Farmers Welfare Department of Agriculture, Cooperation and Farmers Welfare Directorate of Economics and Statistics.

Elias, S. M., Hossain, M. S., Sikder, F. S., Ahmed, J., & Karim, M. R. (1986). Identification of constraints to pulse production with special reference to present farming systems. *Annual Report of the Agricultural Economics Division, BARI, Joydebpur*, p.1

Gomez, K. A and Gomez, A. A (1976). Three or more factor experiment. In: Statistical Procedure for

Agricultural Research 2nd edition, p. 139 -141.

Krishna Reddy, S. V., & Ahlawat, I. P. S. (1996). Growth and yield response of lentil cultivars to phosphorus, zinc and biofertilizers. *Journal of Agronomy and Crop Science.* 177(1), 49-59.

Meena, S. N., Jain, K. K., Prasad, D., & Ram, A. (2012). Effect of nitrogen on growth, yield and quality of fodder pearl millet (*Pennisetum glaucum*) cultivars under irrigated condition of North-Western Rajasthan. *Annals of Agricultural Research*, 33(3), 183-188.

Neha Sahay, S. P. Singh, Javed Ali and Yogesh Kumar Sharma (2015) Effect of cobalt and Sulphur nutrition on yield, quality and uptake of nutrients in lentil. *Legume Research*, 38 (5) 2015: 631-634.

Patel, A. S., Sadhu, A. C., Patel, M. R., & Patel, P. C. (2007). Influence of nitrogen and zinc on maize (*Zea mays* L.). *Forage Research*, 34(4): 209-212.

Parashar A, Tripathi L (2020). Effect of phosphorus and Sulphur on the growth and yield of black gram (*Vigna mungo* L.). *Journal of*

Comment [H3]: Different ref. (in context Debnath et al. (2018).

Comment [H4]: Unify the format

Comment [H5]: 2020 or 2022

*Pharmacognosy and Phytochemistry*;9(5):2585-2588.

Ram, B., & Punia, S. S. (2018). Effect of seed priming and foliar urea spray on yield and economics in lentil (*Lens culinaris*) under rainfed condition. *International Journal of Agricultural Sciences.*, 10: 5801–5803.

Shukla, A.K.; Behera, S.K.; Prakash, C.; Tripathi, A.; Patra, A.K.; Dwivedi, B.S.; Trivedi, V.; Rao, C.S.; Chaudhari, S.K.; Das, S.; et al (2021). Deficiency of Phyto-available Sulphur, zinc, boron, iron, copper, and manganese in soils of India. *Scientific Reports.*,11, 19760.

Shukla, A.K., & Singh, N. (2014). Performance of lentil (*Lens culinaris*) varieties to different levels of Sulphur under rainfed conditions district of Chitrakoot U.P. *Trends in Bio Sciences*,7(14), 1677-1678.

Singh, R., Singh, S.B., Manhas, S.S. and Kumar A. 2012. Effect of different levels of Sulphur and varieties on growth, yield and quality of Indian mustard. *International Journal of Plant Science* 7(2):290-294

Singh, Y. P., Chauhan, C. P. S. and Gupta, R. K. 2000. Effect of Sulphur,

Phosphorous and inoculation on growth, yield and Sulphur utilization by lentil (*Lens culinaris*). *Indian Journal of Agricultural Sciences* 70(7); 491-493.

Upadhyay RG, Singh A (2016). Effect of nitrogen and zinc on nodulation, growth and yield of cowpea (*Vigna unguiculata*). *Legume Research-An International Journal*; 39(1):149-151.

Valenciano, J.B., Boto, J.A. and Marcelo, V. [2010]. Response of chickpea (*Cicer arietinum* L.) yield to zinc, boron and molybdenum application under pot conditions. *Spanish Journal of Agricultural Research* 8(3): 797-807.

Wessells, K.R.; Brown, K.H (2012). Estimating the global prevalence of zinc deficiency: Results based on zinc availability in national food supplies and the prevalence of stunting. *PLoS ONE*.

Comment [H6]: Unify the format

Comment [H8]: 2010 or 2011

Comment [H7]: Reorder 2014 before 2021

Comment [H9]: Not present in context

**Table 1. Influence of Sulphur and Zinc on growth parameters of lentil.**

| S. No. | Treatment combinations         | Plant height<br>(cm) | Number of nodules/plants | Plant Dry weight<br>(%) | Crop growth rate<br>(g/m <sup>2</sup> /day) |
|--------|--------------------------------|----------------------|--------------------------|-------------------------|---|
| 1.     | Sulphur 20 kg/ha + Zinc 5kg/ha | 33.24                | 15.77                    | 15.80                   | 13.4  |
| 2.     | Sulphur 20 kg/ha + Zinc 6kg/ha | 34.10                | 16.44                    | 16.34                   | 13.5  |
| 3.     | Sulphur 20 kg/ha + Zinc 7kg/ha | 34.35                | 17.03                    | 16.73                   | 14.2  |
| 4.     | Sulphur 30 kg/ha + Zinc 5kg/ha | 35.06                | 17.59                    | 16.31                   | 14.6  |
| 5.     | Sulphur 30 kg/ha + Zinc 6kg/ha | 35.43                | 18.19                    | 16.45                   | 15.8  |
| 6.     | Sulphur 30 kg/ha + Zinc 7kg/ha | 35.77                | 17.78                    | 17.14                   | 16.3  |
| 7.     | Sulphur 40 kg/ha + Zinc 5kg/ha | 36.32                | 18.72                    | 17.47                   | 15.6  |
| 8.     | Sulphur 40 kg/ha + Zinc 6kg/ha | 37.27                | 18.61                    | 18.48                   | 15.4  |
| 9.     | Sulphur 40 kg/ha + Zinc 7kg/ha | 37.68                | 19.25                    | 18.83                   | 15.7  |
| 10.    | Control (NPK 20-40-20kg/ha)    | 33.64                | 16.50                    | 15.07                   | 13.2  |
|        | F test                         | S                    | S                        | S                       | S   |
|        | S Em.(±)                       | 0.45                 | 0.37                     | 0.13                    | 0.33  |
|        | CD<br>(p=0.05)                 | 1.32                 | 1.10                     | 0.38                    | 0.97  |

**Table 2. Influence of Sulphur and Zinc on yield attributes and yield of lentil.**

| S. No. | Treatments                     | Pods/plant | Seeds/pod | Test weight (g) | Seed yield (t/ha) | Stover yield (t/ha) | Harvest index (%) |
|--------|--------------------------------|------------|-----------|-----------------|-------------------|---------------------|-------------------|
| 1.     | Sulphur 20 kg/ha + Zinc 5kg/ha | 106.83     | 1.29      | 21.97           | 1.29              | 2.55                | 33.64             |
| 2.     | Sulphur 20 kg/ha + Zinc 6kg/ha | 111.27     | 1.35      | 22.18           | 1.34              | 2.66                | 33.56             |
| 3.     | Sulphur 20 kg/ha + Zinc 7kg/ha | 117.30     | 1.50      | 23.12           | 1.42              | 2.72                | 34.27             |
| 4.     | Sulphur 30 kg/ha + Zinc 5kg/ha | 121.17     | 1.53      | 22.18           | 1.46              | 2.72                | 34.96             |
| 5.     | Sulphur 30 kg/ha + Zinc 6kg/ha | 125.54     | 1.61      | 22.71           | 1.55              | 2.80                | 35.70             |
| 6.     | Sulphur 30 kg/ha + Zinc 7kg/ha | 131.60     | 1.67      | 24.18           | 1.64              | 2.82                | 36.75             |
| 7.     | Sulphur 40 kg/ha + Zinc 5kg/ha | 136.83     | 1.64      | 23.00           | 1.57              | 2.69                | 36.79             |
| 8.     | Sulphur 40 kg/ha + Zinc 6kg/ha | 142.54     | 1.72      | 24.21           | 1.68              | 2.84                | 37.17             |
| 9.     | Sulphur 40 kg/ha + Zinc 7kg/ha | 148.77     | 1.80      | 25.27           | 1.78              | 3.05                | 36.95             |
| 10.    | Control (NPK 20-40-20kg/ha)    | 118.73     | 1.38      | 20.63           | 1.26              | 2.70                | 31.87             |
|        | F-Test                         | <b>S</b>   | <b>S</b>  | <b>S</b>        | <b>S</b>          | <b>S</b>            | <b>S</b>          |
|        | SEm.±                          | 2.19       | 0.04      | 0.69            | 0.03              | 0.05                | 0.67              |
|        | CD (p=0.05)                    | 6.50       | 0.11      | 2.05            | 0.10              | 0.15                | 1.98              |