

Original Research Article

Influence of Nitrogen levels and Biofertilizers on the growth and yield of French bean (*Phaseolus vulgaris* L.)

ABSTRACT

The field experiment entitled "Influence of Nitrogen levels and Biofertilizers on the growth and yield of French bean" was conducted during *rabi* season, 2022 at Crop Research Farm in the Department of agronomy, Naini Agriculture Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj Uttar Pradesh. The treatment consisted of three levels of Nitrogen (75, 100 and 125 kg/ha), Biofertilizers (*Rhizobium*, PSB and *Rhizobium*+PSB) and control. The experiment was layout in Randomized Block Design (RBD) with 10 treatments and replicated thrice. Application of *Rhizobium* and PSB along with Nitrogen 100 kg/ha produces higher plant height (46.19cm), higher dry weight (26.20g), maximum number of pods/plant (19.73), maximum number of seeds/pod (5.08), seed yield (22.47 q/ha) and stover yield (50.08 q/ha).

Comment [KS1]: Abstract is too brief

Keywords: French bean, Nitrogen, Biofertilizer, growth and yield.

Introduction

French bean (*Phaseolus vulgaris* L.) is one of the most important commercially grown pulse crop globally. This crop is well adapted to diverse climatic conditions ranging from tropical to temperate region. The French bean (*Phaseolus vulgaris* L.) is a short duration high yielding legume plant of family Fabaceae known by common bean or kidney bean and was considered to have been introduced from Ethiopia in the 16th century by the Portuguese. It is one of the most important commercially grown legume crops during winter season in India and it can be used both as pulse and vegetable. Seed of French bean is highly nutritious containing 22.90 percent protein, 1.20 percent fat and 60.60 percent carbohydrates and minerals viz. calcium (260 mg/100 g of seed), phosphorus (101 mg/100 g of seed) and iron (5.80 mg/100 g of seed). It is assumed that beans can play an important role in the enhancement of the level of nourishment because this plant

is rich in elements like potassium, calcium, iron and phosphorus and it contains 18-32 percent of protein in every grain averagely. The plant is also rich in vitamins like A, B and D etc.

French bean is considered as a nitrogen responsive crop due to its high responsiveness to fertilizer while it is noticed that increased nutrition has a negative impact on yield. Nitrogen is so vital in every crop because it is a major component of chlorophyll, amino acids and the building blocks of proteins. Without proteins, plants wither and die. Application of N increases leaf N and chlorophyll content. Leaf N, chlorophyll and photosynthetic rate are closed related and high leaf N and chlorophyll increase the photosynthetic rate. Therefore, application of N increases the leaf area and photosynthetic rate and thus increases dry matter production. Prolonged use of chemical fertilizers degrades the soil health and affects crop yield. Application of higher doses of nitrogen especially for seed crop of French bean is imperative for realizing its potential yield (Sardana *et al.*, 2000). There is also report that French bean is in sufficient in trapping atmospheric nitrogen due to lack of nodulation in north Indian plains (Kushwaha, 1994). Therefore, it requires large quantity of nitrogenous fertilizer. Increasing cost of inorganic fertilizers and reduction in soil health with chemical Fertilizers there is a need of present are to use of eco-friendly inputs like vermicompost, bio fertilizer viz., Phosphorus solubilizing bacteria, Rhizobium along with inorganic fertilizers.

Biofertilizer required to restore the fertility of the soil. Prolonged use of chemical fertilizers degrades the soil and affects crop yield. Biofertilizers, on the other hand enhance the water holding capacity of the soil and add essential nutrients such as nitrogen, vitamins, and proteins to the soil. Microbes in biofertilizers provide atmospheric nitrogen directly to plants. They aid in solubilisation and mineralisation of other plant nutrients like phosphates. Better synthesis and availability of hormones, vitamins, auxins, and other growth - promoting substances improves plant growth. The interaction of biofertilizer along chemical fertilizer the requirement of inorganic fertilizer will reduced will be better for ecosystem, as the use of fertilizer is one most important factor to increase all growth parameters and yield of crop (Kumar *et al.* 2014).

Comment [KS2]: Delete, repetition

Comment [KS3]: Revise the introduction with latest references with proper justification for taking up the trial

2. MATERIALS AND METHODS

This experiment was laid out during the *Rabi* season of 2022 at Crop Research Farm, Department of Agronomy, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj (U.P.). The crop research farm is

situated at 25° 39' 42" N latitude, 81° 67' 56" E longitude and at an altitude of 98 m above mean sea level. The experiment was laid out in Randomized Block Design Which consisting of ten treatments with T₁-Nitrogen 75kg + *Rhizobium*, T₂-Nitrogen 75 kg + PSB, T₃- Nitrogen 75 kg + *Rhizobium*+ PSB, T₄ - Nitrogen 100 kg + *Rhizobium*, T₅Nitrogen 100 kg + PSB, T₆ - Nitrogen 100 kg + *Rhizobium*+ PSB, T₇ - Nitrogen 125 kg + *Rhizobium*, T₈ - Nitrogen 125 kg + PSB, T₉- Nitrogen 125 kg + *Rhizobium*+ PSB, T₁₀ - Control (NPK 100-80-60 kg/ha).The soil in the experimental area was sandy loam with pH (8.0), Organic Carbon (0.42%), Available N (180.58 kg/ha), Available P (15.54 kg/ha), and Available K (198.67 kg/ha). Seeds are sown at a spacing of 45×15cm² to a seed rate of 80 kg/ha. The recommended dose of phosphorus (80 kg/ha), potassium (60 kg/ha) and Biofertilizer and Nitrogen were applied as per the treatments. Data recorded on different aspects of crop, viz., growth, yield attributes were subjected to statistically analysis by analysis of variance method. (Gomez and Gomez, 1976) and economic data analysis mathematical method.

Comment [KS4]: delete

Comment [KS5]: is it per ha or acre, what is dose of bio fertilizers and how was it applied

Comment [KS6]: delete, experimental soil

Comment [KS7]: express in g/kg

Comment [KS8]: delete , was

Comment [KS9]: seed and stover yield

RESULT AND DISCUSSION:

Growth parameters

In the present investigation growth parameters were found to be maximum for plant height(cm), plant dry weight(g) and crop growth rate (g/m²/day) in treatment 6 [Nitrogen 100 kg + *Rhizobium*+ PSB] which was followed by treatment 9 [Nitrogen 125 kg + *Rhizobium*+ PSB]. Least growth parameters were observed in treatment 1 [Nitrogen 75kg + *Rhizobium*]. The increase was found to be rapid between 40-80 as compared to 20-40. However maximum plant height(46.19 cm), plant dry weight (26.20 g) and crop growth rate (8.22g/m²/day) was recorded at 80 DAS in treatment 6 [Nitrogen 100 kg + *Rhizobium*+ PSB] and least was recorded in treatment 1[Nitrogen 75kg + *Rhizobium*] of 37.14 cm plant height, 21 g dry weight and 6.13 g/m²/day crop growth rate. The findings showed that the trend of increase in plant growth parameters were observed up to treatment 6 beyond that negative effect observed with increased nutrition. The plant growth parameters declined significantly at 125 kg/ha than 100 kg N/ha due to incomplete assimilation of nitrogen. When a nutrient is present in the soil in excess of plant requirement, the nutrient is absorbed in higher amounts which causes imbalance of nutrients or disorder in physiological processes (Reddy &Reddi, 2005). So, it might had affected on the conversion of NH₃ into glutamic acid in plants resulting in the retardation of crop growth character like plant height at maximum level (125

Comment [KS10]: no supporting data

kg/ha) of nitrogen. Similar effect was recorded in total dry matter production per plant (Table 1) at 80 DAS resulting in lower values of yield attributes (table 2). Moreover, according to Moniruzzaman *et al.* (2008), amount of nitrogen beyond optimum level brings about nutrient imbalance and suppresses growth of the plant. Growth inhibition has been attributed to various factors such as toxic effects of free ammonia and carbohydrate limitation (Cramer & Lewis, 1993) due to excessive consumption of soluble sugars for NH₄⁺ assimilation (detoxification) (Walch-Liu *et al.*, 2000). Ammonium induced inhibition of shoot growth has been reported for *Phaseolus vulgaris* L. (Chaillouet *et al.* 1986). It is due to uncoupling of photophosphorylation, lack of carbohydrate and impairment of water status (Walch-Liu *et al.*, 2000). Increase in plant growth parameters as compared to other lower nutrient levels might be due to the increase in nitrogen availability which might have promoted photosynthesis resulting in more assimilation of food material in plant. Similar results were obtained by Rajput *et al.* (2006) and Ramana *et al.* (2011).

The interaction effect of biofertilizers along with inorganic nitrogen was significant for growth parameters. Significant increase in plant height and plant dry weight with *Rhizobium* and PSB as compared to control might be due to increase in uptake of N and P by the plants, which might be due to more N-fixation and P-solubilisation through micro-organisms. Results were similar to (Ramana *et al.*, 2010) in French bean and (Singh *et al.*, 2018) in chickpea.

Yield attributes and Yield

As depicted in table 2 the statistically analysed data pertaining to pods/plant, seeds/pod, seed yield, stover yield indicated that significant differences observed among the treatments. Treatment 6 [Nitrogen 100 kg + *Rhizobium*+ PSB] recorded maximum pods/plant, seeds/pod, seed yield, stover yield (19.70, 5.08, 22.47 and 50.08) and least was observed in treatment 1 [Nitrogen 75kg + *Rhizobium*].

significantly higher pods per plant was found at Treatment 6 [Nitrogen 100 kg + *Rhizobium*+ PSB] (19.73.) and all the remaining treatments were significantly less. Increased pods/plant might be due to increased assimilation of nitrogen and phosphorus which resulted from positive interaction between fertilizer and biofertilizers at 100 kg N/ha. Beyond this level negative effect observed. Verma and Saxena (1995) also reported that pods per plant of French bean increased significantly with the increase in

Comment [KS11]: recast as follows
There was significant differences in growth parameters among treatments due to application of nitrogen at different rates in the presence of biofertilizers (table 1)
All the growth parameters increased with N levels upto 100 kg N/ha and declined at 125 kg N/ha
Combined application of PSB and rhizobium recorded higher growth parameters compared to individual application in the presence of N
The tallest plant (46.19 cm), DMP (26.20 g/plant) and CGR (8.20) with N @ 100 kg/ha plus PSB + rhizobium and was comparable with T9, T8 and T3 but significantly superior to rest of the treatments
The best treatment caused 14.9%, 14.9% and 8.3% increase in plant height, DMP and CGR over control. However, T1 and T2 recorded lower growth parameters compared to RDF.
Explain why
Support your result with latest references

Comment [KS12]: no question of interaction

Comment [KS13]: it should be supported from your data

Comment [KS14]: delete and write as T6

nitrogen level up to 120 kg/ha. Similar findings of the pods per plant of French bean was significantly higher at the nitrogen level of 120 kg/ha as compared to 80, 40 and 0 kg/ha in 1993 (Waniet *al.*, 1998).

The seeds per pod was found to increase with the level of nitrogen from 0 to 100 kg/ha and then declined at 125 kg/ha (Table 2). The maximum number of grains per pod (5.08)

was obtained with 100 kg N/ha at treatment 6 [Nitrogen 100 kg + *Rhizobium*+ PSB] and minimum (3.87) in treatment 1 [Nitrogen 75kg + *Rhizobium*]. This is in conformity with the results obtained by Singh and Verma (2002) that grains per pod increased significantly with increasing level of nitrogen up to 120 kg/ha and beyond that negative impact observed. These findings are in agreement with Saxena and Verma (1995) that the increase in the nitrogen level from 0 to 120 kg/ha increased grains per pod significantly in French bean in 1989/90 and 1990/91. As seed is treated with biofertilizers better result was observed with recommended nitrogen level.

Increased seed yield and stover yield observed in treatment 6 [Nitrogen 100 kg + *Rhizobium*+ PSB] (22.47 q/ha and 50.08 q/ha) and least in treatment 1 [Nitrogen 75kg + *Rhizobium*] (16.00 q/ha and 43.55 q/ha). Increased yield by the biofertilizer interaction with fertilizer could be due to the greater availability of nutrients in the soil and better nodulation under the influence of inoculation resulting in better growth and development which might be attributed to better mobilization of important nutrients and increased allocation of photosynthates towards the economic parts and also hormonal balance on the plant system. This findings corroborate with the findings of Menaria *et al.* (2004) in soybean, Naagare *et al.* (2004) in cluster bean, Rajput *et al.* (2004) in Garden pea and Jain *et al.* (1999) in chick pea. Increase in yield by dual inoculation of *rhizobium* and PSB might be due to increased number of leaves and leaf area which determines the photosynthetic efficiency of plants, dry matter production and ultimately the yield.

CONCLUSION:

Based on the above findings it can be concluded that French bean with the application of *Rhizobium* and PSB along with the application of Nitrogen 100 kg/ha (Treatment 1) recorded highest plant height, dry weight, no. of pods/plant, no. of seeds/pod, seed yield and stover yield.

Comment [KS15]: Application of 100kg N/ha plus PSB and rhizobium significantly increased number of pods/plant over all the treatments except T3 and T9. The best treatment caused 28.68% increase over control. Similarly, T6 recorded maximum no of seeds / plant and significantly superior rest of treatments except of T9. The best treatment caused 16.25% increase over control. Percent increase in seed yield ranged from 4.7 (T4) to 28.13 % (T6) over control. However, T1 and T2 caused reduction in seed yield over control. Application of 100 kg N/ha Plus PSB and rhizobium recorded the highest seed yield (22.47 q/ha) and was significantly superior to rest of the treatments except T9. However, T9, T7, T8 and T3 was on par, T4, T5, T7 and T8 was on par and T1 and T2 was on par. Percent increase in Stover yield ranged from 0.31% (T5) to 9.21% (T6). Similarly, T1 and T2 caused reduction in Stover yield over control to the tune of 5.04 and 2.10, respectively. Application of 100kg N/ha plus PSB and rhizobium recorded maximum Stover yield (50.08 Q/ha). It was significantly superior to rest of the treatments except T9. However, T9 was on par with T3, T4, T7, T8 and T5 and T1 and T2 is also on par. Discussion need to be more strong with latest literature.

Table: 1 Influence of Nitrogen levels and Biofertilizer on growth of French bean.

S.No.	Treatment combinations	At 80 DAS		
		Plant height (cm)	Dry weight (g/plant)	Crop growth rate(g/m ² /day)
1.	Nitrogen 75kg + <i>Rhizobium</i>	37.14	21.00	6.13
2.	Nitrogen 75 kg + PSB	39.54	22.05	6.44
3.	Nitrogen 75 kg + <i>Rhizobium</i> + PSB	44.63	25.74	7.21
4.	Nitrogen 100 kg + <i>Rhizobium</i>	40.73	23.10	7.45
5.	Nitrogen 100 kg + PSB	41.43	23.50	7.45
6.	Nitrogen 100 kg + <i>Rhizobium</i> + PSB	46.19	26.20	8.22
7.	Nitrogen 125 kg + <i>Rhizobium</i>	42.38	23.77	7.34
8.	Nitrogen 125 kg + PSB	44.07	24.83	7.14
9.	Nitrogen 125 kg + <i>Rhizobium</i> + PSB	45.29	25.96	8.11
10.	Control (NPK 100-80-60 kg/ha)	40.20	22.79	7.59
	F-test	S	S	S
	SEm(±)	0.76	0.60	0.57
	CD (p=0.05)	2.24	1.77	1.68

Table: 2 Influence of Nitrogen levels and Biofertilizers on yield attributes and yield of French bean.

S.No.	Treatment combination	Number of pods /Plants	Number of seeds/pod	Seed Yield (q/ha)	Stover Yield (q/ha)
1.	Nitrogen 75kg + <i>Rhizobium</i>	14.00	3.87	16.00	43.55
2.	Nitrogen 75 kg + PSB	14.80	4.03	16.33	44.90
3.	Nitrogen 75 kg + <i>Rhizobium</i> + PSB	18.27	4.65	20.61	47.65
4.	Nitrogen 100 kg + <i>Rhizobium</i>	15.80	4.37	18.36	46.19
5.	Nitrogen 100 kg + PSB	16.00	4.44	18.76	46.00
6.	Nitrogen 100 kg + <i>Rhizobium</i> + PSB	19.70	5.08	22.47	50.08
7.	Nitrogen 125 kg + <i>Rhizobium</i>	16.67	4.51	19.79	46.03
8.	Nitrogen 125 kg + PSB	17.13	4.59	19.63	46.05
9.	Nitrogen 125 kg + <i>Rhizobium</i> + PSB	18.70	4.81	21.16	48.29
10.	Control (NPK 100-80-60 kg/ha)	15.33	4.37	17.53	45.86
	F-test	S	S	S	S
	SEm(±)	0.82	0.14	0.54	0.75
	CD (p=0.05)	2.45	0.40	1.62	2.40

REFERENCE

1. Adsule, R.N., Deshpande, S.S and Sathe, S.K. (1998). French bean In: Handbook of Vegetable Science and Technology, [Salunkhe, D.K. and Kadam, S.S. (eds.)] Marcel Dekker, Inc., New York pp. 457-469.
2. Alsina I, Dubova L, Karlovska A, Steinberga V, Strauta L. Evaluation of Effectiveness of *Rhizobium Leguminosarum* Strains on Broad Beans, *ActaHorticulturae*, No. 2016; 1142:**63**.
3. Behura, A.K.; Mahapatra, P.K. and Swain, D. (2008). Effect of irrigation and nitrogen on yield, water and nitrogen use efficiency in rajmash (*Phaseolus vulgaris* L.). *Legume Research*, **31** (1): 40 -43.
4. Dwivedi GK. Evaluation of Rhizobium leguminosorum (bv. Phaseoli) Strains in an Acid Iceptisol for N-fixing Ability in French bean under Temperate Region of Himalaya. *Legume Research*. 2007; **30**(4):279-282.
5. Dwivedi, Y.C., Sharma, R.S. and Sengupta, S.K. (1995). Effect of phosphorus and potassium fertilization on seed yield of French bean (*Phaseolus vulgaris* L.). *Vegetable Science***22** (1):36-38.
6. Garg OK, Hemantaranjan A, Ramesh C. Effect of iron and zinc fertilization on senescence in French bean (*Phaseolus vulgaris* L.), *Journal of Plant Nutrition*. 2008; **9**(3-7):257-266.
7. Gopalakrishnan, T.R. (2007). Vegetable Crops. New India Publishing Agency, New Delhi (India).
8. Karasu, A., Oz, M. and Dogan, R. (2011). The effect of bacterial inoculation and different nitrogen doses on yield and yield components of some dwarf dry bean cultivars (*Phaseolus vulgaris* L.). *Bulgarian Journal of Agricultural Science***17** (3): 296-305.
9. Kushwaha, B.L. (1994). Response of French bean to nitrogen application in North Indian plains. *Indian Journal of Agronomy***39** (1):34-37.
10. Manivannan, S., Batamurugan, M., Parthasarathi, K., Gunasekaran, G. and Ranganathan, L. S. (2009). Effect of vermicompost on soil fertility and crop productivity - beans (*Phaseolus vulgaris*). *Journal of Environmental Biology***30** (2): 275-281.
11. Mongi, Rose; Tongoona, Pangirayi; Shimelis, Hussein and Sibiya, Julia (2016). Appraisal of common bean farming systems under angular leaf spot disease prone environments of the southern highlands of Tanzania. *Indian Journal of Agricultural Research*, **50** (5): 428-433.
12. Moniruzzaman, M., Rahman, S.M.L., Kibria, M.G., Rahman, M.A. and Kaiser, M.O. (2007). Performances of vegetable French bean as influenced by varieties and sowing dates in rabi season. *International Journal of Sustainable Crop Production***2** (5):69-73.
13. Panse, V.G. and Sukhatame, P.V. (1984). Statistical Methods for Agricultural Workers. Fourth edition. ICAR Publication, New Delhi.
14. Parithiban, S. and Thamburaj S. (1991). Influence of Rhizobium culture and nitrogen fertilization on French bean. *South Indian Horticulture*, **39** (3):137-138.

15. Rajput, Pankaj Kumar, Singh, O.N., Singh, Yogeshwar and Singh, J.P. (2006). Integrated nutrient management for quantitative and qualitative yield of French bean (*Phaseolus vulgaris* L.). *Vegetable Science***33** (2):155-159.
16. Ramana, V., Ramakrishna, M., Purushotham, K. and Reddy, K. Balakrishna (2010). Effect of biofertilizers on growth, yield attributes and yield of French bean (*Phaseolus vulgaris*). *Legume Research*. **33** (3):178-183.
17. Ramana, V., Ramakrishna, M., Purushotham, K. and Reddy, K. Balakrishna (2011). Effect of biofertilizer on growth, yield and quality of French bean. *Vegetable Science*. **38** (1):35-38.
18. Sardana, Virender; Dhingra, K.K.; M.S. Gill and Singh, InderJit (2000). Production technology of French bean (*Phaseolus Vulgaris* L.) cultivation: A review. *Agric. Rev.*, **21** (3): 141-154.
19. Ssali H, Keya SO. The effect of phosphorus and nitrogen fertilizer level on nodulation, growth and dinitrogen fixation of three bean cultivars. *Trop. Agric*. 1986; **63**:105-109.
20. Singh, Anjani K., Singh, S.B. and Singh, Vineeta (2009). Influence of nitrogen doses on growth and green pod yield parameters of French bean varieties during kharif season under sub tropical area of Jammu region. *Legume Research***32** (2): 142-144.
21. Singer SM, Ali AH, El-Desuki MM, Gomaa AM, Khalafallah MA. Synergistic Effect of Bio-and Chemical Fertilizers to Improve Quality and Yield of Snap Bean Grown in Sandy Soil, *ActaHorticulturae*. 1998; (**513**):**24**.
22. . Tiwari S, Chouhan RK, Singh R, Sukla R, Gaur R. Integrated Effect of Rhizobium and Azatobactor Culture on the Leguminous Crop Black Gram (Vignamungo), *Adv Crop Sci Tech*. 2017; (**5**):**3**.
23. Wange, S.S., Karkeli, M.S., Patil, J.D and Meher, B.B. (1996). Effect of rhizobial inoculation and fertilizer nitrogen on French bean varieties. *Journal of Soils and Crops*. **6** (1):132-133.