

Rebuttals in red and blue fonts
Red = Correct
Blue = Delete
Build on the given document

Case study

A CASE OF MAXILLARY PERIPHERAL OSSIFYING FIBROMA AND ITS SURGICAL MANAGEMENT

Running Title - PERIPHERAL OSSIFYING FIBROMA OF MAXILLA

**Smart title: Surgical Management of Peripheral Maxillary Ossifying Fibroma in Level
..... facility of Udaipur, Rajasthan**

ABSTRACT

Peripheral ossifying fibroma is an enlargement of the gingiva that is fibromatous and non-neoplastic and exhibits regions of calcification/ossification. The present case shows a peripheral ossifying fibroma of the maxilla. The lesion was visible on attached gingiva of maxillary posterior teeth. **State a brief magnitude of the problem, study population, sample size, calculation procedure was done under** local anesthesia, fibroma was surgically removed and L-PRF membrane was placed over the underlying exposed bone. Histological examination was done to confirm the **diagnosis of peripheral ossifying fibroma of the maxilla.** L-PRF membrane as a biomaterial to cover the exposed areas of the bone after surgical excision is an effective option for better secondary intention healing and it provides acceptable results for tissue regeneration.

Key words– Fibroma; ossifying; calcification; granuloma; pyogenic; peripheral; surgical; biopsy,

INTRODUCTION

Introduce the topic briefly, On the gingiva, isolated reactive lesions ranging from various kinds, which include 'fibrous hyperplasia', 'peripheral ossifying fibroma', 'peripheral giant cell granuloma', and 'pyrogenic granuloma', may develop.¹ Most prevalent kind of gingival finding is a single gingival enlargement, and the cause of these lesions is usually a reactive

response to local irritation, however other potential causes include microorganisms, plaque, calculus, restorations, irritants, and dental appliances.

Mainly there are two categories of fibromas: 'central' type and 'peripheral' type. The peripheral type primarily impacts the soft tissues covering the tooth-containing areas of the jaws, whereas the central type causes an expansion of the medullary cavity and originates from the endosteum or periodontal ligament closer to the root apex.² POF is a gingival growth that is fibromatous, non-cancerous, and distinguished by regions of calcification or ossification. It is made up of a cellular fibroblastic connective tissue stroma and haphazardly distributed calcification foci that are composed of bone or cementum-like tissue.³ 'Shepherd' described it for the first time as "*maxillary exostosis*" in 1844. 'Eversol and Robin' initially coined the term "*Peripheral Ossifying Fibroma*" in 1972. Various other synonyms for POF include "*Calcifying or ossifying fibroid epulis*", "*peripheral fibroma with calcification*", "*calcifying fibroblastic granuloma*", "*peripheral cementifying fibroma*", or "*mineralizing ossifying pyogenic granuloma*".^{citation} POF manifests itself clinically as ossifying fibroma that seems sessile or pedunculated nodular mass that is generally red to pink in color, and most frequently found in children and young adults. It tends to affect the maxillary incisor and cuspid region.²

Clinical case presentation^{citation}:

The primary complaint of a 42 year old female patient who reported to the department of Periodontics at Pacific Dental College and Hospital (PDCH) in Udaipur, Rajasthan was that her upper left front and back teeth had been loosening for the past month, and a lump-like **swelling** in gums of the same area since previous seven months (Figure 1). There was no pertinent medical history, injuries, or ulceration history.^{citation}

The oral cavity was clinically examined intraorally, and on the keratinised gingiva, a single, pedunculated nodular growth that expanded mesiodistally from the distal aspect of the canine (23) to the mesial aspect of the first molar (26) and covered up to more than two-third of the teeth was seen (Figure 2). The lesion was well-defined, reddish, circular to oval-shaped, non-tender, solid in consistency, and measured roughly 3 cm by 2.5 cm by 2 cm (Figure 3). The lesion did not bleed when provoked and had pathological migration of 24 (Figure 3) and mobility of grade III in 22, 23, 24, and 25. In addition, generalised Miller's Class II gingival recession was clearly evident.^{citation}

Upon extraoral examination, it was found that the left side, from the philtrum to the angle of the mouth, was enlarged. Pyogenic granuloma and Peripheral ossifying fibroma were included in the differential diagnosis along with a provisional diagnosis of traumatic fibromacitation.

The IOPA revealed bone loss upto the middle of the root with diffuse ill-defined periapical radiolucency suggestive of periapical abscess(Figure 12). CBCT revealed bone loss uptotheapical third of the root along with an expansile lesion evident on the left side of the maxilla extending from the 24 – 26 tooth region with small radio-opaque foci of calcification noted within the lesion along with the expansion of the buccal cortex(Figure 13). Routine haematological investigations were normal.citation

CASE PRESENTATION:

Following thorough scaling and root planing, the patient was instructed to come back seven days later for an excisional biopsy under local anaesthesia. After appropriate antisepsis, local anaesthesia was administered, the lesion was excised (Figure 4), and the accompanying grade III mobile teeth were extracted (Figure 5)citation. Curettage of the area was done and as bony spicules were present, bone file was used to smoothen the bone. L-PRF membrane was placed over the area as there was a tissue deficiency and for better healing (Figure 6). A periodontal dressing (COE -PAK™ AUTOMIX) was applied (Figure 7), and the patient was summoned back for a check-up and the dressing was removed after 7 dayscitation.

On the 7th day, adequate healing was seen at the site and patient was kept on follow-up for 2 months and no recurrence was observed. (Figure 9,10).

Histopathological report:

Upon histopathological evaluation, it was discovered that the squamous epithelium was stratified and parakeratinized and the connective tissue consisted of collagen fibers, fibro-cellular, and plump fibroblasts between collagen bundles. The stroma displayed lamellar bone trabeculae, erratic basophilic, cementum-like materials, and a persistent inflammatory infiltrate of lymphocytes and plasma cells.citation(Figure 11,14)

DISCUSSIONdiscuss your results and findings in relation to other scholars works

With a 2% prevalence across all oral biopsies, the peripheral ossifying fibroma is by far the most common lesion. It is a clearly defined mass of tissue with a sessile or pedunculated

base that is situated on the gingiva. The surface may be intact (34%) or ulcerated (66%) and the colour may correlate with normal mucosa or lightly reddish⁵[send to first paragraph of introduction](#). POF typically measures less than 1.5 cm, but in some instances, sizes as large as 6 to 9 cm have also been documented.⁶ Although the etiopathogenesis of POF is still unresolved it is thought that periodontal ligament cells are the cause of the lesion in either case. Some researchers thought that POF was a '*neoplastic process*', while others thought it was a '*reactive process*'.⁸[discuss your results and findings in relation to other scholars works](#)It is thought to originate from the PDL because POF exclusively occurs in the gingival; especially interdental papilla, the gingival area closer to the PDL, and some lesions have oxytalan fibres within the matrix that is mineralized.⁸ Since many other conditions, such as pyogenic granuloma or peripheral giant cell granuloma, might display similar clinical manifestations and clinical courses, a conclusive diagnosis of POF cannot be established based solely on clinical findings. Therefore, a biopsy and histopathologic examination are necessary for the absolute diagnosis in order to be certain. Comprehensive overviews of the broad histologic spectrum of POF have been given by '*Buchner et al.*'¹⁰ All the fibromatous lesion will not show ossification or calcification.⁸ Although POF does not always result in radiographic changes, occasionally a central point of radiopaque material may be visible, especially in more severe or mineralised lesions.^{11, 12} Surgical excision of POF is the absolute treatment after elimination of local etiological factors like dental plaque, calculus, ill-fitting dentures, and subpar restorations. In order to prevent recurrence, the periosteum and periodontal ligament beneath the lesion should be eliminated. Recurrence rates have been observed to range from 8% to 20%.^{10, 12}[discuss your results and findings in relation to other scholars works](#)

When '*Kennedy et al.*' compared 'peripheral ossifying fibroma' and 'peripheral odontogenic fibroma', they discovered that factors like a high female predilection and a peak incidence in the second decade of life suggests hormonal influences.⁵ By analysing biopsy tissues histopathologically, definitive diagnosis of POF can be made. When performing a microscopic assessment, the following characteristics are frequently seen: (1) Presence of connective tissue that is fibrous and benign, (2)irregular to frequent endothelial growth, (3)Mineralized substance that appear as mature, lamellar, or woven osteoid, cementum-like, or dystrophic calcified material. Inflammatory cells, either acute or chronic, can also be discovered in lesions.⁸ In our case, most of the above features were present. In order to successfully treat POF, local surgical excision is advised which was done in this particular

case along with the extraction of the related grade III mobile teeth.[discuss your results and findings in relation to other scholars works](#)

In this case, after surgical excision and extraction of teeth, we placed an L-PRF membrane over the exposed bone surface as L-PRF is a simple to handle naturally affordable substance that is created by centrifuging patient blood without the use of anticoagulants. It promotes faster healing and releases growth factors like ‘platelet-derived growth factor’, ‘transforming growth factor beta 1’, and ‘vascular endothelial growth factor’ for seven days. In conjunction with important coagulation and matrix proteins like thrombospondin-1, fibronectin, and vitronectin, L-PRF membrane acts as an adjuvant for promoting and boosting up healing and/or tissue regeneration.[discuss your results and findings in relation to other scholars works](#)

The case of gingival overgrowth that was described here has been identified histopathologically as peripheral ossifying fibroma as no giant cells were found in the connective tissue stroma and the lesion had less vascularity, the possibility of peripheral giant cell granuloma and pyogenic granuloma was ruled out.[discuss your results and findings in relation to other scholars works](#)

The patient underwent a successful excision of POF, was maintained on periodic maintenance therapy for three months, and is still being monitored for signs of recurrence.[discuss your results and findings in relation to other scholars works](#)

CONCLUSION

POF is a chronic lesion that develops gradually. Because there generally no symptoms present, the growth frequently grows to a huge size before the patient chooses to seek treatment. POF can be misdiagnosed with other reactive lesion, so a histopathological study of the specimen is required for an absolute diagnosis. The possibility of recurrence is significant in presence of local factors and incomplete surgical removal. The lesion and the nearby tissues must be meticulously removed using curettage in order to prevent the recurrence.[Sent on material on clinical presentation, not smart conclusion.](#)

REFERENCES, use APA referencing system. Name of author in alphabetical order, year,

title page

1. Farquhar T, MacLellan J, Dymont H, Anderson RD. Peripheral ossifying fibroma: a case report. *J Can Dent Assoc.* 2008; 74: 809-12.
2. Keluskar V, Byakodi R, Shah N. Peripheral ossifying fibroma. *J Indian Acad Oral Med Radiol.* 2008; 20: 54.
3. Das UM, Azher U. Peripheral ossifying fibroma. *Indian Soc Pedod Prev Dent.* 2009; 27: 49.
4. Chatterjee A, Ajmera, Singh A. Peripheral cemento-ossifying fibroma of maxilla. *J Indian Soc Periodontol.* 2010; 14: 186.
5. Kenney JN, Kaugars GE, Abbey LM. Comparison between the peripheral ossifying fibroma and peripheral odontogenic fibroma. *J Maxillofac Surg.* 1989; 47: 378-82.
6. Poon C-K, Kwan P-C, Chao S-Y. Giant peripheral ossifying fibroma of the maxilla: report of a case. *J Maxillofac Surg.* 1995; 53: 695-8.
7. Kumar SK, Ram S, Jorgensen MG, Shuler CF, Sedghizadeh PP. Multicentric peripheral ossifying fibroma. *J Oral Sci.* 2006; 48: 239-43.
8. Buchner A, Hansen LS. The histomorphologic spectrum of peripheral ossifying fibroma. *Oral Surg.* 1987; 63: 452-61.
9. Flaitz C. Peripheral ossifying fibroma of the maxillary gingiva. *Am J Dent.* 2001; 14: 56.
10. Cuisia Z, Brannon R. Peripheral ossifying fibroma--a clinical evaluation of 134 pediatric cases. *Pediatr Dent.* 2001; 23: 245-8.




Fig 1: Frontal View




Fig 2: Apico-coronal extension



Fig 3: Mesio-distal extension




Fig 4: Surgical Excision and preservation of associated teeth




Fig 6: L-PRF membrane placement dressing placement




Fig 8: Excised mass

Fig 9: 1 month re -evaluation

Fig 10: 2 months re-evaluation



Fig 11: Histopathological examination

Fig 12: CBCT

UNDER PEER REVIEW

UNDER PEER REVIEW