

Original Research Article

Different Agricultural Practices Adopted by Farmers in **Kharar** Division, Punjab

Title could be changes as follows:

A study on adoption level of different recommended agricultural practices by farmers in Kharar division of Punjab

Abstract:

Since agriculture employs half of all workers worldwide, it is the largest employer and the economic engine of rural India. Rural India's development in relation to national development would be reflected in the development of agriculture [11]. Agriculture and allied industries are without a doubt the greatest employer in India, particularly in the vast rural areas. A vital basis for the Gross Domestic Product is also provided by it (GDP). The agriculture industry in India contributes considerably to the economy, accounting for around **16%** of GDP. All of the procedures that a farmer follows from seed to seed, from seed sowing through seed harvesting and storage, are referred to as agronomical practices. A study on agronomic practices followed by 120 farmers from five villages viz. Shakrullapur (28 farmers), Rora (30 farmers), Bibipur (22 farmers), Batta (18 farmers) and Fatehpur-Theri (22 farmers) was considered. The data collected from the respondents include major crops grown, seed rate followed by respondents, fertilizer dose followed, major weeds affecting, major plant diseases, major insect and pests, harvesting methods and yield records. As per the data collected, wheat is cultivated by nearly 87% of the farmers in this region and 100% farmers who are growing wheat, agreed that the *Phalaris minor* is the major weed affecting the wheat production. Also Sheath blight and Yellow rust **were** the two major diseases affecting the paddy and wheat respectively. The study's conclusions **illustrated that the farmers were adopted latest techniques** in the fields to ensure the best crop stand and highest yields.

Key words: *Agronomy, insects, pests, fertilizer, diseases, yield, weeds*

1. Introduction:

The agricultural sector contributes strategically to a nation's economic growth. It has already had a substantial impact on the economic prosperity of developed **nations and its** contribution to the economic growth of less developed nations is crucial. In other words, the focus is on agriculture and other primary industries where per capita real income is low. One of the most established financial sectors in our country is agriculture. Different districts use different development strategies. Undoubtedly the largest source of employment in India is agriculture and its related

sectors, especially in the vast rural areas. Additionally, it provides a crucial foundation for the Gross Domestic Product (GDP). In India, the agricultural sector contributes significantly to the economy, making up about 16% of GDP (GDP). India's population depends on it for survival to the tune of about 62% [4]. Given that it accounts for roughly 16% of GDP, agriculture is a vital sector of the Indian economy [2].

The most significant factor and largest contributor to the Indian economy is agriculture (This statement is invalid as agril sector contributes only 16% to GDP and hence cannot say largest contributor to indian economy). As the "Agricultural Hub of India," states like Punjab, Haryana, and Uttar Pradesh are referred to (this statement is incomplete and to be redrafted). In rural India, the main source of employment is agriculture, which accounts for half of all employment worldwide [12]. Nearly 58% of Indians' income comes from agriculture, and their population is still expanding quickly [1]. Agriculture contributes \$400 billion to India's GDP, which is second only to China [12]. Wheat (*Triticumaestivum* L.), the most important cereal crop in the world, is a staple food for around one-third of the world's population [5]. It accounts for about 21% of all cultivated land (30.597 million hectares) and about 35% of the nation's total food production (98.38 million tonnes) (2016–17) [8]. For India's food security and economic growth, rice (*Oryza sativa*), a staple crop, is essential [3]. It provides over one-fourth of all caloric intake and occupies more than one-fifth of all gross cropland [7].

A vital component of the sustainability of food systems is agricultural technology. An illustration of how scale-independent technology has changed agricultural productivity is the Green Revolution. The Green Revolution has improved harvests, decreased poverty, built infrastructure, increased access to food, and decreased food prices, among other things [6]. A farmers activities from seed to seed are referred to as farming practices from planting to gathering and storing. The relationship between agronomic practices and yields is direct. Farmers must use the finest agricultural techniques for the season and agro-climatic zone in order to get optimum harvests. By region, these traditions change. It has been crucial in assuring improved agricultural practices in nations like India, allaying worries that the country has reached the point of food excess, which happens when population growth outpaces agricultural production. India is second in the world for agricultural production despite having a little portion of the world's agricultural land.

2. Material and Methods:

The study was conducted in Kharar block, district SAS Nagar (Punjab) in which five villages were selected randomly Shakrullapur, Rora, Bibipur, Batta and Fatehpur-Theri. A total of 28 respondents were selected from village Shakrullapur, 30 were from Rora, 22 were from Bibipur, 18 were from Batta and 22 respondents are from Fatehpur-Ther making total sample of 120 for the study. A detailed interaction was directed with the farmers regarding their social status, and the Agronomical practices they follow throughout the year. The respondents were interviewed at

their homes and in their fields. To cover every aspect regarding the agronomical practices followed by the farmers an interview schedule was prepared to analyze each parameter properly in a sequence.

After data collection from respondents, data were classified and analyzed with the help of suitable statistical measures such as percentages, graphical representation, bar graphs, and Pie charts.

Formulae used for calculation:

$$\text{Percentage} = \frac{\text{Number of respondents}}{\text{Total number of respondents}} * 100$$

$$\text{Average} = \frac{\text{Sum of total observations}}{\text{Total number of observations}}$$

3. Results and Discussion

3.1 Major Crops Grown by the Farmers:

The data represented in the Table 1 shows the major crops grown by the farmers in their respective villages. As per the data collected overall, 87% farmers are growing wheat, 85% farmers are growing paddy and 52% farmers are growing mustard. Apart from these crops, 44% farmers are growing sugarcane, 41% farmers are growing berseem as a fodder crop, 35% farmers are growing maize, 34% farmers are growing bajra and 31% farmers are growing sorghum. Vegetable crops are grown less in this region. Less than 10% farmers are growing potato and onion.

Sr. No.	Crops grown	Shakrullapur (n=28)	Rora (n=30)	Bibipur (n=22)	Batta (n=18)	Fatehpur-Theri (n=22)	Overall (N=120)
1	Wheat	22 (79%)	28 (93%)	20 (91%)	15 (83%)	20 (91%)	105 (87%)
2	Paddy	18 (64%)	28 (93%)	22 (100%)	12 (67%)	22 (100%)	102 (85%)
3	Mustard	12 (43%)	13 (43%)	10 (45%)	11 (61%)	16 (73%)	62 (52%)
4	Sugarcane	13 (46%)	10 (33%)	14 (64%)	6 (33%)	10 (45%)	53 (44%)
5	Berseem	10 (36%)	11 (37%)	7 (32%)	11 (61%)	10 (45%)	49 (41%)
6	Maize	15 (54%)	8 (27%)	4 (18%)	5 (28%)	10 (45%)	42 (35%)
7	Bajra	12 (43%)	8 (27%)	7 (32%)	6 (33%)	8 (36%)	41 (34%)
8	Sorghum	11 (39%)	7 (23%)	6 (27%)	5 (28%)	8 (36%)	37 (31%)

9	Potato	3 (11%)	2 (7%)	1 (4%)	2 (11%)	3 (14%)	11 (9%)
10	Onion	2 (7%)	1 (3%)	3 (14%)	1 (6%)	3 (14%)	10 (8%)

Table 1: Major crops grown by respondent farmers




Fig 1: Crops grown by total percentage of farmers

3.2 Seed Rate Followed by the Farmers:

The data represented in Table 2 shows the total percentage of farmers using seeds less than the recommended rate, recommended rate and above the recommended rate. As per the data, most of the farmers are using recommended seed rate. In Shakrullapur village 7% farmers are using seeds below the recommended rate, 61% farmers using recommended seed rate and 32% farmers are using above recommended rate. In Rora village 10% farmers are using seeds below recommended rate, 53% farmers are using recommended rate and 37% farmers are using above recommended rate. In Bibipur village 55% farmers are using recommended seed rate and 45% farmers are using above than recommended seed rate. In Batta village 11% farmers are using below recommended seed rate, 61% farmers are using recommended seed rate and 28% farmers are using above than the recommended seed rate. In the Fatehpur-theri village 68% farmers are using recommended seed rate and 32% farmers are using more than recommended seed rate.

Sr.	Seed Rate	Shakrulla	Rora(n=3)	Bibipur(n)	Batta(n=1)	Fatehpur-	Overall
-----	-----------	-----------	-----------	------------	------------	-----------	---------

No		pur (n=28)	0)	=22)	8)	Theri (n=22)	(N=120)
1	Below recommended rate	2 (7%)	3 (10%)	0	2 (11%)	0	7 (6%)
2	Recommended rate	17 (61%)	16 (53%)	12 (55%)	11 (61%)	15 (68%)	71 (59%)
3	Above recommended rate	9 (32%)	11 (37%)	10 (45%)	5 (28%)	7 (32%)	42 (35%)

Table 2: Recommended seed rate followed by the farmers in percentage




Fig 2: Recommended seed rate followed by farmers in percentage

3.3 Fertilizer Dose Applied by the Farmers:

Since major crops grown by the farmers of these villages are wheat, paddy, mustard and sugarcane. The results shown in Table 3 depicts that the overall mean amount of fertilizers used by the respondents are more than the recommended doses. The recommended dose for the wheat crop is 120N: 60P: 40K [10], recommended dose for the paddy crop is 125N: 30P: 30K [9], the recommended dose for mustard crop is 130N: 120P [10] and the recommended dose for sugarcane is 300N:100P:200K [9]. The results revealed that farmers were

adopted excess/indiscriminate fertilizers doses in all crops. Potash fertilizers are accorded less importance when compared to nitrogenous and phosphoric fertilizer respectively. However, most of these fertilizer application do not take into account the needs of the soil for N, P, and K.

Sr. no	Parameters	Shakrulapur	Rora	Bibipur	Batta	Fatehpur-Theri	Average	Recommended dose/ha
Wheat								
1	Urea/ha	300 kg/ha	340 kg/ha	282 kg/ha	326 kg/ha	293 kg/ha	308.2 kg/ha	120 kg/ha
2	DAP/ha	182 Kg/ha	206 kg/ha	160 kg/ha	175 kg/ha	200 kg/ha	184.6 kg/ha	60 kg/ha
3	Potassium/ha	-	-	-	-	-	-	40 kg/ha
Paddy								
1	Urea/ha	150 kg/ha	195 kg/ha	225 kg/ha	186 kg/ha	210 kg/ha	193.2 kg/ha	125 kg/ha
2	DAP/ha	373 kg/ha	356 kg/ha	340 kg/ha	330 kg/ha	347 kg/ha	349.2 kg/ha	30 kg/ha
3	Potassium/ha	-	-	-	-	-	-	30 kg/ha
Mustard								
1	Urea/ha	195 kg/ha	225 kg/ha	202 kg/ha	190 kg/ha	212 kg/ha	204.8 kg/ha	130 kg/ha
2	DAP/ha	254 kg/ha	297 kg/ha	285 kg/ha	260 kg/ha	270 kg/ha	273.2 kg/ha	120 kg/ha
Sugarcane								
1	Urea/ha	540 kg/ha	460 kg/ha	402 kg/ha	512 kg/ha	520 kg/ha	486.8 kg/ha	300 kg/ha
2	DAP/ha	267 kg/ha	288 kg/ha	285 kg/ha	300 kg/ha	276 kg/ha	283.2 kg/ha	100 kg/ha

3	Potassium/ha	291 kg/ha	321 kg/ha	326 kg/ha	298 kg/ha	310 kg/ha	309.2 kg/ha	200 kg/ha
---	--------------	-----------	-----------	-----------	-----------	-----------	-------------	-----------

Table 3: Represents the mean value of N:P:K amount applied by the respondents

3.4 Major Weeds:

As seen in the Table 4 different kinds of weeds (Crop specific) are present in their fields throughout the year. The main weed that all of the responders saw was *Phalaris minor* (87%). According to them, this weed shows resistance to most of the chemicals used to eradicate or eliminate this weed. It reduces the yield up to a significant level. This was the major concern of the farmers during the Rabi season, especially in wheat crop.

Sr. No	Name of weeds	Crop affected	Shakrulla pur (n=28)	Rora (n=30)	Bibipur (n=22)	Batta (n=18)	Fatehpur-Theri (n=22)	Overall I (N=120)
1	<i>Phalaris minor</i>	Wheat	22 (79%)	28 (93%)	20 (91%)	15 (83%)	20 (91%)	105 (87%)
2	<i>Avenafatua</i>	Wheat	22 (79%)	28 (93%)	19 (86%)	15 (83%)	19 (86%)	103 (86%)
3	<i>Avenaludoviciana</i>	Wheat	20 (71%)	27 (90%)	18 (82%)	14 (78%)	19 (86%)	98 (82%)
4	<i>Echinochloa crus-galli</i>	Paddy	16 (57%)	26 (87%)	19 (86%)	10 (56%)	21 (96%)	92 (77%)
5	<i>Partheniumhyster ophorus</i>	Sugar cane	11 (39%)	9 (30%)	11 (50%)	5 (28%)	9 (41%)	45 (38%)

Table 4: various crops affected by major weeds




Fig 3: Percentage of farmers affected by different weeds

3.5 Plant Diseases:

According to information acquired from five villages indicated in Table 5, around 78% of farmers producing rice reported occurrence of sheath blight, 63% reported occurrence of brown leaf spot, and 74% reported occurrence of dwarf plant diseases in their fields. 71% of farmers producing wheat reported occurrence of Powdery mildew, 83% reported yellow smut, and 72% reported loose smut disease in their wheat crop. In maize fields 22% of farmers reported occurrence of Seed rot and 27% reported occurrence of Sheath blight in their maize crop. Similarly in sugarcane crop, 39% of farmers witnessed red rot disease and 32% of farmers witnessed smut disease in their fields. In mustard crop 46% of Farmers reported *Alternaria blight* in their field and 39% farmers witnessed *Sclerotinia stem rot* in their fields.

Sr No	Crop	Parameters	Shakrullapur (n=28)	Rora (n=30)	Bibipur (n=22)	Batta (n=18)	Fatehpur -Theri (n=22)	Overall (N=120)
1	Rice	Sheath Blight	17 (61%)	25 (83%)	19 (86%)	12 (67%)	20 (91%)	93 (78%)
		Brown leaf spot	12 (43%)	22 (73%)	15 (68%)	8 (44%)	18 (82%)	75 (63%)

		Dwarf plant disease	15 (54%)	24 (80%)	21 (95%)	10 (56%)	19 (86%)	89 (74%)
2	Wheat	Powdery Mildew	20 (71%)	21 (70%)	16 (72%)	12 (67%)	16 (73%)	85 (71%)
		Yellow rust	22 (79%)	27 (90%)	18 (82%)	13 (72%)	20 (91%)	100 (83%)
		Loose smut	21 (75%)	24 (80%)	13 (59%)	10 (56%)	18 (82%)	86 (72%)
3	Maize	Seed rot	9 (32%)	5 (17%)	3 (14%)	2 (11%)	7 (32%)	26 (22%)
		Sheath blight	12 (43%)	6 (20%)	3 (14%)	3 (17%)	8 (36%)	32 (27%)
4	Sugarcane	Red rot	11 (39%)	8 (27%)	12 (55%)	6 (33%)	10 (46%)	47 (39%)
		Smut	9 (32%)	10 (33%)	7 (32%)	4 (22%)	8 (37%)	38 (32%)
5	Mustard	Alternaria blight	11 (39%)	13 (43%)	10 (45%)	6 (33%)	15 (68%)	55 (46%)
		Sclerotinia Stem rot	12 (43%)	9 (30%)	7 (32%)	9 (50%)	10 (45%)	47 (39%)

Table 5: Represents the data of farmers affected by various crops diseases




Fig 4: Percentage of farmers affected by diseases in rice




Fig 5 : Percentage of farmers affected by diseases in wheat




Fig 6: Percentage of farmers affected by diseases of maize




Fig 7: Percentage of farmers affected by diseases of sugarcane




Fig 8: Percentage of farmers affected by diseases of mustard

3.6 Insect and Pest:

According to data analysis gathered from 5 villages shown in Table 6 indicated that in paddy crop roughly 83% of farmers witnessed grasshoppers, 69% of farmers witnessed leaf folders, and 70% of farmers witnessed Rice Hispa in their fields. In fields of wheat crop 85% of farmers witnessed aphids while 68% farmers witnessed termites. In fields of maize crop 24% of farmers witnessed

mites, 26% of farmers witnessed stem borer and 17% farmers witnessed fall armyworm in their field. Similarly in sugarcane crop, 38% of farmers witnessed termites in their field, while 41% of farmers observed top borer. In Mustard, 43% of farmers witnessed hairy caterpillars and 45% of farmers reported aphids attack in their fields.

Sr No	Crop	Parameters	Shakrullapur (n=28)	Rora (n=30)	Bibipur (n=22)	Batta (n=18)	FatehpurTheri (n=22)	Overall (N=120)
1	Rice	Grasshopper	18 (64%)	28 (93%)	21 (95%)	12 (67%)	20 (91%)	99 (83%)
		Leaf folder	16 (57%)	24 (80%)	17 (77%)	10 (56%)	16 (3%)	83 (69%)
		Rice Hispa	17 (61%)	19 (63%)	15 (68%)	12 (67%)	21 (95%)	84 (70%)
2	Wheat	Aphids	22 (79%)	28 (93%)	17 (77%)	15 (83%)	20 (91%)	102 (85%)
		Termites	19 (68%)	17 (56%)	14 (63%)	13 (72%)	19 (86%)	82 (68%)
3	Maize	Mites	9 (32%)	6 (20%)	3 (14%)	2 (11%)	9 (41%)	29 (24%)
		Stem borer	12 (43%)	5 (17%)	3 (14%)	3 (17%)	8 (36%)	31 (26%)
		Army worm	7 (25%)	6 (20%)	1 (4%)	2 (11%)	4 (18%)	20 (17%)
4	Sugarcane	Top Borer	12 (48%)	9 (30%)	14 (64%)	6 (33%)	8 (36%)	49 (41%)
		Termites	13 (46%)	7 (23%)	11 (50%)	4 (22%)	10 (45%)	45 (38%)
5	Mustard	Aphids	10 (37%)	13 (43%)	10 (45%)	5 (28%)	16 (73%)	54 (45%)
		Hairy Caterpillar	12 (43%)	9 (30%)	8 (36%)	9 (50%)	14 (64%)	52 (43%)

Table 6: Represents the data of farmers affected by different pests infestation




Fig 9: Percentage of farmers affected by different pests in rice




Fig 10: Percentage of farmers affected by different pests of Wheat




Fig 11: Percentage of farmers affected by different pests of maize




Fig 12: Percentage of farmers affected by different pests of sugarcane




Fig 13: Percentage of farmers affected by different pests of mustard

3.7 Harvesting Method:

The data shown in Table 7 revealed that majority of farmers favor use of combine harvester for harvesting crops like wheat and rice as it requires less time and effort. Among rice producing farmers combine harvester is used by about 72% of farmers, where as 13% preferred hand harvesting. In the wheat crop 74% of farmers used combine harvesters while 13% were preferred manual harvesting. In crops like maize, sugarcane and mustard farmers were preferred by manual harvesting. According to results 50% of farmers' were manually harvested mustard, 44% of farmers' manually harvested sugarcane and 28% of farmers' manually harvested maize crop.

Sr No	Crops	Parameters	Shakrullapur (n=28)	Rora (n=30)	Bibipur (n=22)	Batta (n=18)	FatehpurTheri (n=22)	Overall (N=120)
1	Rice	Combine harvester	12 (43%)	25 (83%)	20 (91%)	8 (44%)	21 (95%)	86 (72%)
		Manual	6 (21%)	3 (10%)	2 (9%)	4 (22%)	1 (5%)	16 (13%)
2	Wheat	Combine harvester	16 (57%)	25 (83%)	18 (82%)	11 (61%)	19 (86%)	89 (74%)
		Manual	6 (21%)	3	2 (9%)	4	1 (5%)	16

				(10%)		(22%)		(13%)
3	Maize	Manual	12 (43%)	6 (20%)	3 (14%)	3 (17%)	9 (41%)	33 (28%)
4	Sugarcane	Manual	13 (46%)	10 (33%)	14 (64%)	6 (33%)	10 (45%)	53 (44%)
5	Mustard	Manual	12 (43%)	13 (43%)	10 (45%)	9 (50%)	16 (73%)	60 (50%)

Table 7: Represents the data of farmers following different harvesting methods




Fig 14: Percentage of farmers following different harvesting methods in rice




Fig 15: Percentage of farmers following different harvesting methods in wheat




Fig 16: Percentage of farmers following manual harvesting method in maize, sugarcane and mustard

3.8 Yield Record:

As per the results shown in Table 8, total average production of paddy is 61.4 quintals/ha with highest yield of 63 quintals/ha in Bibipur and lowest yield of 60 quintals/ha in Rora and Batta. The average yield of wheat is 46.6 quintals/ha with Rora village having the highest yield of 48 quintals/ha and Batta village with lowest yield of 45 quintals/ha. Average yield of Maize is 37.8 quintals/ha with lowest yield of 37 quintals/ha in Shakrullapur and FatehpurTheri and highest yield of 39 quintals/ha in Rora village. Average production of Sugarcane is 810 quintals/ha with lowest yield of 790 quintals/ha in Shakrullapur and highest yield of 830 quintals/ha in FatehpurTheri village. Similarly Average yield of Mustard is 13.6 quintals/ha with lowest yield of 12 quintals/ha in Bibipur and highest yield of 14 quintals/ha in Shakrullapur.

Sr No	Crop	Shakrullapur	Rora	Bibipur	Batta	Fatehpur- Theri	Average production
1	Rice	62 quintals/ha	60 quintals/ha	63 quintals/ha	60 quintals/ha	62 quintals/ha	61.4 quintals/ha
2	Wheat	47 quintals/ha	48 quintals/ha	47 quintals/ha	45 quintals/ha	46 quintals/ha	46.6 quintals/ha
3	Maize	37 quintals/ha	39 quintals/ha	38 quintals/ha	38 quintals/ha	37 quintals/ha	37.8 quintals/ha

			a	a	ha	ha	
4	Sugarcane	790 quintals/ha	820 quintals/ha	810 quintals/ha	800 quintals/ha	830 quintals/ha	810 quintals/ha
5	Mustard	14 quintals/ha	15 quintals/ha	12 quintals/ha	13 quintals/ha	14 quintals/ha	13.6 quintals/ha

Table 8: Represents the average yield of major crops grown by the respondents

Conclusion:

Based on findings it could be concluded that a wide variety of crops are grown in this region like Wheat, paddy, sugarcane, maize, mustard, sorghum, berseem, potato, onion etc. Wheat and paddy are the two main cereal crops and mustard is the major oilseed crop cultivated in this area whereas major fodder crops grown were berseem and sorghum. Majority of the farmers using recommended seed rate followed by more than recommended rate. Most of the farmers (87%) facing the weed problem of *Phalaris minor*, commonly known as 'Gullidanda' in Punjabi. This was primarily due to the reason that farmer's non adoption of crop rotation in their fields. They were following both manual methods and weedicides to control the weeds in the fields. Majority of the farmers observed Sheath blight (78% farmers) in rice and yellow rust in wheat (83% farmers). To deal with the diseases farmers preferred different chemical control measures. Grasshopper and aphids were the major pests causing maximum damage in paddy and wheat crops respectively. Harvesting was mainly done by combine harvester in the case of cereal crops like wheat and rice whereas manual harvesting was adopted in maize, mustard, sugarcane, bajra, berseem etc. Most of the farmers were using excess amount of fertilizers. Farmers were using more complex fertilizers containing nitrogen and phosphorus fertilizers and paid less attention to potash fertilizers. To overcome this problem farmers should undergo routine soil testing to stay informed about the macro- and micronutrient state of their soil and should supply the crop with the ideal amount of nutrients based on the soil nutrient status. By using these agronomical techniques, productivity and economic standing could be improved.

References:

- [1] Agarwal, H. P., & Sinha, R. (2017). Urban farming-A sustainable model for Indian cities. *International Journal on Emerging Technologies*, 8(1), 236–242.
- [2] Department of Agriculture, Cooperation & Farmers' Welfare (DAC&FW). Annual Report (2020-21). <https://agricoop.nic.in/en/whatsnew>

- [3] Devkota, K. P., Sudhir-Yadav, Khanda, C. M., Beebout, S. J., Mohapatra, B. K., Singleton, G. R., & Puskur, R. (2020). Assessing alternative crop establishment methods with a sustainability lens in rice production systems of Eastern India. *Journal of Cleaner Production*, 244. <https://doi.org/10.1016/j.jclepro.2019.118835>
- [4] Gupta. & Nagar, M. (2017). Agriculture Sector in India: As a Career. *International Journal on Arts, Management and Humanities* 6(2): 01-06
- [5] Husnain, M., Bukhsh, M. A. H. A., Iqbal, J., Khaliq, T., & Zamir, S. I. (2011). Agro-economic response of two wheat varieties under different tillage practices. *CROP & ENVIRONMENT*, 2011(2), 1–7. www.psa.net.pk
- [6] McCullough, E. B., Pingala, P. L., & Stimuli's, K. G. (2012). Small farms and the transformation of food systems: an overview. *The Transformation of Agri-Food Systems*: 27-70
- [7] Mohanty, S., & Yamano, T. (2017). Rice Food Security in India: Emerging Challenges and Opportunities. In *The Future Rice Strategy for India* (pp. 1–13). Elsevier. <https://doi.org/10.1016/B978-0-12-805374-4.00001-4>
- [8] Nath, B., Bhattacharya, D., Debasis Bhattacharya, C., & Dhakre, D. (2019). Forecasting wheat production in India: An ARIMA modelling approach. ~ 2158 ~ *Journal of Pharmacognosy and Phytochemistry*, 8(1). <http://indiastat.com>
- [9] Package of practices Kharif season. PAU; 2022-23. Retrieved from: https://www.pau.edu/content/ccil/pf/pp_kharif.pdf
- [10] Package of practices Rabi season. PAU; 2022-23. Retrieved from: https://www.pau.edu/content/ccil/pf/pp_rabi.pdf
- [11] Singh AK, Singh AK, Kumar R, Prakash V Sundaram PK and Yadav SK. 2017. Indian Cereals Saga: Standpoint and Way Forward. *Journal of AgriSearch* 4 (1): 1-10.
- [12] Singh, A. K., Upadhyaya, A., Kumari, S., Sundaram, P. K., & Jeet, P. (2020). Role of Agriculture in making India \$5 trillion Economy under Corona Pandemic Circumstance: Role of agriculture in Indian economy. *Journal of AgriSearch*, 7(2), 54–58.