

## **Evaluation of Different Varieties and Nitrogen Levels on Growth and Yield of Black rice (*Oryza sativa* L.)**

### **ABSTRACT**

The field experiment was conducted at Crop Research Farm during *kharif* season, in the Department of Agronomy, Naini Agriculture Institute, Sam Higginbottom University of Agriculture, Technology And Sciences, Prayagraj Uttar Pradesh. The experiment comprised three varieties *viz.*, Chakhao Poireiton (V<sub>1</sub>), Chakhao Amubi (V<sub>2</sub>) and Chakhao Angangba (V<sub>3</sub>); and three rates of nitrogen *viz.*, 60 kg/ha (N<sub>1</sub>), 70 kg/ha (N<sub>2</sub>) and 80 kg/ha (N<sub>3</sub>). The experiment was laid out in a randomized block design with three replications. The growth analysis result was observed with the application of [Chakhao Poireiton and nitrogen (80 kg/ha)] produces higher plant height (142.53 cm), maximum number of tillers/hill (13.97), higher dry weight (32.49 g), higher panicle length (30.1 cm), maximum number of grains/panicle (153.60), higher test weight (24.01 g), higher grain yield (5.67 t/ha), higher straw yield (11.27 t/ha). However, the maximum gross return (2,38,070.00 INR/ha), maximum net return (1,91,800.00 INR/ha) and maximum benefit cost ratio (3.23) was also obtained with the same treatment 3 [Chakhao Poireiton + nitrogen (80 kg/ha)].

**Keywords:** *Black rice, N level, Growth, Yield and Economics.*

### **INTRODUCTION**

Rice (*Oryza sativa* L.) is the most important cereal crop in the developing world and is the most staple food of over half the world's population. It is generally considered a semi-aquatic annual grass plant. About 20 species of the genus *Oryza* are recognized, but nearly all cultivated rice is *O. sativa* L, family of Gramineae. 35-60% of the total calorie approximately one-fifth of all calories consumed by humankind. (Tayefe *et al.*, 2014). Presently more than 90% of total rice production and consumption is in Asia. In Asia, above two billion people receive 60-70 % of their energy necessities from rice (Sridhar *et al.*, 2019).

Black rice is a type of the rice species *Oryza sativa* L. which is glutinous, packed with high level of nutrients and mainly cultivated in Asia. Black rice is also known as purple rice, forbidden rice, heaven rice, imperial rice, king's rice and prized rice. This rice includes several varieties with a long history of cultivation in Southeast Asian countries such as China, India and Thailand. There are more than 200 types of black rice varieties in the world. Only China is responsible for 62 % of global production of black rice and it has developed more than 54 modern black rice varieties with high yield characteristics and multiple resistances. China cultivates most of the black rice followed by Sri Lanka, Indonesia, India and Philippines etc. Thailand occupies the ninth position for black rice cultivation. Black rice is a medium size grain, non-glutinous heirloom rice. Black rice has a dark purplish -Black colour with a nutty, slightly sweet flavour. Black rice is black due to the presence of the Black colour pigment called anthocyanin which contain high amounts of certain antioxidants. Black rice also has higher levels of protein, vitamins and minerals. Compared to white rice, black rice is relatively rich in mineral content such as Fe, Zn, Mn and P and has a high variability in mineral content that depends on varieties and soil types of the planting area. Supplementation of black scented rice in the diet will have a great impact on human health. A health benefit of black rice includes prevention of cancer, diabetes, heart diseases, Alzheimer's disease, gallstones etc.

In world rice growing in 165.12 million ha, the production is 509.42 million tonnes and the yield is about 4.61 metric tons/ha (**USDA, 2022**). Among the rice growing countries of the world, India has the largest rice acreage and ranked second in production after wheat in the world. In India rice is grown in 45.07 million ha, the production level is 122.27 million tones and the yield is about 2713 kg/ha (**GOI, 2021**). In Uttar Pradesh state ranks third in the country in production of rice. it grown over area about 5.68 m/ha which comprise of (13.5%) of total rice in India. Annual rice production is around 15.66 million metric tonnes, the average yield is 2759 kg/ha (**GOI, 2021**).

Nitrogen deficiency is one of the major constraints to crop production. The unique feature of N is its soils having very low soil organic matter content, soils with particular constraints on indigenous N supply, high potential of ammonia (NH<sub>3</sub>). Due to a deficiency of N, older leaves turn an orange-yellow color and die from the tip down; young leaves are thin, short, and rigid. Similarly, a deficiency of N causes a decrease in plant height and the number of tillers. The roots decrease in number and become slender and lengthy. It is one of Asia's most prevalent rice concerns and frequent in all rice-growing soils where modern varieties are planted in the absence

of adequate mineral N fertilizer. It frequently occurs during important growth stages of the plant, such as tillering and panicle initiation (**Shrestha et al., 2022**).

Chakhao Poireiton is cultivated by most of the farmers (43 %) because of its higher productivity and delicacy. As per local belief, black rice was first cultivated by the 12<sup>th</sup> Meitei king Poireiton Khunthokpa during 38-18 BC in his capital Poi located at the foothills of Heirok range. Thus the rice landrace came to be known as Chakhao poireiton. Black rice plants attain 136-166 cm height which was comparatively higher than the other traditional non-black rice varieties cultivated in the study area. Chakhao amubi was the tallest black rice. The highest panicle length number of grains, and tillers were recorded for Chakhao poireiton. The kernel length of Chakhao amubi and Chakhao poireiton are considered as long and medium (**Borah et al., 2018**). Chakhao poireiton and Chakhao amubi, were shown to have high anthocyanin and phenolics content and strong antioxidant activity. Chakhao poireiton was classified as waxy, as their amylose content was less than (2%). Chakhao amubi and chakhao Angangba was with very low amylose content (3.16%). The black chakhao cultivars are quite tall in height. Majority of chakhao cultivars ranges in height between 130-165cm and flowering is initiated between 95-116 DAT. The rice culture Chakhao Poireiton is a good alternative variety, it exhibits high yield potential (5-6 t/ha) and milling 60-70%.

Nitrogen is an essential nutrient element for plants and is necessary for photosynthesis, growth and development, yield, quality, and biomass production in rice. Nitrogen promotes rapid plant growth and increases grain yield and quality by increasing tillering, leaf area development, grain formation, grain filling, and protein synthesis. Nitrogen is highly mobile within the plants and soil. During the early and mid-tillering, panicle initiation, booting, and grain development phases of ripening, N is the most typically required nutrient element (**Bai 2019**). N has an important role in the creation of rice quality. Increasing N fertilization can help improve the nutritional quality and processing quality of rice, but excessive N fertilization can increase rice chalkiness and worsen the rice appearance quality, cooking and eating quality (**Li et al., 2019**). In order to optimize rice yield, different agricultural practices are applied and one of them is fertilizer application. Among the major nutrient elements, nitrogen (N) is the most important and limiting nutrient for rice crop growth and yield which is required in higher amounts compared to other nutrients (**Djaman et al., 2018**). Nitrogen (N) influences rice yield by playing major role in the photosynthesis, biomass accumulation, effective tillering, and spikelets formation. (**Yoshida et al., 2006**). Keeping in view the above facts, the experiment was conducted to find out the

## **“Evaluation of different varieties and nitrogen levels on growth and yield of Black Rice (*Oryza sativa* L.)”.**

### **MATERIAL AND METHODS:**

The field experiment was conducted during *khariif* season 2022 at the Crop Research Farm, Department of Agronomy, Naini Agricultural Institute, Sam Higginbottom University of Agriculture Technology And Sciences, Prayagraj (U.P.). The soil of the experimental field was sandy loam in texture, nearly neutral in soil reaction (pH 8), organic carbon level in medium condition (0.87%), medium available N (225 Kg/ha), high in available P (41.8 kg/ha) and medium available K (261.2 kg/ha). The experiment was laid out in Randomized Block Design with three replication having two factors viz. three black rice varieties (Chakhao poireiton, Chakhao Amubi, Chakhao Angangba) and three Nitrogen levels (60, 70, 80 kg/ha). The three varieties of black rice with 118days duration was used for this experiment. Twenty-five days old seedling were transplanted at plant spacing 20x15 cm with 2-3 seedling per hill. The treatment combinations are treatment 1- Chakhao Poireiton + Nitrogen (60 kg/ha), treatment 2- Chakhao Poireiton + Nitrogen (70 kg/ha), treatment 3- Chakhao Poireiton + Nitrogen (80 kg/ha), treatment 4- Chakhao Amubi + Nitrogen (60 kg/ha), treatment 5- Chakhao Amubi + Nitrogen (70 kg/ha), treatment 6- Chakhao Amubi + Nitrogen (80 kg/ha), treatment 7- Chakhao Angangba + Nitrogen (60 kg/ha), treatment 8- Chakhao Angangba + Nitrogen (70 kg/ha), treatment 9- Chakhao Angangba + Nitrogen (80 kg/ha). All agronomic practices are followed in order in the crop period. Experimental data collected was subjected to statistical analysis of variance (ANOVA) as outlined by **Gomez and Gomez (1984)**. Critical Difference (CD) values were calculated wherever the ‘F’ test was found significant at 5 percent level.

### **RESULTS AND DISCUSSION**

#### **Growth parameters**

##### **Plant height(cm)**

The data revealed that a significantly higher plant height (142.53) was recorded in treatment 3 [Chakhao poireiton + Nitrogen (80kg/ ha)]. However, treatment 2 [Chakhao poireiton + Nitrogen (70kg/ ha)] was found to be statistically at par with treatment 3 [Chakhao poireiton + Nitrogen (80kg/ ha)] (Table 1). The significant and higher plant height was observed with the variety Chakhao Poireiton might be due to differences in the genetic makeup of the varieties and their

response ability to different levels of fertilizer applied. This finding is similar reported by **Ndaeyo *et al.* (2008)**. Further increase in plant height nitrogen (80 kg/ha) might be due to enough nutrition (NP) in termed of possible increase in nutrient absorption capacity of plant as a result of better root development and increased translocation of carbohydrates from source to growing points. These results confirmed the finding of **Bhiah *et al.* (2010)**.

### **Number of tillers/hill**

The data revealed that a significantly maximum number of tillers/hill (13.97) was recorded in treatment 3 [Chakhao poireiton + Nitrogen (80kg/ ha)] (Table 1). The significant and maximum number of tillers/hill was observed with the variety Chakhao Poireiton might be due to varietal variation and their genetic characters attributed to greater translocation of photosynthates to the reproductive part of the plant. Similar result was reported by **Thenmozhi *et al.*, (2013)**. Further increase number of tillers was observed with the application of N (80 kg/ha) might be due to increase levels of nitrogen favours greater absorption of nutrients resulting in rapid expansion of foliage, better accumulation of photosynthates. Similar result was reported by **Murthy *et al.* (2015)**.

### **Plant Dry Weight (g/hill)**

The data revealed that a significantly higher plant dry weight (32.93 g) was recorded in treatment 3 [Chakhao poireiton + Nitrogen (80kg/ ha)]. However, treatment 2 [Chakhao poireiton + Nitrogen (70kg/ ha)] was found to be statistically at par with treatment 3 [Chakhao poireiton + Nitrogen (80kg/ ha)] (Table 1). The significant and higher plant dry weight was observed with the variety Chakhao Poireiton of black rice might be due to their genetic potential and differential plant height and more number of tillers/hill. Similar result was reported by **Dillep *et al.* (2018)**. Further higher plant dry weight was observed with Nitrogen (80 kg/ha) might be due to nitrogen is the main component of the protoplasm involves in various metabolic processes *viz.* photosynthesis, stimulation of cell division and elongation, resulted increased plant dry weight. Similar result was reported by **Meena *et al.* (2003)**.

### **Crop Growth Rate (g/m<sup>2</sup>/day)**

The data revealed that a significantly higher Crop Groeth Rate (13.50 g/m<sup>2</sup>/day) was recorded in treatment 9 [Chakhao Angangba+ Nitrogen (80 Kg/ha)] as compared to other treatments. However, the treatment 2 [Chakhao Poireiton + Nitrogen (70 Kg/ha)], treatment 3 [Chakhao Poireiton + Nitrogen (80 Kg/ha)] and treatment 7 [Chakhao Angangba+ Nitrogen (60 Kg/ha)]

were found statistically at par with treatment 9 [Chakhao Angangba+ Nitrogen (80 Kg/ha)] (Table 1). The significant and higher crop growth rate was observed with the variety Chakhao Angangba might be due to the natural endowments of crop cultivars to optimally utilize available nutrients and subsequently partition its photosynthates. Similar result was reported by **Ndon and Ndaeyo (2001)**. Further significant and higher crop growth rate was observed with Nitrogen (80 Kg/ha) might be due to the role of nitrogen in production and translocation of cytokinin from the root to the shoots increases cell division rate and the growth of rice with the increased in nitrogen level. Similar result was reported by **Azarpour *et al.*, 2014**.

#### **Relative Growth Rate (g/g/day)**

The data revealed that a significantly higher Relative Growth Rate (0.0169 g/g/day) was observed and recorded in treatment 9 [Chakhao Angangba + Nitrogen (80 Kg/ha)] as compared to other treatments (Table 1). The significant and higher relative growth rate was observed with the variety Chakhao Angangba might be due to increase in photosynthetic tissues and faster vegetative growth, which may have leads to higher relative growth rate. Similar result was report by **Azarpour *et al.* (2014)**. Further higher relative growth rate with nitrogen (80 kg/ha) might be due to timely and adequate amount of nitrogen supplied during initial crop growth period. Similar result was reported by **Sathiya and Ramesh (2009)**.

### **POST-HARVEST OBSERVATION**

#### **Panicle length (cm)**

The data revealed that a significantly higher panicle length (30.1 cm) was recorded in treatment 3 [Chakhao Poireiton + Nitrogen (80 kg/ha)]. However, the treatment 2 [Chakhao Poireiton + Nitrogen (70 kg/ha)] and treatment 1 [Chakhao Poireiton + Nitrogen (60 kg/ha)] were found statistically at par with treatment 3 [Chakhao Poireiton + Nitrogen (80 kg/ha)] (Table 2). The significant and higher panicle length was observed with the variety Chakhao Poireiton might due to genetic characters of the varieties primarily influenced by the heredity. Similar result was reported by **Diaz *et al.* (2000)**. Further significant and higher panicle length with nitrogen (80 kg/ha) might be due to higher nitrogen application influenced better growth characters which ultimately resulted in higher production and translocation of photosynthates towards panicle. Similar result was reported by **Dahi and Singh (2018)**.

#### **Number of grains/panicle**

The data revealed that a significantly maximum number of grains/panicle (153.60) was recorded in treatment 2 with application of [Chakhao Poireiton + Nitrogen (70 kg/ha)]. However, the treatment 1 [Chakhao Poireiton + Nitrogen (60 kg/ha)] and treatment 3 [Chakhao Poireiton + Nitrogen (80 kg/ha)] were found statistically at par with treatment 2 [Chakhao Poireiton + Nitrogen (70 kg/ha)] (Table 2). The significant and higher number of grains/panicle was observed with the variety Chakhao Poireiton might be due to their genetic makeup and variation in photosynthetic assimilate accumulation. Similar result was reported by **Chamely *et al.*, (2015)**. Further significantly higher number of grains/panicle was observed with Nitrogen (80 kg/ha) might be due to high nitrogen level, greater assimilation of photosynthates resulting in higher number of filled grains. Similar result was reported by **Uddin *et al.* (2013)**.

### **Test weight (g)**

The data revealed that a significantly higher Test Weight (24.01 g) was observed in treatment 3 with application of [Chakhao Poireiton + Nitrogen (80 Kg/ha)]. However, the treatment 1 [Chakhao Poireiton + Nitrogen (60 Kg/ha)] and treatment 6 [Chakhao Amubi + Nitrogen (80 Kg/ha)] were found statistically at par with treatment 3 [Chakhao Poireiton + Nitrogen (80 Kg/ha)] (Table 2). The significant and higher test weight was observed with the variety Chakhao poireiton might be due to genetic character and least influenced by environment, resulted higher test weight. Similar result was reported by **Saha *et al.*, (2017)**. Further significant and higher test weight was observed with Nitrogen (80 kg/ha) might be due to the adequate supply of nitrogen, probably favoured the proper cellular activities during panicle formation and development which led to increase higher test weight. Similar result was reported by **Aparna *et al.* (2022)**.

### **Grain yield (t/ha)**

The data revealed that a significantly higher grain yield (5.67 t/ha) was recorded in treatment 3 [Chakhao Poireiton + Nitrogen (80 Kg/ha)]. However, the treatment 2 [Chakhao Poireiton + Nitrogen (70 Kg/ha)] and treatment 1 [Chakhao Poireiton + Nitrogen (60 Kg/ha)] were found statistically at par with treatment 3 [Chakhao Poireiton + Nitrogen (80 Kg/ha)] (Table 2). The significant and higher grain yield was observed with the variety Chakhao Poireiton might be due to higher number of effective tillers attributed to higher affinity of this variety and yield contributes associate with favourable weather condition followed by differential ability of cultivars and photoassimilate, which resulted in the higher grain yield. Similar result was reported by **Anil and Siddi (2020)**. Further significantly higher grain yield was observed with the application of nitrogen (80 kg/ha) might due to the availability more water enhanced nutrient

availability which improved nitrogen and other macro and micro-elements absorption enhancing the production. Similar result was reported by **Bhiah and Aljbori (2020)**.

### **Straw yield (t/ha)**

The data revealed that a significantly higher straw yield (11.27 t/ha) was observed and recorded in treatment 3 with the application of [Chakhao Poireiton + Nitrogen (80 Kg/ha)]. However, the application of [Chakhao Poireiton + Nitrogen (70 Kg/ha)] was found statistically at par with the application of [Chakhao Poireiton + Nitrogen (80 Kg/ha)] (Table 2). The significant and higher straw yield was observed with the variety Chakhao Poireiton might be due the production of more tillers and high accumulation of dry matter, resulted increased in straw yield. Similar result was reported by **Srimathi and Subramanian (2022)**. Further significantly higher straw yield was observed with nitrogen (80 Kg/ha) might be due to nitrogen levels improved vegetative growth and increase in the chlorophyll concentration in leaf of plants. Similar result was reported by **Biswas *et al.* (2020)**.

### **Harvest index (%)**

The highest harvest index (34.53) was found in treatment 3 [ Chakhao Poireiton + nitrogen (80 kg/ha)], though it was non-significant (Table 2).

### **Economics**

The result showed that [Table 3] maximum gross return (2,38,070.00 INR/ha), net return (1,91,800.00 INR/ha) and B:C ratio (3.23) was recorded in treatment 3 with application of Chakhao Poireiton + Nitrogen 80 kg/ha (Table 2). Net returns and benefit cost ratio (3.23) was recorded with the variety Chakhao Poireiton may be due to its superiority than the rest of the varieties, therefore increased in grain yield, resulted higher benefit cost ratio. Further, higher net return was found with application nitrogen (80 kg/ha) might be due to variation in levels of nitrogen monitoring the major inputs, which ultimately increases economic yield with higher net return. Similar result was reported by **Mishra *et al.* (2015)**.

### **CONCLUSION**

Based on above findings, it can be concluded that the use of various levels of nitrogen improved the growth parameters and yield attributes of Black rice varieties. Maximum gross return, net return and benefit cost-ratio was recorded with application of Chakhao Poireiton + Nitrogen 80 kg/ha.

## REFERENCES

1. **Anil D. and Sreedhar Siddi (2020)**. Performance of pre released rice (*oryza sativa* L.) genotypes under different sowing windows in rabi season. *Internation Journal of Environment and Climate Change*. **10**(10): 43-51.
2. **Aparna Vijaya B, Anny Mrudhula K, Prasad PVN and Ravi Babu M. (2022)**. Effect of planting geometry and nitrogen levels on growth parameters, yield attributes and yield of black rice. *The Pharma Innovation Journal*. **11**(9): 2260-2264.
3. **Azarpour E, Motamed MK, Moraditochae M, Bozorgi HR.(2011)**. Effect of nitrogen fertilizer and nitroxin biofertilizer manageent on growth analysis and yield of rice cultivars (Iran). *World Applied Sciences Journal* **14**(2),193-198.
4. **Azarpour, E., Moraditochae, M., & Bozorgi, H. (2014)**. Effect of nitrogen fertilizer management on growth analysis of rice cultivars. *International Journal of Biosciences*, **4**, 35-47.
5. **Bai ZG. 2019**. Effects of Nitrogen Fertilizer Operation on Nitrogen Metabolism of Rice and Nitrogen Utilization Rate in Rice Field. Ph.D. Thesis, Chinese Academy of Agricultural Sciences, Guangzhou, China.
6. **Bhiah Karim M. and Albori Flayeh A. (2020)**. Response of different growth and yield parameters of four black rice lines to recommended doses of minerals fertilizers. *Plant Archives*. **20**(2): 5932-5936.
7. **Bhiah, K.M., C. Guppy, P. Lockwood and R. Jessop (2010)**. Effect of potassium on rice lodging under high nitrogen nutrition. Brisbane (AU): In 19<sup>th</sup> World Congress of Soil Science “*Soil Solution for a Changing World*, 1-6.
8. **Biswas Papia, Jitendra Singh Bohra and Nitesh Kumar. (2020)**. Effect of nutrient management on growth, yield and yield attributes of rice (*oryza sativa* L.) *International Journal of Current Microbiology and Applied Sciences*. **9**(3): 3136-3146.
9. **Borah N, Athokpam FD, Semwal RL, Gorkotti SC. Chakhao (blackrice; Oryza sativa L.)**. The culturally important and stress-tolerant traditional rice variety of Manipur. *Indian Journal of Traditional Knowledge*. 2018; **17**(4): 176-180.
10. **Chamley SG, N Islam, S Hoshain, MG Rabbani, MA Kader, MA Salam. (2015)**. Effect of variety and nitrogen rate on the yield performance of boro rice. *Progressive Agriculture* **26**(1): 6-14.

11. **Dahi, Singh (2018)**. Effect of different N levels on rice growth parameters, *IJCS*. 3:35-40.
12. **Diaz SH, Castro R, Morejon R (2000)**. Morphoanatomic characterization of varieties of rice. *Instituto Nacional de ciencias Agrícolas*, Gaveta Postall, San Jose, de las, Lajsa, La Habna. **21**(3):81-86
13. **Dileep K, Pasupalak S, Baliarsingh A (2018)**. Effect of establishment methods and sowing time on growth and yied of rice varieties (*Oryza sativa* L.). *The Pharma Innivation Journal*. **7**(4):904-907.
14. **Djaman, K. M.N. (2018)**. Effect of nitrogen fertilizer dose and application timing on yield and nitrogen use efficiency of irrigated hybrid rice under semi-arid conditions. *Journal of Agricultural Science and Food Research*, **9**(2).
  - a. [eands.dacnet.nic.in/PDF/Agricultural\\_Statistics\\_At\\_Glance-2021.pdf](https://eands.dacnet.nic.in/PDF/Agricultural_Statistics_At_Glance-2021.pdf)
15. **GOI. (2021)**. Agricultural statistics at a glance: *Ministry of Agriculture & Farmers Welfare, Government of India*.
16. **Gomez, K.A., Gomez, A.A., (1984)**. Statistical Procedure for Agricultural Research 2<sup>nd</sup> Edition., Jhon Wiley and Sons, New York, 680.
17. **Li S, S Pu, F Deng, LWang, H Hu, S Liao, W Li and W Ren. (2019)**. Influence of optimized nitrogen management on the quality of medium hybrid rice under different ecological conditions (Article). *Chinese Journal of Eco-Agriculture* **27**(7): 1042–1052.
18. **Meena , S., Singh, S., & Shivay, Y. (2003)**. Response of hybrid rice to nitrogen and potassium application in sandy clay-loam soils. *Indian Journal of Agricultural Sciences*, **73**(1), 8-11.
19. **Mishra Dushyant, J.D. Sharma, Asheesh Kumar Pandey, Rajeeva Kumar Mishra, U.N. Shukla and Jitendra Kumar (2015)**. Effect of Nitrogen levels on yield, quantity and economics of rice (*Oryza sativa* L.) Varieties. *International Journal of Agricultural Sciences*. **11**(1):0973-1903.
20. **Murthy Dakdhima KM, Upendra Rao A, Vijay D, Sridhar TV. (2015)**. Effect of levels of nitogen, phosphorus and potassium on performance of rice. *Indian Journal of Agricultural Research*. **49**(1):83-87.
21. **Ndaeyo, N.U, Iboko, K.U., Harry, G.I., and Edem, S.O. (2008)**. Growth and yield performances of some upland rice (*Oryza sativa* L.) cultivars as influenced by varied rates

- of NPK (15:15:15) fertilizer on an ultisol. *Journal of Tropical Agriculture, Food, Environment and Extension*, **7**(3), 249-255.
- 22. Ndon, B.A., and Ndaeyo, N.U. (2001).** Evaluation of some cowpea (*Vigna unguiculata* (L.) cultivars adaptable to acid soils of Udyo, Southeastern Nigeria. *Global Journal of Pure and Applied Sciences*, **7**(3), 437-441.
- 23. Saha Biswajit, Parimal Panda, Partha Sarathi Patra, Ranajit Panda, arindam Kundu, A.K. Singha Roy and Nabakishor Mahato (2017).** *International Journal of Current Microbiology and Applied Sciences*. **6**(7):2408-2418.
- 24. Satiya, K and Ramesh, T. (2009).** Effect of split application of nitrogen on growth and yield of aerobic rice. *Asian Journal of Experimental Sciences*. **23**(1): 303-306.
- 25. Shrestha Jiban, Tika Bahadur Karki and Mohammad Anwar Hossain (2022).** Application of Nitrogenous Fertilizer in Rice Production. *Journal of Nepal Agricultural Research Council*. **18**(5):2392-4543.
- 26. Srimathi K and Subramanian E. (2022).** Evaluation of tillering behaviour and yielding ability of different rice varieties under unpuddled conditions. *Current Journal of Applied Science and Technology*. **41**(28): 26-32.
- 27. Takashi I, Bing X, Yoichi Y, Masaharu N, Tetsuya K. (2001).** Antioxidant activity of anthocyanin extract from purple black rice. *Journal of Medicinal Food*. **4**:211-218.
- 28. Tayefe, M., Akif, G., Amiri, E., & Azin, N.Z. (2014).** Effect of nitrogen on rice yield, yield components and quality parameters. *African Journal of Geology, Ecology, and Landscapes*, **13**(1), 91-105.
- 29. Thenmozhi P, Rajasekaran C (2013).** Genetic diversity and relationship among 40 rice accessions from North-Eastern Zone of Tamil Nadu using morphological and SSR markers. *Research Journal of Biotechnology*, **8**(9):32-41.
- 30. Uddin S, M.A.R. Sarkar and M.M.Rahman (2013).** Effect of nitrogen and potassium on yield of dry direct seeded rice cv. *NERICA1* in aus season. Vol., **4**,69-75.
- 31. USDA, 2022.** Foreign Agriculture service, World Agricultural production.  
<https://apps.fas.usda.gov>.
- 32. Yoshida, H., Horie, T., & Shiraiwa, T. (2006).** A model explaining genotypic and environmental variation of rice spikelet number per unit area measured by cross-locality experiments in Asia. *Field Crops Research*, **97**(2-3), 337-343.

S.No.	Treatment	100 DAT			80-100DAT	
		Plant Height (cm)	Number of tillers/hill	Plant dry weight (g/hill)	Crop growth rate (g/ m <sup>2</sup> / day)	Relative growth rate (g/ g/ day)
1.	Chakhao Poireiton + Nitrogen 60 kg/ha	140.90	9.77	31.77	12.11	0.0128
2.	Chakhao Poireiton + Nitrogen 70 kg/ha	141.40	11.97	31.93	13.11	0.0138
3.	Chakhao Poireiton + Nitrogen 80 kg/ha	142.53	13.97	32.49	12.39	0.0128
4.	Chakhao Amubi + Nitrogen 60 kg/ha	138.13	9.57	30.68	11.80	0.0127
5.	Chakhao Amubi + Nitrogen 70 kg/ha	139.03	10.03	30.33	11.43	0.0136
6.	Chakhao Amubi + Nitrogen 80 kg/ha	139.43	9.83	29.85	11.18	0.0138
7.	Chakhao Angangba + Nitrogen 60 kg/ha	136.27	11.67	29.67	12.21	0.0147
8.	Chakhao Angangba + Nitrogen 70kg/ha	137.57	9.67	28.97	11.89	0.0119
9.	Chakhao Angangba + Nitrogen 80kg/ha	138.43	11.73	29.40	13.50	0.0169
	F-test	S	S	S	S	S
	SEm(±)	0.74	0.82	0.30	0.65	0.0008
	CD (p=0.05)	1.56	1.74	0.64	1.38	0.0017

Table 1: Response of nitrogen on growth of Black rice varieties.

Table 2: Response of nitrogen on yield and yield attributes of Black rice varieties

<b>S. No.</b>	<b>Treatments</b>	<b>Panicle length (cm)</b>	<b>Number of grains/panicle</b>	<b>Test weight (g)</b>	<b>Grain yield (t/ha)</b>	<b>Straw yield (t/ha)</b>	<b>Harvest index (%)</b>
1.	Chakhao Poireiton + Nitrogen 60 kg/ha	29.80	150.13	23.12	5.17	9.80	33.47
2.	Chakhao Poireiton + Nitrogen 70 kg/ha	29.93	153.47	22.53	5.13	10.20	33.46
3.	Chakhao Poireiton + Nitrogen 80 kg/ha	30.1	153.60	24.01	5.67	11.27	34.53
4.	Chakhao Amubi + Nitrogen 60 kg/ha	28.73	133.00	22.03	4.50	8.83	33.75
5.	Chakhao Amubi + Nitrogen 70 kg/ha	28.90	131.53	21.80	4.90	9.33	34.43
6.	Chakhao Amubi + Nitrogen 80 kg/ha	28.53	134.60	23.20	5.00	10.80	31.64
7.	Chakhao Angangba + Nitrogen 60 kg/ha	28.27	135.47	21.19	3.50	9.17	27.62
8.	Chakhao Angangba + Nitrogen 70kg/ha	28.13	134.80	21.20	3.33	9.83	25.30
9.	Chakhao Angangba + Nitrogen 80kg/ha	28.17	135.40	22.70	3.67	10.07	26.71
	F-test	S	S	S	S	S	NS
	SEm(±)	0.18	1.06	0.58	0.31	0.46	1.72
	CD (p=0.05)	0.39	2.25	1.24	0.66	0.97	-

<b>S. No.</b>	<b>Treatment combinations</b>	<b>Cost of cultivation (INR/ha)</b>	<b>Gross return (INR/ha)</b>	<b>Net return (INR/ha)</b>	<b>B:C ratio</b>
1.	Chakhao Poireiton + Nitrogen 60 kg/ha	46110	216600	170490	3.16
2.	Chakhao Poireiton + Nitrogen 70kg/ha	46190	215400	169210	3.05
3.	Chakhao Poireiton + Nitrogen 80kg/ha	46270	238070	191800	3.23
4.	Chakhao Amubi + Nitrogen 60 kg/ha	46110	188830	143120	2.91
5.	Chakhao Amubi + Nitrogen 70 kg/ha	46190	205330	159540	2.75
6.	Chakhao Amubi + Nitrogen 80 kg/ha	46270	210800	164930	2.35
7.	Chakhao Angangba + Nitrogen 60 kg/ha	46110	149170	103460	1.78
8.	Chakhao Angangba + Nitrogen 70 kg/ha	46190	143030	97240	1.84
9.	Chakhao Angangba + Nitrogen 80 kg/ha	46270	156870	111000	1.92

Table 3. Response of nitrogen on yield and yield attributes on post-harvest yield of Black rice varieties.

UNDER PEER REVIEW