

## EFFECT OF INTEGRATED NUTRIENT MANAGEMENT ON YIELD AND ECONOMICS OF *KHARIF* SORGHUM. (*Sorghum bicolor* L.)

### Abstract

The field experiment was laid out at Sorghum Research Station, VNMKV, Parbhani. Dist. Parbhani during *kharif* season of 2015 to study the “Integrated Nutrient Management in *Kharif* Sorghum (*Sorghum bicolor* L.)”. The experiment was laid out in randomized block design with three replications and variety CSH -16 as test crop along with nine (09) treatment combinations. The quantity of organic and inorganic fertilizer dose was calculated and applied in the plots as per the treatments. Sowing was done on 21<sup>th</sup> June 2015 by seeds are dibbling and distance between 45cm x 15 cm. The crop was harvested at physiological maturity and data on yield attributes and yield were recorded. The study revealed that the application of 75% RDN through inorganic fertilizer + 25% RDN through vermicompost + seed treatment with PSB + *Azospirillum* (T<sub>8</sub>) gave significantly higher GMR (76,068 ₹ ha<sup>-1</sup>), NMR (37,999 ₹ ha<sup>-1</sup>) and B:C ratio (2.07) over rest of the treatments, however it was at par with application of 75 % RDN through inorganic fertilizer + 25% RDN through FYM + seed treatment with PSB + *Azospirillum* (T<sub>7</sub>).

**Key Words** : INM, Economics, Sorghum, Yield

### Introduction

Sorghum (*Sorghum bicolor* (L.) is an important cereal crop in India popularly known as ‘Jawar’ and large size of among other grain millets is called ‘Great millet’. It is probably originated in East Central Africa and it was introduced to India from East Africa in the year 1500 BC. Sorghum is the 5<sup>th</sup> most important cereal crop in the world after rice, wheat, maize and barley. . The advantage of this cereal crop it can be cultivated in both *Kharif* and *Rabi* season. In the world, USA is the largest producer of sorghum occupying 20.03 percent of area with 16.41 per cent production. India is a major sorghum growing country in the

world. It stood second largest crop till green revolution but after green revolution, it occupies the third place among food grains after rice and wheat. Maharashtra is the largest sorghum growing state in the area nearly 6.80 million hectares with an annual production of 5.85 million tonnes and productivity of 860 kg ha<sup>-1</sup>. The area, production, productivity of *Kharif* jowar in Marathwada region was 3.66 lakh ha, 1.97 lakh tonnes and 539 kg ha<sup>-1</sup>, respectively. It is observed that area of *Kharif* sorghum is decreasing day by day while the production and productivity is in increasing trend. Increase in productivity is only due to high yielding varieties and advanced technology. According to recent estimation of planning commission of India, 30-35 million tonnes of food grains and 14-15 million tonnes of nutrients require for commercial crops.

Hence integrated nutrient management approach is the option today's to mitigate the gap of nutrients in Indian agriculture. Integrated nutrient management is the alternative to increase the productivity through proper management. Keeping in view this fact, the present investigation was undertaken to find out a suitable nutrient management system to boost yield of *kharif* sorghum.

Integrated and balanced use of nutrients through inorganic and organic sources like FYM, vermicompost and biofertilizers is pre-requisite to produce to sustain soil fertility, supply of nutrients at an optimum level and to produce maximum crop yield with minimum inputs (Dahiphale *et al.*, 2003). According to Sarkar (2000), the basic principal of INM is the maintenance of soil fertility by using all sources of nutrient. In plant nutrition, organic matter levels of a soil are a key property that decides the availability status of essential nutrients. The INM however, helps to maintain the productivity of soil and improves fertilizer use efficiency. Thus, it economizes the use of chemical fertilizers by influencing the yield of *kharif* crop besides nutrients availability, FYM and vermicompost also improves soil physical characteristics such as structures,

porosity and water holding capacity. Vermicompost when applied in conjunction with biofertilizer, supplies energy to beneficial microorganism including *Azotobacter* and PSB

## **MATERIALS AND METHOD**

The experiment was conducted during *Kharif*, 2015 at AICSIP, Sorghum Research Station, VNMKV, Parbhani (MS). Parbhani located at 19° 16' N latitude and 76° 41' East longitudes and has sub – tropical climatic conditions. Parbhani is grouped under assured rainfall zone. The normal rainfall of this region is around 954.9 mm, precipitating mostly between mid June–mid Novembers. The average maximum and minimum temperature recorded 31.2°C and 22.6°C, respectively. The soil was medium deep black and well drained. The topography of the experimental field was fairly uniform and leveled. Soil samples up to 30 cm were randomly collected from different locations of field before start of the experiment during *Kharif*, 2015.

The precipitation is assured for *Kharif* crops so need of providing life saving irrigation. The experiment was laid out in Randomized block design with three replications constituting nine treatments of integrated nutrient management in *Kharif* sorghum. The treatments were (T<sub>1</sub>):-100% RDN through inorganic fertilizer, (T<sub>2</sub>):- 50% RDN through inorganic fertilizer + 50% RDN through FYM, (T<sub>3</sub>):- 75% RDN through inorganic fertilizer + 25% RDN through FYM, (T<sub>4</sub>) 50% RDN through inorganic fertilizer + 50% RDN through vermicompost, (T<sub>5</sub>) 75% RDN through inorganic fertilizer + 25% RDN through vermicompost, (T<sub>6</sub>):- 50% RDN through inorganic fertilizer + 25% RDN through FYM + 25% RDN through Vermicompost, (T<sub>7</sub>):- 75% RDN through inorganic fertilizer + 25% RDN through FYM+ seed treatment with Microbial fertilizers like PSB + *Azospirillum*, (T<sub>8</sub>):- 75% RDN through inorganic fertilizer + 25% RDN through Vermicompost + seed treatment with Microbial fertilizers like PSB + *Azospirillum* and (T<sub>9</sub>) 75% RDN

through inorganic fertilizer + seed treatment with Microbial fertilizers like PSB + *Azospirillum*.

The sorghum cultivar CSH-16 was obtained from AICSIP Sorghum Research Station, VNMKV, Parbhani. Sowing was done on 21<sup>st</sup> June, 2015 by dibbling 2-3 seed at each hill on a recommended spacing of 45 cm x 15cm. Seeds were treated with Emamectin benzoate @ 3 g kg<sup>-1</sup> seeds in order to protect the crop from shoot-fly attack. Inorganic sources *viz.*, Urea, SSP, MOP and organic sources *viz.*, FYM, vermicompost, bio fertilizers were applied to respective plots as per the recommendation uniformly in the lines opened for sowing. During early growth stages of crop, to control of shoot fly (*Atherigona soccata*) and stem borer (*Chilopartellus*) in addition to seed treatment with Emamectin benzoate two sprayings of Quinolphos and Emamectin benzoate were taken. Five plants from each net plot were randomly selected and labeled for taking biometric observations at pre-harvest and post harvest growth stages. Available nitrogen, phosphorus and potassium were determined by standard methods. Data obtained on various variables were analyzed by “analysis of variance method” (Panse and Sukhatme, 1967).

## **RESULTS AND DISCUSSION**

### **Yield**

The application of 75% RDN through inorganic fertilizer + 25% RDN through vermicompost + seed treatment with PSB + *Azospirillum* (T<sub>8</sub>) gave significantly higher grain yield (2356.89 Kg ha<sup>-1</sup>), fodder yield (5450.31 Kg ha<sup>-1</sup>) and biological yield (7807.20 Kg ha<sup>-1</sup>) over rest of the treatments, however it was at par with application of 75 % RDN through inorganic fertilizer + 25% RDN through FYM + seed treatment with PSB + *Azospirillum* (T<sub>7</sub>) and (T<sub>1</sub>) 100% RDN

through inorganic fertilizer. Similar results were also reported by Kalibhavi *et al.* (1996) and Gawai and Pawar (2007). However highest harvest index recorded in treatment 75% RDN through inorganic fertilizer + 25% RDN through vermicompost + seed treatment with PSB + *Azospirillum* (29.41) whereas mean harvest index was (28.52). Test weight does not influence significantly among all the integrated nutrient management treatment.

## **Economics**

### **Cost of cultivation**

Data presented in Table 2 implies total cost of sorghum cultivation. The mean value of total cost was 36552 ₹ ha<sup>-1</sup>. Glimpse of the Table 2 insinuated that, the cost of cultivation was higher under application of 50% RDN through inorganic fertilizer + 50% RDN through vermicompost (T<sub>4</sub>) (38500 ₹ ha<sup>-1</sup>) followed by (T<sub>2</sub>) 50% RDN through inorganic fertilizer + 50% RDN through FYM and lowest cost of cultivation (35200 ₹ ha<sup>-1</sup>) was under treatment (T<sub>1</sub>) 100% RDN through inorganic fertilizer.

### **Gross returns**

Data on gross return are presented in Table 2. The mean gross returns were (612367 ₹ ha<sup>-1</sup>)Scrutiny of the Table 1 implies that the application of 75% RDN through inorganic fertilizer + 25% RDN through vermicompost + seed treatment with PSB + *Azospirillum* (T<sub>8</sub>) recorded significantly higher gross returns ₹ha<sup>-1</sup> *i.e.* (73599 ₹ ha<sup>-1</sup>) over rest of the treatments. But it was at par with (T<sub>7</sub>) 75 % RDN through inorganic fertilizer + 25% RDN through FYM + seed treatment with PSB + *Azospirillum* and (T<sub>1</sub>) 100% RDN through inorganic fertilizer.

Application of 75% RDN through inorganic fertilizer + seed treatment with PSB + *Azospirillum* (T<sub>9</sub>) (47718 ₹ ha<sup>-1</sup>) is recorded significantly lower gross returns ₹ha<sup>-1</sup>. Similar results were also reported by Jat *et al.* (2013).

### Net returns

Data presented in Table 2 revealed that the mean net return of sorghum was 24715 ₹ ha<sup>-1</sup>. The application of 75% RDN through inorganic fertilizer + 25% RDN through vermicompost + seed treatment with PSB + *Azospirillum* (T<sub>8</sub>) recorded significantly higher net returns ₹ha<sup>-1</sup> i.e. (37999 ₹ ha<sup>-1</sup>) over rest of the treatments, but it was at par with (T<sub>7</sub>) and (T<sub>1</sub>). Application of 75% RDN through inorganic fertilizer + seed treatment with PSB + *Azospirillum* (T<sub>9</sub>) (12818 ₹ ha<sup>-1</sup>) is recorded significantly lower net returns ₹ha<sup>-1</sup>. Similar results were also reported by Mahajan *et al.* (1994).

### B:C ratio

Data on B:C ratio as influenced by various treatments are presented in Table 2. The mean B:C ratio was (1.68). The application 75% RDN through inorganic fertilizer + 25% RDN through vermicompost + seed treatment with PSB + *Azospirillum* (T<sub>8</sub>) was found highest (2.07) B:C ratio over rest of the treatments. But it was at par with (T<sub>7</sub>) and (T<sub>1</sub>). Similar results were also reported by Nawale *et al.* (2007).

## CONCLUSIONS

On the basis of present investigation following conclusions can be drawn.

1. In *kharif* sorghum the application of 75% RDN through inorganic fertilizer along with 25% organic sources + seed treatment with *Azospirillum* and PSB produced higher grain and fodder yield and also improve the soil fertility.
2. The application of 75% RDN through inorganic fertilizer + 25% RDN through vermicompost + seed treatment with PSB + *Azospirillum* recorded higher gross monetary returns (76,068 ₹ ha<sup>-1</sup>), net monetary returns (37,999 ₹ ha<sup>-1</sup>) and B:C ratio (2.07).

Above conclusions are based on single season research finding and it needs further confirmation by repeating the trial for at least one more season.

**Table no. 1:** Grain yield, fodder yield, biological yield and harvest index of sorghum as influenced by different treatments.

Treatment	Grain yield (Kg ha <sup>-1</sup> )	Fodder yield (Kg ha <sup>-1</sup> )	Biological yield (Kg ha <sup>-1</sup> )	Harvest Index (%)
T <sub>1</sub> : 100% RDN through inorganic fertilizer	2190.65	5303.00	7493.65	29.23
T <sub>2</sub> : 50% RDN through inorganic fertilizer + 50% RDN through FYM	1578.28	4013.03	5591.30	28.23
T <sub>3</sub> : 75% RDN through inorganic fertilizer + 25% RDN through FYM	1662.45	4208.73	5871.18	28.32
T <sub>4</sub> : 50% RDN through inorganic fertilizer + 50% RDN through vermicompost	1698.22	4229.78	5928.00	28.72
T <sub>5</sub> : 75% RDN through inorganic fertilizer + 25% RDN through vermicompost	1752.94	4492.82	6245.76	28.06
T <sub>6</sub> : 50% RDN through inorganic fertilizer + 25% RDN through FYM + 25% RDN through vermicompost	1763.46	4566.48	6329.93	27.86
T <sub>7</sub> : 75% RDN through inorganic fertilizer + 25% RDN through FYM+ seed treatment with PSB + <i>Azospirillum</i>	2220.11	5366.14	7586.24	29.27
T <sub>8</sub> : 75% RDN through inorganic fertilizer + 25% RDN through vermicompost + seed treatment with PSB + <i>Azospirillum</i>	2243.25	5387.18	7630.43	29.41
T <sub>9</sub> : 75% RDN through inorganic fertilizer + seed treatment with PSB + <i>Azospirillum</i>	1397.30	3667.91	5065.21	27.59
S.E. ±	19.49	80.59	79.8963	-
C.D. at 5 %	58.66	242.59	240.511	-
<b>G. Mean</b>	<b>1834.07</b>	<b>4581.67</b>	<b>6415.75</b>	<b>28.52</b>

**Table 2. Cost of cultivation, gross and net return (₹ ha<sup>-1</sup>) and B:C ratio of sorghum as influenced by different treatments**

<b>Treatment</b>	<b>Cost of cultivation (₹ ha<sup>-1</sup>)</b>	<b>Gross returns (₹ ha<sup>-1</sup>)</b>	<b>Net returns (₹ ha<sup>-1</sup>)</b>	<b>B : C ratio</b>
T <sub>1</sub> : 100% RDN through inorganic fertilizer	35200	72126	36926	2.05
T <sub>2</sub> : 50% RDN through inorganic fertilizer + 50% RDN through FYM	38200	53118	14918	1.39
T <sub>3</sub> : 75% RDN through inorganic fertilizer + 25% RDN through FYM	36200	55841	19641	1.54
T <sub>4</sub> : 50% RDN through inorganic fertilizer + 50% RDN through vermicompost	38500	56626	18126	1.47
T <sub>5</sub> : 75% RDN through inorganic fertilizer + 25% RDN through vermicompost	36500	59211	22711	1.62
T <sub>6</sub> : 50% RDN through inorganic fertilizer + 25% RDN through FYM + 25% RDN through vermicompost	38100	59847	21747	1.57
T <sub>7</sub> : 75% RDN through inorganic fertilizer + 25% RDN through FYM+ seed treatment with PSB + <i>Azospirillum</i>	35500	73047	37547	2.06
T <sub>8</sub> : 75% RDN through inorganic fertilizer + 25% RDN through vermicompost + seed treatment with PSB + <i>Azospirillum</i>	35600	73599	37999	2.07
T <sub>9</sub> : 75% RDN through inorganic fertilizer + seed treatment with PSB + <i>Azospirillum</i>	34900	47718	12818	1.37
S.E. ±	-	555	555	-
C.D. at 5 %	-	1671	1671	-
<b>G. Mean</b>	<b>36522</b>	<b>61237</b>	<b>24715</b>	<b>1.68</b>

## REFERENCES

- Abdelmuniem Yousif Elamin and K. Madhavi (2012). Influence of integrated nutrient management on growth and yield parameters of kharif Sorghum. *American Journal of scientific and Industrial Research*, 6 (5): 90-96.
- Afzal M., Ahamad A and AU.H. Ahamad (2011). Effect of nitrogen on growth and yield of Sorghum forage under three cutting system. *Cercetari Agronomice in Moldova*; 15. (4) 152-53.
- Gawai P.P. and Pawar V.S. (2003). Integrated Nutrient Management in Sorghum chickpea cropping sequence under irrigated conditions. *Indian Journal of Agronomy* 51 (1): 17-20.
- Kachapur M.D, Roodgi L.I., Chittapur B.M, Basavaraj B. (2001). Influence of vermicompost on the productivity of kharif sorghum. *Karnataka Journal of Agricultural sciences*: 14 (3): 779-80.
- Kalibhavi C.M., Kachapur M.D. and Patil R.H. (1996). Performance of Rabi sorghum under Integrated Nutrient management system. *Indian Journal Dryland agri. Res & Dev.* 16 (1): 45-50.
- Nawale S.S., Pawar A.D., Lambade B.M., and Ugale N.S. (2007). Yield Maximization of chick pea through INM Applied to sorghum-Chickpea Cropping Sequence under Irrigated Condition. *Legume Res.* 32 (4): 282-85.
- Patil H.M., Tuwar S.S. and Wani A.G. (2002). Integrated nutrient management in sorghum-chickpea cropping sequence under irrigated conditions. *Inter. J. Agric. Sci. Vol.* 4 (1): 220-24.

UNDER PEER REVIEW