

## ASSESSMENT OF THE EFFECT OF GINGER FLOUR ADDITION ON THE CHEMICAL AND STORAGE STABILITY OF SOYMILK

Comment [S1]: 'ginger powder' will be effective

### Abstract

This study investigated the effect of ginger flour addition on the storage stability of soymilk. Fresh ginger rhizome was processed to flour by oven drying (OD), sun drying (SD) and ambient drying (AD). The ginger flour were incorporated into processed soymilk at 1 g and 2 g levels, respectively to obtain ginger spiced soymilk, which was stored for 21 days. The chemical composition and microbiological load of the soymilk were determined. The pH ranged from 6.6 to 4.7. The values decreased with storage time. Total titratable acidity (TTA) ranged from 0.0015 to 0.0698 % and increased with storage time. Total soluble solids (TSS) decreased from 26.60 to 3.99 ° Brix as storage time increased. Total solids (TS) decreased from 7.76 to 0.31 % during storage. The total viable counts of the control sample increased from  $4.0 \times 10^3$  to  $71 \times 10^3$  cfu/mL while those of the treated samples increased from 0 to  $33 \times 10^3$  cfu/mL, Coliform was not detected. Yeast and mould growth did not occur from day 0 to day 7 but increased from  $2.0 \times 10^3$  to  $5.0 \times 10^3$  cfu/mL on day 14 and 21, respectively. The highest microbial load was obtained in the unspiced soymilk sample. Two grams of ginger flour preserved the soymilk better than one gram. It is concluded that the incorporation of ginger flour affected the chemical composition and microbial load of soymilk during storage at refrigeration conditions.

Comment [S2]: Language can be modified.

Keywords: Soymilk, ginger flour, chemical composition, storage stability, microbial load.

Comment [S3]: Each key word should start with capital letter.

### Introduction

Soymilk is an aqueous, white, creamy extract produced from soybeans (*Glycine max*) [1]. It is a highly nutritious food drink which contains proteins, fats, carbohydrates, vitamins and minerals [2]. The oldest evidence of soya milk production is from China where a kitchen scene proving use of soya milk is incised on a stone slab dated around AD 25–220 [3]. Soymilk quality depends on the change observed on its physiochemical, microbiological and sensory characteristics [4]. In addition, its shelf-life is limited due to the deterioration of its properties during storage [5]. In order to improve the soymilk quality, different flavouring ingredients have been added during its manufacturing [6]. Other than the whole seed, many processed soy products are available in the market. They include soya milk, soya flour, soya

curd and *tofu* (soya paneer) [7]. Soybean is the main raw material used in the preparation of soymilk.

Soybean (*Glycine max*) is a non-native and non-staple crop in Sub-Saharan Africa (SSA) with potential to be a commercial crop owing to its wide range of uses as food, feed, and industrial raw material [8]. In 1908, soybean was introduced to Nigeria and was cultivated as an export crop in a small area in Benue state, where the introduced variety ‘Malayan’ was adopted [9]. The crop is usually in small holdings in mixed cropping with sorghum or maize or as an intercrop in citrus orchards. Zaire has a long history of soybean production by indigenous farmers [3]. Soybeans were introduced and promoted first by missionaries before the nation gained independence, and are considered a medicinal food to prevent and cure the wasting effects of malnutrition in Zaire [10]. In Sub-Saharan Africa (SSA), South Africa is the largest soybean producer in 2016, followed by Nigeria, Zambia, and Uganda [8]. Since the value of soybean as a high-protein food source has been recognized, the utilization and consumption of soybean-based foods are becoming popular in SSA [9]. Food products such as *nshima*, also called *bidia*, *dawadawa*, *mahewu*, *soy-ogi* (fortifying maize with soybean), *soy kebab* (spicy *tofu*), biscuits, soy flour, soy yogurt, and soymilk have been accepted by local people in many SSA countries [8]. In SSA, soymilk is the daily meal of malnourished children and HIV/AIDS-infected patients in public schools and hospitals [3]. Nigeria offers the best example in SSA for promoting utilization of soy-based food products in rural and urban areas. The high utilization and consumption of soy-based food products in Nigeria is attributed to development of home-level and small-scale processing technologies for soy-based foods, training of technical and extension staff, training of local farmers and village leaders, dissemination of food processing technologies, and support from the public and private sectors [8].

Comment [S4]: Should be in full form

Most flavourants used in the industrial production of soymilk are synthetic [5]. The use of natural flavourants such as fruits or vegetables will be more beneficial to the consumers' health because they are a good source of minerals, fibres, vitamins, and bioactive compounds [8]. Amongst many other spices, ginger, a root vegetable is one of the mostly used culinary spices [11]. The need for use of inexpensive local resources in the preservation of staple foods to limit post-harvest losses is of high importance in our society today. As a local beverage produced by traditional methods, soymilk offer only a short lifespan [6]. The milk is underutilized due to its short shelf-life, which hinders widespread consumption of the beverage due to the deteriorating effects of some microorganisms on the milk [5]. There have been many attempts to industrialize the locally prepared soymilk, but the inability to preserve the milk for a long time has been a major problem [8]. Some authors have evaluated the preservative effect of ginger on food products [12, 13, 14, 15, 16] but highlighted the need for intensification in order to obtain improved flavour, aroma, as well as to extend the shelf life. Addition of ginger powder to soymilk could improve the antimicrobial and antioxidant (bioactive) properties of soymilk and thus rendering it more functional with improved shelf life [17]. The aim of the present study therefore is to assess the effects of ginger flour on the chemical and storage stability of soymilk.

Comment [S5]: Should use accurate word

Comment [S6]: (.) should be used

## Materials and methods

Comment [S7]: Study periodis missin in the methodology

### Sample Collection

The fresh roots (rhizomes) of ginger (*Zingiber officinale*) and soybeans (*Glycine max*) were purchased from Wurukum market, Makurdi, Benue State and taken to the Food Technology Laboratory of the Center for Food Technology and Research (CEFTER), Benue State University for processing and analysis.

Comment [S8]: Spelling must be checked

### Preparation of ginger flour

Comment [S9]: Should start with capital letter

The ginger was washed several times with tap water (potable water). It was peeled, rewashed with tap water, sliced into fillets of 2- 3 diameters thickness.

- i. Oven dried at 50°C for 12 h.
- ii. Sun dried at 40°C to 42 °C for 3 days, with relative humidity at 75 % to 75.3 % using a thermo-hydrometer.
- iii. Ambient dried at 30.35°C to 32.2°C for 5 days, with relative humidity at 59.5 %, 60 % and 63.5 %.

The dried ginger was blended into powder in three proportions using a kitchen blender (model: Binatone BLG-452), and sieved through a sieve of pore size 0.5 mm [18]. The sieved ginger powder was stored in well-sealed plastic containers at room temperature in a closed cupboard to avoid sun light.

Comment [S10]: the information drawn from ref 18 should be clear.

### Preparation of ginger spiced soymilk

Comment [S11]: Should start with caopital letter such as Ginger Spiced Soymilk.

Soybean was sorted and cleaned to remove stones, damaged and deformed seeds. Then soybean was further weighed using an electronic weighing balance (capacity 2,500 g), washed and soaked in water (500 g in 1 L) overnight for 8-12 h. It was then rinsed and blanched in 1.25% NaHCO<sub>3</sub> (baking soda) for 30 min. The rehydrated soybean was then washed, manually dehulled and rinsed. The soybean seeds were wet milled (in the ratio of 3:1 water to beans on a weight basis) using a kitchen blender (model: SB736). The milk obtained was sieved (0.5 mm) and pasteurized at the temperature 70°C for 15 seconds, cooled (18°C) [18]. The soymilk and ginger flour were homogenised to ensure uniformity throughout the product. The mix was subsequently packaged and stored at refrigeration temperature (4 to 6°C).

Comment [S12]: the information drawn from ref 18 should be clea

Comment [S13]: Alignment should be corrected

### Formulation of ginger spiced soymilk

The obtained milk was then formulated by adding ginger powder. The ginger spiced soymilk was used to produce 7 samples in all (A - G). Sample A was plain soymilk which served as the control and samples B - G were prepared by addition of 1 g and 2 g ginger powder to every 500 mL of soymilk as shown in Table 1.

**Table 1 Sample formulation for ginger spiced soymilk**

Samples	Soymilk (mL)	flour (g)
A (control)	500	0
OB	500	1
OC	500	2
SD	500	1
SE	500	2
AF	500	1
AG	500	2

A= plain soymilk, OB=500 mL soymilk and 1 g oven dried ginger flour, OC = 500 mL soymilk and 2 g oven dried ginger flour, SD = 500 mL soymilk and 1 g sundried ginger flour, SE = 500 mL soymilk and 2 g sundried ginger flour, AF= 500 mL soymilk and 1 g ambient dried ginger flour, AG = 500 mL soymilk and 2 g ambient dried ginger flour.

### Chemical properties of ginger spiced soymilk

#### Determination of pH

1. The pH was determined by the use of a glass electrode pH meter (Model: OAKLON, SN: 1564140) at ambient temperature according to the method of AOAC [19]. Ten millilitres of sample was measured in to a beaker. The pH was then standardized using a buffer solution. The electrode of the pH meter was rinsed with distilled water and then inserted into the beaker containing the sample. The pH reading was recorded in triplicates and pH electrode rinsed with distilled water.

#### **Determination of total titratable acidity**

Comment [S14]: Serial should be denoted as 2.

Titratable acidity was determined according to the method of AOAC [19]. Blended portion of the samples (2 mL) was weighed and transferred into 50 mL centrifuge tube. About 10 mL of distilled water was added. One millilitre aliquot of the solution was taken into another 50 mL centrifuge tube and 10 mL of distilled water added to dilute the sample. The diluent (10 mL) was titrated against 0.1N NaOH solution using phenolphthalein (2 drops) indicator and percentage titratable acidity was calculated:

$$\text{TTA} = \frac{\text{mL of NaOH used} \times 0.1\text{M NaOH} \times \text{milliequivalent} \times 100}{\text{Volume of sample}} \quad (1)$$

#### **Determination of total solids content**

Total solids content of ginger spiced soymilk samples was measured by a method described by Matin *et al.* [20] and the results expressed in terms of percentage. Two millilitres of ginger spiced soymilk sample was measured into a beaker, warmed slowly to 35 - 40 °C on a water bath with careful mixing to incorporate any cream adhering to the sample. The sample was then cooled quickly to room temperature. A dish was heated with its lid alongside in the drying oven at least 1 h. The lid was then placed on the dish and immediately transferred to a desiccator. It was allowed to cool to room temperature (at least 30 min) and weighed to 0.1 mg. A 5 mL of prepared sample was added and lid placed on the dish and weighed again. The dish was further placed without the lid on the vigorously boiling water bath in such a way that the bottom of the dish was directly heated by the steam. There was continuous heating till water was removed. The dish was removed from the water bath, with the underside wiped and placed it in the oven alongside the lid for 2 h. The lid was placed on the dish and transferred to the desiccator. The dish was allowed to cool and weighed to the nearest 0.1 mg. The dish was heated in the oven for 1 h. It was allowed to cool and weighed again. The

operation was repeated until the difference in the two consecutive weighing did not exceed 1 mg. The lowest mass was recorded.

#### **Determination of total Soluble Solids content**

The total soluble solids in the ginger spiced soymilk samples was determined by using a digital refractometer (Model: ATAGO RX-5000) and expressed in terms of °Brix [21].

#### **Determination of total viable count**

All media were prepared by weighing appropriate amount of the flour according to manufacturer's specification [22]. Following sterilization of media by autoclaving at 121 °C for 15 min, 1 mL of ginger spiced soymilk samples was measured with the aid of a sterile syringe. After vigorous shaking, a plastic rack was arranged with sterile test tubes containing 9 mL of sterile distilled water. A tenfold serial dilution [23], was carried out by dropping 1 mL of the sample into the first test tube labelled  $10^{-1}$ . This was mixed properly. 1 mL was again taken from the  $10^{-1}$  dilution tube and transferred into the next test tube labelled  $10^{-2}$ . The dilutions continued to dilution  $10^{-3}$ . Each test tube was vigorously shaken before each transfer. All plate samples were duplicated. Colonies were then counted and recorded after incubation [24].

#### **Determination of coliform count**

Eosin methylene blue agar (EMB) was inoculated with serial dilutions of the sample by the pour plate method and incubated at 37 °C for 24-48 h. The different representative colonies from Eosin methylene blue agar (EMB) plates were then counted and recorded after incubation [25].

#### **Determination of fungi count**

Fungi counts were determined on Potato dextrose agar (PDA). The PDA media was modified with Chloramphenicol capsule to inhibit bacterial contamination. One millilitre of the diluted

samples was measured with the use of a sterile syringe and poured into an empty sterile petri dish, followed by pouring PDA into the prepared plates. The plates were rotated clockwise for easy mix up of the sample and the media and incubated for 3-5 days at 37 °C. Colonies were then counted and recorded after incubation period [26].

### **2.7.3 Counting of the Colonies**

After incubation of all the plates, counts of the number of colonies in each plate was done with a hand tally counter [22]. A mean of the count was obtained and multiplied with the appropriate diluting factor. The mean count was calculated as shown below:

Mean = total viable count/ number of plates

The estimation of viable number of microorganisms (total viable counts) in each sample was made in colony forming units (cfu)

TVC = 1 Weight of sample \*N\*D

Where N = average number of colonies

D = dilution factor

### **Storage Stability Study**

Storage stability was determined within 21 days (3 weeks) of storage at refrigeration conditions (4 – 6 °C). Data were collected at 0, 7, 14 and 21 days interval on pH, total titratable acidity, total dissolved solids and total soluble solids.

### **Statistical Analysis**

Data obtained were subjected to the Analysis of variance (ANOVA) using Statistical Package for Social Science (SPSS) version 21. The means obtained were separated using Tukey HSD test. Differences were considered at 95 % ( $P \leq 0.05$ ) confidence interval.

### **Results and discussion**

Comment [S15]: Should start with capital letter

### pH of ginger spiced soymilk during storage

Comment [S16]: Sequence number missing

There was a significant decrease in pH of ginger spiced soymilk during storage at refrigeration conditions between unspiced and spiced soymilk samples. The initial pH of all the samples was near to neutrality ranging from 6.6 to 6.9. Similar results were reported by Arekemase *et al.* [27] on the assessment of the effectiveness of ginger (*Zingiber officinale*), clove (*Syzygium aromaticum*) and Sodium Benzoate on the shelf life of soymilk and Odom *et al.* [28] on comparative studies of ginger (*Zingiber officinale*) and black pepper (*Piper guinenses*). The pH of the entire samples followed the same decreasing pattern as storage progressed. During storage, and in refrigerated conditions, activity of organic acids is not completely stopped and there is acidification by organic acid leading to a drop of pH. The reduction in pH might also be due to the fact that the initial pH of the soymilk samples favoured the growth of bacteria and then led to subsequent release of metabolic products into the medium. Similar results were reported by Afroz *et al.* [29] on the preparation of soymilk using different method. This results were also in line with values reported by Amadou *et al.* [18] on the physicochemical and sensory properties of ginger spiced yoghurt. In a study by Odu *et al.* [30], it was reported that the reduction in pH during fermentation might be as a result of acidification by acetic acid fermentation of breadfruit which reduces the chance of microbial spoilage. At the end of storage, the unspiced soymilk had the lowest ( $P<0.05$ ) pH value while soymilk sample spiced with 2 g ginger flour had the highest. The low values ( $P<0.05$ ) observed for the spiced soymilk samples compared to the unspiced soymilk sample at the end of storage could be as a result of low level of fermentation due to the antimicrobial activity of the ginger, thus the ginger could contribute to the preservation of soymilk. The significant ( $P<0.05$ ) difference between the initial and the final values of soymilk samples pH could also be related to the continuous fermentation at refrigerated conditions.

Comment [S17]: Sentence should be grammatically correct

**Table 2:** pH of Ginger Spiced Soymilk during Storage

Samples	Day 0	Day 7	Day 14	Day 21
A	6.9 <sup>a</sup> ±0.05	5.5 <sup>cd</sup> ±0.10	4.2 <sup>d</sup> ±0.00	3.7 <sup>e</sup> ±0.05
OB	6.8 <sup>ab</sup> ±0.05	5.5 <sup>cd</sup> ±0.00	5.3 <sup>ab</sup> ±0.05	4.7 <sup>cd</sup> ±0.05
OC	6.7 <sup>bc</sup> ±0.05	5.7 <sup>ab</sup> ±0.00	5.4 <sup>b</sup> ±0.05	4.9 <sup>a</sup> ±0.00
SD	6.8 <sup>b</sup> ±0.05	5.4 <sup>d</sup> ±0.05	5.2 <sup>c</sup> ±0.05	4.7 <sup>d</sup> ±0.00
SE	6.7 <sup>bc</sup> ±0.05	5.6 <sup>c</sup> ±0.00	5.3 <sup>ab</sup> ±0.05	4.8 <sup>bc</sup> ±0.00
AF	6.8 <sup>b</sup> ±0.00	5.5 <sup>cd</sup> ±0.00	5.3 <sup>ab</sup> ±0.05	4.8 <sup>ab</sup> ±0.05
AG	6.6 <sup>c</sup> ±0.05	5.8 <sup>a</sup> ±0.00	5.5 <sup>a</sup> ±0.05	4.9 <sup>a</sup> ±0.00
LSD	0.08	0.08	0.07	0.17

Values are means ± SD of 3 replicates. Means within a column with the same superscript were not significantly ( $P>0.05$ ) different.

#### Total Titratable Acidity of ginger spiced soymilk during storage

There was a gradual increase ( $P<0.05$ ) in titratable acidity of all samples during storage. Moreover, the unspiced sample presented the highest increase in titratable acidity ( $P<0.05$ ) especially on day 14 and at the end of storage. For spiced samples, the initial and the final values of titratable acidity were significantly ( $P>0.05$ ) different while those of unspiced sample showed a similar difference ( $P>0.05$ ). The high increase in titratable acidity obtained with the unspiced sample was due to fermentation (activity of bacteria) which is not completely stopped during refrigeration thus leading to the production of organic acid. The antimicrobial activity of ginger could be responsible for the non-significant variation of titratable acidity in the spiced soymilk samples during storage. The gradual increase in titratable acidity during storage was also observed in Sudanese yoghurt [18]. The titratable acidity values obtained during storage at refrigeration temperature is comparable to the values

**Comment [S18]:** Sequence number should be added with previous number

obtained by [2], who reported mean values of titratable acidity in the range of 0.87 to 1.13 %. This was different from the values of 2.5 to 2.07 % reported by [31] on soymilk stored at refrigeration temperature. It was also observed that titratable acidity increased with decrease in pH of the samples. Similar increasing trend of titratable acidity with a decrease in pH was observed in soymilk samples preserved with local spices during storage by [25]. These values were in line with the values reported for soy-corn milk (0.063 and 0.057 % in white and yellow maize, respectively) by [32]. The high titratable acidity of unspiced sample compared to spiced samples could indicate a positive effect of ginger on the preservation of soymilk.

**Table 3:** Total Titratable Acid (%) of Ginger Spiced Soymilk during Storage

Samples	Day 0	Day 7	Day 14	Day 21
<b>A</b>	0.002 <sup>a</sup> ±0.00	0.006 <sup>a</sup> ±0.00	0.062 <sup>a</sup> ±0.00	0.069 <sup>a</sup> ±0.00
<b>OB</b>	0.002 <sup>a</sup> ±0.00	0.005 <sup>b</sup> ±0.00	0.055 <sup>c</sup> ±0.00	0.063 <sup>f</sup> ±0.00
<b>OC</b>	0.002 <sup>a</sup> ±0.00	0.005 <sup>b</sup> ±0.00	0.053 <sup>d</sup> ±0.00	0.066 <sup>e</sup> ±0.00
<b>SD</b>	0.002 <sup>a</sup> ±0.00	0.006 <sup>a</sup> ±0.00	0.061 <sup>a</sup> ±0.00	0.067 <sup>c</sup> ±0.00
<b>SE</b>	0.002 <sup>a</sup> ±0.00	0.005 <sup>b</sup> ±0.00	0.057 <sup>b</sup> ±0.00	0.068 <sup>b</sup> ±0.00
<b>AF</b>	0.002 <sup>a</sup> ±0.00	0.006 <sup>a</sup> ±0.00	0.057 <sup>b</sup> ±0.00	0.067 <sup>d</sup> ±0.00
<b>AG</b>	0.002 <sup>a</sup> ±0.00	0.005 <sup>b</sup> ±0.00	0.051 <sup>e</sup> ±0.00	0.061 <sup>g</sup> ±0.00
<b>LSD</b>	0.002	0.003	0.007	0.003

Values are means ± SD of 3 replicates. Means within a column with the same superscript were not significantly ( $P>0.05$ ) different.

### 3.3 Total soluble solids of ginger spiced soymilk during storage

Statistically TSS of samples showed significant ( $P < 0.05$ ) difference with respect to increased number of storage days and change in treatment. According to [33] about half of the solids in soymilk consist of soybean protein. As a common problem with soymilk is lack of stability sediment precipitation of proteins and other added solid particles such as minerals or flavours it might be the reason for lowered value of TSS during storage. Similar results were reported by khodke *et al* [33] on a study on the storage of sterilized soymilk.

Comment [S19]: Should be used in proper way

**Table 4:** Total Soluble Solids (<sup>o</sup>brix) of Ginger Spiced Soymilk during Storage

Samples	Day 0	Day 7	Day 14	Day 21
A	21.19 <sup>c</sup> ±0.00	13.32 <sup>e</sup> ±0.02	6.66 <sup>e</sup> ±0.00	3.99 <sup>e</sup> ±0.01
OB	23.14 <sup>b</sup> ±0.12	13.35 <sup>de</sup> ±0.05	6.67 <sup>e</sup> ±0.00	4.01 <sup>e</sup> ±0.00
OC	23.02 <sup>b</sup> ±0.00	14.86 <sup>c</sup> ±0.04	7.43 <sup>c</sup> ±0.00	4.27 <sup>b</sup> ±0.01
SD	22.94 <sup>b</sup> ±0.00	13.52 <sup>d</sup> ±0.02	6.77 <sup>d</sup> ±0.01	4.05 <sup>d</sup> ±0.00
SE	23.26 <sup>b</sup> ±0.00	16.47 <sup>b</sup> ±0.09	8.24 <sup>b</sup> ±0.01	4.11 <sup>c</sup> ±0.01
AF	26.53 <sup>a</sup> ±1.00	13.35 <sup>de</sup> ±0.13	6.66 <sup>e</sup> ±0.00	4.01 <sup>e</sup> ±0.00
AG	26.60 <sup>a</sup> ±0.00	18.52 <sup>a</sup> ±0.02	9.26 <sup>a</sup> ±0.00	4.64 <sup>a</sup> ±0.01
LSD	0.67	0.12	0.01	0.02

Values are means ± SD of 3 replicates. Means within a column with the same superscript were not significantly ( $P > 0.05$ ) different.

### 3.4 Total solid content (TS) of ginger spiced soymilk during storage

There was a significant ( $p < 0.05$ ) difference in the initial TS of ginger spiced soymilk and the untreated sample. The ginger spiced samples had the highest TS and the least value was recorded by sample A. This could be due to the higher levels of suspended particles in the treated samples. These results were in line with findings of Ajala *et al.*, [32] on physicochemical and sensory qualities of spiced soy-corn milk.

**Table 5:** Total Solids (%) content of Ginger Spiced Soymilk during Storage

Samples	Day 0	Day 7	Day 14	Day 21
A	3.76 <sup>f</sup> ±0.00	1.70 <sup>c</sup> ±0.00	1.50 <sup>e</sup> ±0.01	0.31 <sup>e</sup> ±0.01
OB	4.76 <sup>e</sup> ±0.00	2.70 <sup>b</sup> ±0.00	1.51 <sup>e</sup> ±0.01	0.31 <sup>e</sup> ±0.01
OC	4.78 <sup>e</sup> ±0.02	2.76 <sup>b</sup> ±0.00	1.85 <sup>a</sup> ±0.02	0.81 <sup>b</sup> ±0.01
SD	4.81 <sup>d</sup> ±0.00	1.77 <sup>c</sup> ±0.01	1.67 <sup>c</sup> ±0.00	0.83 <sup>b</sup> ±0.00
SE	6.81 <sup>b</sup> ±0.01	2.80 <sup>a</sup> ±0.00	1.86 <sup>a</sup> ±0.01	0.97 <sup>a</sup> ±0.01
AF	5.50 <sup>c</sup> ±0.00	1.76 <sup>c</sup> ±0.00	1.57 <sup>d</sup> ±0.01	0.36 <sup>d</sup> ±0.01
AG	7.76 <sup>a</sup> ±0.00	2.76 <sup>b</sup> ±0.00	1.76 <sup>b</sup> ±0.00	0.53 <sup>c</sup> ±0.01
LSD	0.016	0.008	0.019	0.017

Values are means  $\pm$  SD of 3 replicates. Means within a column with the same superscript were not significantly ( $P > 0.05$ ) different.

### 3.5 Microbiological analysis of ginger spiced soymilk during storage

#### 3.5.1 Total Viable Count (TVC) of ginger spiced soymilk during storage

Microbial growth causes degradation of food quality which leads to visible changes in colour, odour and texture. Changes in colour, flavour, taste, whey separation and acid production

were observed in the preserved ginger spiced soymilk samples at different time intervals. There was a gradual increase in counts as storage progressed from day 0 to day 21 in all samples. Lower viable counts were observed in samples treated with ginger powder as compared to the untreated sample. The lower counts observed in treated samples was attributed to the antimicrobial properties (gingerol, shigoal and gingerdoil) of ginger. Similar results were reported by Arekemase *et al.* [27] on the assessment of the effectiveness of ginger, clove and sodium Benzoate on the shelf life of soymilk. It is also in line with the work of Amadou *et al.* [10] on the effect of ginger extract on the physicochemical and sensory properties of yoghurt. This result were in accordance with the work of Akponah *et al.* [34] who reported on the potency of ginger in extending the shelf life of orange juice. The result dovetails with the findings of Adesokan *et al.* [35], who reported the effectiveness of ethanolic extract of ginger in prolonging the shelf life of West African soft cheese for up to 3 days. This is also in agreement with the work of Obi *et al.* [36], who reported that the prolonged shelf life of ginger treated zobo drink for 6 days at room temperature was due to it antimicrobial property.

**Table 6: Total Viable Bacterial Count (TVBC) ( $\times 10^3$  cfu/mL)**

Samples	Day 0	Day 7	Day 14	Day 21
<b>A</b>	4.0 <sup>a</sup> ±0.00	17 <sup>a</sup> ±0.10	30 <sup>a</sup> ±0.10	71 <sup>a</sup> ±2.00
<b>OB</b>	3.0 <sup>b</sup> ±0.05	4.0 <sup>b</sup> ±0.00	23 <sup>b</sup> ±0.50	33 <sup>b</sup> ±0.30
<b>OC</b>	2.0 <sup>b</sup> ±0.00	4.0 <sup>b</sup> ±0.05	15 <sup>c</sup> ±0.10	22 <sup>bc</sup> ±0.20
<b>SD</b>	3.0 <sup>b</sup> ±0.05	5.0 <sup>b</sup> ±0.05	10 <sup>d</sup> ±0.45	22 <sup>bc</sup> ±1.15
<b>SE</b>	2.0 <sup>b</sup> ±0.05	3.0 <sup>b</sup> ±0.00	9.0 <sup>d</sup> ±0.00	11 <sup>c</sup> ±0.10
<b>AF</b>	-	-	6.0 <sup>e</sup> ±0.10	14 <sup>bc</sup> ±0.36
<b>AG</b>	-	-	4.0 <sup>e</sup> ±0.00	10 <sup>c</sup> ±0.25

Values are means  $\pm$  SD of 3 replicates. Mean values within a column with the same superscript were not significantly ( $P>0.05$ ) different.

### 3.5.2 Total Coliform Counts (TCC) of ginger spiced soymilk during storage

There was no coliform detected in all samples throughout the storage period of 21 days. These results were in accordance with the SON standard which stipulates no coliform count is allowed in soymilk. The inhibition of coliform in all samples could be due to proper method of production and preservation of samples. This could also be attributed to the antimicrobial properties of ginger in spiced samples. Arekemase *et al.* [27] reported that, phytochemical screening of ginger and clove extracts showed the presence of tannins, phenol, alkaloid, flavonoid, terpenoids, steroids, glycosides, coumarin and saponins. The antimicrobial activities of ginger might be due to these phytochemicals. This finding agrees with those of other workers[36, 28]. The findings of Odom *et al.* [28] suggest that the preservative effect of ginger and black pepper on soymilk was as a result of their essential oil fraction which was inhibitory to microbial growth. Obi *et al.* [36] reported that the antimicrobial activity of spice in zobo drink was as a result of their volatile oil. This study confirms that the use of ginger flour was effective in the inhibition of coliforms in the product.

**Table 7:** Total Coliform Count (TCC) ( $\times 10^3$ cfu/mL)

Samples	Day 0	Day 7	Day 14	Day 21
A	Nil	Nil	Nil	Nil
OB	Nil	Nil	Nil	Nil
OC	Nil	Nil	Nil	Nil
SD	Nil	Nil	Nil	Nil
SE	Nil	Nil	Nil	Nil

<b>AF</b>	Nil	Nil	Nil	Nil
<b>AG</b>	Nil	Nil	Nil	Nil

Values are means of 3 replicates. Means within a column with the same superscript were not significantly ( $P>0.05$ ) different.

### 3.5.3 Total Yeast and Mould counts of ginger spiced soymilk during storage

According to the standards organisation of Nigeria (SON), a maximum of 5 cfu/mL yeast and mould count is allowed in soymilk. From the results, there was no yeast and mould count detected from 0 to 7 days. During day 14 and day 21 of storage there were fungi counts (1cfu/mL) grew in each of the treated samples, while the untreated sample had the highest count. All the treatments showed a higher suppressive action against fungal, than the bacterial isolates. The findings agreed with those of other workers including [37, 38]. Similar findings were reported by Akponah *et al.* [34] that all extracts of ginger, garlic and rosemary exhibited more antimicrobial action against fungal isolates than the bacterial isolates. [27] research work point to the fact that the antifungal effects might be due to monoterpene, which destroyed the integrity of fungi membrane. [31] reported that the inability of the ginger, clove and a combination of ginger-clove extract to completely inhibit fungal and bacterial growth might be due to the low concentration of the extracts used, as higher concentrations of plant extracts were required before antimicrobial properties were observed. In addition, it was observed that many spices have the ability to inhibit microbial growth in food than in culture media [27]. However, large amounts of the spice extracts required for complete microbial inhibition would affect the organoleptic properties of the milk products [31].

**Comment [S20]:** Reference should be used in proper way

**Comment [S21]:** Reference should be used in proper way

**Table 8:** Total Yeast and Mould Counts ( $\times 10^3$  cfu/mL)

Samples	Day 0	Day 7	Day 14	Day 21
A	-	-	2.0 <sup>a</sup> ±0.50	5.0 <sup>a</sup> ±0.57
OB	-	-	1.0 <sup>b</sup> ±0.00	1.0 <sup>b</sup> ±0.00
OC	-	-	1.0 <sup>b</sup> ±0.00	1.0 <sup>b</sup> ±0.00
SD	-	-	1.0 <sup>b</sup> ±0.00	1.0 <sup>b</sup> ±0.00
SE	-	-	1.0 <sup>b</sup> ±0.00	1.0 <sup>b</sup> ±0.00
AF	-	-	1.0 <sup>b</sup> ±0.00	1.0 <sup>b</sup> ±0.00
AG	-	-	1.0 <sup>b</sup> ±0.00	1.0 <sup>b</sup> ±0.00

Values are means  $\pm$  SD of 3 replicates. Means within a column with the same superscript were not significantly ( $P > 0.05$ ) different.

### Conclusion

The study established that the ginger spiced soymilk had a longer shelf stability compared to the untreated (control).

**Comment [S22]:** Conclusion should be more meaningful.

### References

- [1] Kohli, D., Kumar, S., Upadhyay, S. and Mishra, R. (2017). Preservation and processing of soymilk: A review. *International Journal of food science and Nutrition*, 2(6): 66–70.
- [2] Borode, O. F. (2017). The Effect of Water and Ethanol Extracts of Ginger and Garlic on the Nutritional Quality and Physico-Chemical Properties of Stored Soymilk. *International Journal of Food Science and Biotechnology*, 2( 2): 43–50.

- [3] Khojely, D. M., Ibrahim, S. E., Sapey, E. and Han, T. (2018). History, current status, and prospects of soybean production and research in sub-Saharan Africa. *Journal of Crop Production*, 6(3):226–235.
- [4] Momoh, J. E., Udobi, C. E. and Orukotan, A. A. (2011). Improving the Microbial Keeping Quality of Home Made Soymilk Using a Combination of Preservatives, Pasteurization and Refrigeration. *British Journal of Dairy Sciences*, 2(01): 1–4.
- [5] Shakeel, A., Saeed, M., Khuram, H., Aslam, W. and Naheed, N. (2015). Extraction of Soya Milk from Different Varieties of Soya Beans and Comparative Study for better Nutrition with Buffalo Milk. *Journal for Global Innovation Agriculture and Social Science*, 3(4): 146 - 151.
- [6] Oyeniyi, A. O., Aworh, O. C. and Olaniyan, J. O. (2014). Effect of Flavourings on Quality and Consumer Acceptability of Soy-Yoghurt. *Journal of Environmental Science, Toxicology and Food Technology*, 8(1): 38–44.
- [7] Nadifah, F. and Sari, R. M., F. (2016). The Effect of Ginger (*Zingiber officinale*) and Green Tea ( *Camellia sinensis* ) Against Bacteria Growth on Soymilk. *Journal of American institute of physics*, 2(1): 1- 4.
- [8] Akakpo, A. Y., Somda, M. K., Kabore, D., Mihin, H. B. and Ouattara, A. S. (2019). Natural tropical and chemical preservatives effects on shelf-life and hedonic acceptability of tiger-nut milk and soymilk under storage conditions : A review, *International Journal of Food Science and Nutrition*, 4(6): 80 - 87.
- [9] Erhirhie, E. O. and Moke, G. E. (2016). A Review of the Ethnomedicinal , Chemical and Pharmacological Properties of *Xylopi aethiopica*. *American Journal of Pharmtech Research* 4(6): 1-37.
- [10] Ghosh, A. K, Banerjee, S., Mullick, H. I. and Banerjee, J. (2011). *Zingiber officinale*: A natural gold. *International Journal of Pharmacy and Biological Science*, 2(1): 283–294
- [11] Amadou, N. M., Richard, E. A., Roger, K. J., Waingeh, N. C., Ateh, K. D., Mbiydzengh, A. F., Che, N. S. and Yunenyui, M. P. (2018). Effect Of Ginger Extract On The Physicochemical And Sensory Original Research Article Effect Of Ginger

Extract On The Physicochemical And Sensory Properties Of Yoghurt. *International Journal of Development Research*, 8(05):1–11.

- [12] Amadou, N. M., Richard, E. A., Waingeh, N. C., Hélène, I., Ndombow, Y. K. and Jules-roger, K. (2019). Physicochemical and Sensory Properties of Ginger Spiced Yoghurt Physicochemical and Sensory Properties of Ginger Spiced Yoghurt. *Journal of Nutritional Therapeutics*, 6(3):1–8.
- [13] Dauda, A. O. and Abiodun, O. A. (2017). Comparative study of the effect of dry and wet ginger (*Zingiber officinale Roscoe*) spice on the proximate and microbial safety of soybean beverage. *Journal for Food Science and Technology*, 9(2): 130–135.
- [14] Izah, S. C., Kigigha, L. T., Aseibai, E. R. and Okowa, I. P. (2016). Advances in Preservatives and Condiments used in Zobo (a food-drink) production. *Journal of Biotechnoly and Research*, 2(3): 104–119.
- [15] Odom, T. C., Udensi, E. A., Dike, C. O., Ogbuji, C. A., Kanu, A. M. and Aji, R. U. (2012). Comparative studies of ginger (*Zingiber officinale*) and black pepper (*Piper guinenses*) extracts at different concentrations on the microbial quality of soymilk and kunuzaki. *African Journal of Biotechnology*, 11(70): 13494–13497.
- [16] Aisha, I. Ephraim, B. E. and Jamila, W. M. (2017). Potential of Soyabeans Seasoning with Garlic and Ginger Spices for Sustainable Health in Bauchi State Nigeria. *International Journal of Development Strategies in Humanities, Management and Social Sciences*, 7(1): 22–39, 2017.
- [17] Sharif, Z. I. M., Mustapha, F. A., Jai, J., Yusof, N. M. and Zaki, N. A. M. (2017). Review on methods for preservation and natural preservatives for extending the food longevity. *Journal of Chemical Engineering and Research*, 19(01): 145–153.
- [18] Zamal, S. S., Uddin, M. B. and Huda, M. S. (2011). Effect of Preservatives on the Shelf-Life of Soymilk, *Eco-friendly Agricultural Journal*, 4 (01): 520–523.
- [19] AOAC. (2010). Official methods of analysis, Association of Official Analytical Chemists, 15th Edition, *Washington, D.C., USA*, 807-928.
- [20] Matin, A., Banik, T., Badsha, R., Hossain, A., Haque, M. M. and Ahmad, M. (2018).

Microbiological quality analysis of yoghurt in some selected areas of Bangladesh. *International Journal of Natural and Social Sciences*, 5(4): 82- 86.

- [21] Pratap, D, Maurya, V. K., Kumar, N., Singh, R. and Upadhyay, A. (2017). Studies on Physico-chemical and Organoleptic Properties of Soymilk Blended Dahi (Curd) with Toned Milk (Cattle Milk). *Current Nutrition and Food Science Journal*, 13(3): 1- 7.
- [22] Stanley, M. C., Ifeanyi, O. E., Ifediora, A. C. and Uzoma, U. G. (2014). Isolation And Identification Of Microorganisms Involved In The Spoilage Of Soymilk. *Journal of Pharmacy and Biological Science*, 9(5): 28–36.
- [23] Edet, N. and Peter, S. (2017). Microbiological and Biochemical Analysis of Soymilk Produced and Sold within Calabar Metropolis. *International journal of Microbiology*, 1(1) 1-7.
- [24] Varghese, N. and Joy, P. P. (2016). Microbiology laboratory manual, 1<sup>st</sup> Edition, *Pineapple Research Station*, 1-76.
- [25] Liamngee, K., Terna, T. P., Bem, A. A., Orpin, J. B, Mzungu, I. and Obaje, M. (2013). Microbial Analysis of Soyabean Milk Sold In Makurdi Metropolis. *Journal for Environmental Science, Toxicological and Food Technology*, 3(3): 97–104.
- [26] Datti, Y., Salihu, I., Nura, T. and Imrana, B. (2019). Proximate Composition and the Mineral Contents of Soya Beans (*Glycine max*) Available in Kano State, Nigeria. *Chemistry Search Journal*, 10(2): 62–65.
- [27] Arekemase, M. O. and Babashola, D. R. (2019). Assessment of the Effectiveness of Ginger (*Zingiber officinale*), Clove (*Syzygium aromaticum*) and Sodium Benzoate on the Shelf Life of Soymilk. *Journal of Natural Biological Science*, 11(4):400–409.
- [28] Odom, T. C., Udensi, E. A., Dike, C. O., Ogbuji, C. A., Kanu, A. M. and Aji, R. U. (2012). Comparative studies of ginger (*Zingiber officinale*) and black pepper (*Piper guinenses*) extracts at different concentrations on the microbial quality of soymilk and kunuzaki. *African Journal of Biotechnology*, 11(70): 13494–13497.
- [29] Afroz, M. F., Anjum, W., Kobir, A., Hossain, K. and Sayed, A.(2016). Preparation of Soymilk Using Different Methods. *International Journal of Food Science*, 4(1): 11–

17.

- [30] Odu, N. N. and Egbo, N.N. (2012). Assessment of the Effect of Different Preservatives on the Keeping Quality of Soymilk Stored at Different Temperatures. *Journal of Nature and Sciences*, 10(9): 1–9.
- [31] Kabiru, S. A., Makun, Y. A., Saidu, H. A., Muhammad, A. N., Nuntah, L. H. and Amoo, L. C. (2012) Soymilk Preservation Using Extracts Of Cloves (*Syzygium aromaticum myrtaceae*) And Guinea-Pepper (*Xylopia aethiopica annonaceae*). *Journal of Pharmacy and Biological Sciences*, 3(5):44–50.
- [32] Khodke, S., Shinde, K. and Ashta, T. (2014). A study on the storage of sterilized soymilk. *International Journal of Food Science*, 2(1): 1- 12.
- [33] Kaushik, P. and Goyal, P. (2015). Evaluation of Various Crude Extracts of *Zingiber officinale* Rhizome for Potential Antibacterial Activity, *Journal of Advanced Microbiology*. 1(1): 7–12.
- [34] Akponah, F. E., Ibokoro, E., Ubogu, I.O. and Ejukonemu, M. (2013). Effects Of Ethanolic Extracts Of Garlic, Ginger And Rosemary On The Shelf-Life Of Orange Juice. *International Journal of Agriculture and Policy Research*, 1(01): 197–204.
- [35] Adesokan, I., Odetoynbo, B. B, T. and Okanlawon, M. (2014). Optimization of Lactic Acid Production by Lactic Acid Bacteria Isolated from Some Traditional Fermented Food in Nigeria. *Pakistan Journal of Nutrition*, 8(5): 611- 615.
- [36] Izah, S. C., Kigigha, L. T., Aseibai, E. R. and Okowa, I. P. (2016). Advances in Preservatives and Condiments used in Zobo (a food-drink) production. *Journal of Biotechnoly and Research*, 2(3): 104–119.
- [37] Ubogu, M. (2018). Effects Of Ethanolic Extracts Of Garlic , Ginger And Rosemary On The Shelf-Life Of Orange Juice. *International Journal of Agricultural and Policy Research*, 1(7):1–9.
- [38] Ajayi, O., Ola, O.O. and Akinwunmi, O. O. (2017), Effect of drying nethod on nutritional composition, sensory and antimicrobial properties og ginger (*Zingiber officinale*). *Journal of international Food Research*, 24(2): 614- 620. .