

## **Original Research Article**

# **PREVALENCE AND PATTERN OF WORKPLACE VIOLENCE AMONG NURSES AND OTHER HOSPITAL PERSONALS IN TWO SELECTED HOSPITALS IN OGUN STATE.**

## **ABSTRACT**

Workplace violence against nurses and other health workers is a complex and persistent occupational hazard facing the healthcare profession. The job sector with the mission to care for people appears to be at the highest risk of workplace violence. Nurses are among the most assaulted workers in the Nigeria workforce. Too frequently, nurses are exposed to violence, primarily from patients, patients' families, visitors and other health care workers. This violence can take the form of intimidation, harassment, stalking, beatings, stabbings, shootings, and other forms of assault. It has been shown that workplace violence affects every professional group in every country and sometimes to an 'epidemic' extent. In the health care sector, all categories of healthcare workers are at risk of violence though at different degrees, hence this study point out some of the violence melted against nurses and other healthcare workers.

## **INTRODUCTION**

Workplace violence is an act or threat of physical violence, psychological violence, harassment, intimidation or other threatening disruptive behaviour that occurs at the worksite between and among workers. It ranges from threats and verbal abuse to physical assaults and even homicide. Workplace violence is not restricted to particular occupation or geographical location, but rather cuts across all work environments and countries (Khoshknab et al, 2016). The workplace violence acts, committed by patients, visitors, and/or coworkers and many others; usually raises concerns about personal safety of every worker (Papa & Venella, 2018). Though, many cases of

workplace violence are not reported, those reported are substantially disturbing; despite the various violence preventive control that is emplaced; due to so many damaging effects of such violence, which sometimes resulted to the absenteeism of the receiving victims from work (Tribune newspaper, 2020; NIOSH, 2002). Surprisingly, the health sector accounts for about 25% of all violent workplace accidents, and more than half of all health workers have already been victims of violence attacks (Hahn et al, 2017). According to Clark (2016), nurses are the most attacked group in the health-care industry, accounting for almost 80% of all assaults, a figure that is likely underestimated. Owing that, it is generally agreed that the health care industry is the most susceptible to workplace violence due to rivalry among professionals and a host of unruly or disgruntled patients; it is important to establish policies, awareness, physical security and training that will reduce workplace violence to the barest minimum. Violence at work claims the lives of nearly a million people worldwide (WHO, 201 b2). While almost a million Britons were victims of workplace violence. between 2009 and 2011 (Occupational Safety and Health Administration (OSHA), 2015). Nearly 2 million Americans are victims of workplace violence each year (OSHA, 2015; Wurim, 2016), as many more cases go unreported (United States Department of Labour, 2011; Findorff, McGovern, Wall & Gerberich, 2005 in OSHA, 2015). The situation is similar in Africa, Asia, Latin America and in many developing countries (Wurim,2016). Workplace violence among nurses varies by country and location, with rates as high as 29.9% in Ethiopian public health facility (Fute et al, 2015) and 86.1% in Egyptian obstetrics and gynecology departments (Samir et al, 2015). Nigeria is no exception when it comes to workplace violence against nurses. In a tertiary hospital in Abia state (Ogbonnaya et al, 2015), alarming rates as high as 88.1% were observed, and 53.5% in a South-Western City (Abodunrin et al, 2014). According to a study conducted in Osun State, there is

67% incidence rate of workplace violence within the hospital. Hence every establishment, department and unit must maintain a violence free workplace environment, as part of ways of maintaining that, threatening behaviour is not allowed, by providing preventive guidelines and quick response mission for unforeseen violent accidents. As a result, the purpose of this study is to look into the prevalence and patterns of workplace violence exposure and its reported impacts between Nurses and other health practitioners in few but major health-care settings in Ogun State, Nigeria. After seeking for ethical approval from Ministry of Health via the Federal Medical Centre Abeokuta, authority of Registered Nurses and hospital management team; a self-developed questionnaire which aimed at capturing prevalence and pattern of workplace violence was sent to the forum of registered nurses in Ogun state to access a cross section of nurses that can be conveniently sampled. Possibly due to the fact that many nurses don't like reporting violence attack, most especially when there is no temporary or permanent damage, a total of 372 filled questionnaire was returned out of 1000 after consistent persuasion. Out of the 372 returned questionnaires, only 257 were properly filled. The data which was entered into statistical package of social science software (SPSS) was analyzed via descriptive statistics and tested for differences and association using recently published discrete statistical analysis (Soyinka and Olosunde, 2022). The questionnaire which allowed respondents chose from likert scales developed for each of the measured interest in the questionnaire was able to access the prevalence and pattern of violence attack to Nurses by Nurses and other health professionals. The questionnaire which contains some social demographics also imbibed the WHO-CDC workplace violence categorization. An acceptable Cronbach's alpha value of 0.812 was obtained for the questionnaire.

Table 1. Respondents Characteristics

<b>Socio-demographics</b>	<b>Frequency</b>	<b>Percentage</b>
<b>Age</b>		
20-30	131	51
31-40	49	19.1
41-50	57	22.2
>50	20	7.8
<b>Qualification</b>		
BNSC	151	58.8
RN	36	14
RN/RM/RPHN	70	27.2
<b>Years of Experience</b>		
1-10	142	55.3
11-20	85	33.1
21-30	14	5.4
31-35	16	6.2
<b>Marital Status</b>		
Married	164	63.8
Single	79	30.7
Widow	14	5.4
<b>Hours per week</b>		
<20	24	9.4
20-40	91	35.4
>40	142	55.3

Majority of the respondents were in the age bracket of 20-40 (n=180, 70.1%) with a bachelor of nursing certificate (n=151, 58.8%) and at least 20 years of experience in service (n=227, 88.4%). About 63.8% of the respondents are married and they experienced more than 40 hours of week per week. The demographic statistics showed that 87.5% (225) of the respondents have experienced violence attack in the course of their job.

Prevalence of workplace violence attack based on nurses experienced is presented in table 2 and

<b>Socio-</b>							
---------------	--	--	--	--	--	--	--

<b>demographics</b>							
<b>Age</b>	Nurse	Senior nurse	Health	Doctors	Patients	Patient relative	Security
20-30	48 (53.3)	48 (44)	53 (52)	54 (54.5)	89 (50)	92 (46.7)	26 (33.8)
31-40	8 (8.9)	27 (24.8)	27 (26.5)	25 (25.3)	39 (21.9)	39 (19.8)	27 (35.1)
41-50	34 (37.8)	34 (31.2)	22 (21.6)	20 (20.2)	34 (19.1)	50 (25.4)	24 (31.2)
>50					16 (19)	16 (8.1)	
<b>Qualification</b>							
BNSC	51 (56.7)	53 (48.6)	48 (47.1)	53 (53.5)	97 (54.5)	111 (56)	40 (51.9)
RN/RM/RPHN	22 (24.4)	35 (32.1)	36 (35.3)	30 (30.3)	56 (31.5)	62 (32)	22 (28.6)
<b>RN</b>	17 (18.9)	21 (19.3)	18 (17.6)	16 (16.2)	25 (14)	24 (12)	15 (19.5)
<b>Years of Experience</b>							
1-10	50 (55.6)	51 (46.8)	52 (51)	51 (51.5)	94 (52.8)	99 (50.3)	26 (33.8)
11-20	36 (40)	54 (49.5)	44 (43.1)	40 (40.4)	62 (34.8)	76 (38.6)	47 (61)
21-30	4 (4.4)	4 (3.7)	6 (5.9)	8 (8.1)	8 (4.5)	8 (4.1)	4 (5.2)
31-35					14 (7.9)	14 (7.1)	
<b>Marital Status</b>							
Married	68 (75.6)	85 (78)	68 (66.7)	69 (69.7)	102 (57)	117(59)	61 (79.2)
Single	22 (24.4)	24 (22)	34 (33.3)	30 (30.3)	62 (35)	66(34)	16 (20.8)
Widow					14 (8)	14 (7)	
<b>Hours per week</b>							
<20	2 (2.2)	8 (7.3)	5 (5.1)	5 (4.9)	15 (8.4)	15 (8)	7 (9)
20-40	41 (45.6)	44 (40.4)	30 (30.3)	19 (18.6)	70 (39.3)	68 (34)	6 (8)
>40	47 (52.2)	57 (52.3)	64 (64.6)	78 (76.5)	93 (52.2)	114 (58)	64 (83)

Table 2 Demographic survey results

it revealed the statistical properties in table 3. From table 3, the study showed that at an average an approximately 41%-45% of the nurses experiences workplace violence regularly from their colleagues, senior colleagues doctors, other health professionals and outsiders; with a ratio tendency that one out of every three nurses is evidently attacked violently at his/her workplace regularly. This violent attack on nurses will definitely continue since  $\alpha_3 > 2.72$  and  $\alpha_4 > 4.9$  till a large proportion of the nurses 2%-77% are now attacked with impunity on what looks like a daily basis, if something drastic is not done urgently to curtail such a bizarre situation.

	<b>Socio-demographics</b>						
<b>Statistic</b>	Nurse	Senior nurse	Health	Doctors	Patients	Patientrelative	Security
Shape (b)	0.9429	0.9394	0.9418	0.9427	0.9420	0.9398	0.93
Proportion (p)	0.421	0.4345	0.417	0.4173	0.422	0.4254	0.4409
Ratio (q)	0.3337	0.333	0.3438	0.3346	0.3383	0.3405	0.347
Mean	5.911	5.9	5.888	5.911	5.898	5.887	5.8485
Variance	53.61	53.58	53.53	53.61	53.57	53.54	53.4
Skewness	2.7234	2.7239	2.724	2.7234	2.7237	2.724	2.7263
Kurtosis	4.9294	4.9344	4.9397	4.9294	4.9347	4.9396	4.96
95% CI (p)							
Lower	0.02	0.02	0.02	0.02	0.02	0.02	0.02
Upper	0.7685	0.774	0.76	0.768	0.765	0.7663	0.7735

Table 3 Socio-demographics analysis

Keen investigation into the pattern of violence that are more common within the hospital environment showed that, every two nurses from three nurses, about 70% of nurses above, experience various pattern of violence at regular interval across its ranks and file. It is obvious the violence in the hospital environmental is at an alarming rate as this study showed that nobody is spear from attack which ranges from worker to worker attack ( $p=0.421$ ,  $q=0.3362$ ), to attack due to personal interaction ( $p=0.422$ ,  $q=0.3426$ ), and to attack as a result of patients' relative attack on other health workers ( $p=0.4254$ ,  $q=0.3437$ ), not in the ward; this may be due to perceived hostile, hospital environment or hospital policy. This implies that all the standard categories of workplace violence are present in the hospital setting at almost equal rate.

	<b>Pattern of Workplace Violence</b>				
<b>Statistic</b>	Undermining Work	Verbal Abuse	Physical Abuse	Continual Criticism	Unfriendly engagement
Proportion (p)	0.4167	0.4516	0.3537	0.3495	0.2054
Ratio (q)	0.7715	0.76	0.757	0.7624	0.737
Correlation	0.016	0.0154	0.0261	0.026	0.0582
Non-linear association	0.3645	0.336	0.201	0.2292	0.1513

Table 4 Pattern of

## Workplace Violence

The pattern of violence the respondents agreed that are common includes undermining work, verbal abuse, continual criticism, unfriendly engagement and physical abuse while sex abuse was almost unanimously agreed to be insignificant. The respondents currently experiencing the different pattern of workplace violence was averagely between 35%-46%. Though the workplace violence was not directed to any particular nurse rank, as all nurses irrespective of status experience various attacks, the range of pattern of workplace violence experienced by nurses from different attackers which are fellow nurses, senior nurses, doctors, health workers, security, patients and patients relative, varies from 15%-37% as presented in the descriptive statistics below.




Figure 1. Pattern of work place violence.

Note that patterns like destroying confidence often, ignoring opinions and views, exposing to unmanageable work, humiliation and ridiculing, bullying by patient are all categorized under unfriendly engagement.

## CONCLUSION

This study has been able to show that workplace violence is common among the nurses. Undermining work, due to unmanageable and unhealthy workload, was the most prevalent form of violence experienced by nurses followed by verbal abuse and continuous criticism. Though violence from doctors was relatively high, the proportion was however not statistically significantly different from that of attacks from other party considered in the study. The most disheartening of all the unhealthy attacks is that perpetrators suffered no consequences. It is thus

important that, problematic environment leading to physical abuses and unfriendly engagement (humiliation and ridiculing) at work place, should be checkmated by policies, while health personals should be advised to tolerate and respect each other.

## REFERENCES

Khoshknab, M., Tamizi, Z., Ghazanfari, N., & Mehrabani, G. (2012). Prevalence of workplace violence in psychiatric wards, Tehran, Iran. *Pakistan Journal of Biological Sciences*, 15(14), 680-684. Retrieved January 20, 2016, from <https://www.ncbi.nlm.nih.gov/pubmed/24171249>.

Papa, A., & Venella, J. (2013). Workplace violence in health care: Strategies for advocacy. *Online Journal of Issues in Nursing*, 18(1), 5. Retrieved February 28, 2016, from <https://www.ncbi.nlm.nih.gov/pubmed/23452201>.

National Institute for Occupational Safety and Health (NIOSH). (2015). Workplace violence prevention for nurses. Retrieved January 20, 2016, from <http://www.cdc.gov/niosh/topics/violence/>

Hahn, S., Hantikainen, V., Needham, I., Kok, G., Dassen, T., & Halfens, R. (2012). Patient and visitor violence in general hospital, occurrence, staff interventions and consequences: a cross sectional survey. *Journal of Advanced Nursing*, 68(12), 2685-2699.

Occupational Safety and Health Administration (OSHA). (2015). Guidelines for preventing workplace violence for healthcare and social service workers. Retrieved January 20, 2016, from <https://www.oshgov/Publications/OSHA3826.pdf>.

Clark, K. (2016). The aftermath of violence: The lived experience phenomena of assault in nursing. Walden university scholar works.

World Health Organization. Global campaign for violence prevention: Violence prevention alliance. (2012). Retrieved January 20, 2016, from <http://www.who.int/violenceprevention/approach/definition/en/>

Wurim, B. (2013). Workplace Violence and the Cost-Benefit Trade Off of Zero-Tolerance Safety Policies in Central Nigerian Hospitals. *International Journal of Business and Social Science*, 4(8). Retrieved February 8, 2016, from <http://ijbssnet.com/journals/Vol 4 No 8 Special Issue July 2013/21>

U.S. Department of Labour Occupational Safety and Health Administration. (2015). Retrieved January 12, 2016, from <https://www.osha.gov/publications/osha3148.pdf>.

Samir, N., Mohammed, R., Moustafa, E., & Abou Saif, H. (2012). Nurses' attitudes and reactions to workplace violence in obstetrics and gynaecology departments in Cairo hospitals. *East Mediterranean Health Journal*, 18(3), 198-204.

Occupational Safety and Health Administration (OSHA, 2015)

United States Department of Labour, 2011; Findorff, McGovern, Wall & Gerberich, 2005 in OSHA, 2015.

*South American Journal of Nursing Special Edition*, 2016.

Soyinka, A.T. and Olosunde, A.A. (2022). On discretization of continuous random variables for contingency tables: discrete Johnson systems of distribution as a case study with applications.

*Journal of Statistical Theory and Practice*. 16(4), Published by

Springernature.com.Url:doi.org/10.1007/s42519-022-00290.

The Emergency Nurses Association (ENA) position paper in 2016.

The National Institute for Occupational Safety and Health (NIOSH, 2015).

Guardian newspaper, 2016 Nurse assaulted by patient.

Papa & Venella, 2018 Department of Health and Human Services [DHHS] 2018.

*Safety and Health at work journal*, 2021.

Sue 2015, The first step to freedom from domestic abuse retrieved from

<http://www.thefirststep.org.uk>.

*International journal of public health* 2021.

*Panafrican medical journal* 2017.

*BMC Oral health article*, 2019.

*Ijamhr Journal*, Prevalence and Patterns of Psychological and Physical Violence among Nurses in a Public Tertiary Health Facility in Enugu, Southeast Nigeria, 2020.

Nigerian journal of nursing, workplace violence among nurses in general hospitals in osun state,  
Nigeria, 2019.

UNDER PEER REVIEW