

CONSTRAINTS IN RICE FALLOW PULSES PRODUCTION IN SRIKAKULAM DISTRICT OF ANDHRA PRADESH

ABSTRACT

Rice fallow cultivation constitutes the major portion of pulse production particularly black gram and green gram in Srikakulam district of A.P. Farmers are reaping lower yields of 3.75-5.00 q/ha against 15.0-20.0 q/ha of its potential. Keep this in view an extension study was conducted on the study of constraints and reasons for low productivity in Pulses in Srikakulam District. Ex-post-facto research design was used for the study with a total sample size of 60. Results revealed that the low soil fertility was the major constraint in pulses production (100%), the incidence of yellow vein mosaic virus (93.33), labour scarcity and incidence of low temperatures and fog during the flowering period i.e., the first fortnight of January (88.33%), weed infestation particularly vicia, cuscuta, xanthium and Echinochloa (86.67%). Lack of YVMV resistant varieties and suitable varieties (83.33%) and lack of a regulated market (96.67%) were the major technical or situational constraints. Non-adoption of irrigation management (93.33%), non-adoption of seed treatment (91.67%), non-adoption of proper sowing (way of broadcast) method and seed rate (90.00), non-adoption of weed management practices (88.33%), non-adoption of foliar nutrient management in rice fallow pulses (86.67%) were the adoption gaps perceived as constraints. There is a need to conduct more Front Line Demonstrations (FLDs) in the farmer fields, more training programmes to the farmers and timely supply of inputs or make available to the farmers prior to the season.

Key words: Knowledge, Adoption, Partial adoption, Constraints, Situational constraints

1. INTRODUCTION

Pulses are major sources of proteins among vegetarians in India and complement the staple cereals in diets with proteins, essential amino acids, vitamins and minerals. They contain 22-24% protein, which is almost twice the protein in wheat and thrice that of rice. Pulses contribute 11 per cent of the total

intake of proteins in India (1). Rice fallow cultivation constitutes the major portion of pulse production particularly black gram and green gram in Srikakulam district of A.P. Farmers are reaping lower yields of 3.75-5.00 q/ha against 15.0-20.0 q/ha of its potential. Keep this in view an extension study was conducted on the study of constraints and reasons for low

productivity in Pulses in Srikakulam District as per the approval of the State Level Technical Programme (SLTP)2021(2).This study was conducted by the Extension Scientist of the District Agricultural Advisory and Technology Transfer Centre(DAATTC) of Srikakulam, an extension unit of Acharya N.G Ranga Agricultural University.

2. MATERIALS AND METHODS

The study was conducted in selected six Mandals purposively based on crop coverage (pulses). They are Santhakaviti (3838ha), Palakonda (3660ha), Veeraghattam(3646ha), Burja(3286ha), Polaki (3840ha) and Narasannapeta (3000ha) Mandals. Two villages from each Mandal were selected. They are Mandavakurthy (200ha) and Manthina (145ha) of Santhakaviti Mandal, Velagada (112ha) and Vadama ((128ha) of Palakonda Mandal, Nadimikella (98ha) and Nadukuru (95ha) of Veeraghttam Mandal, Neelakantapuram (102ha) and Lakkupuram(95ha) of Burja mandal, Thotada(96ha) and Vurjam(92ha) of Polaki Mandal, Vurlam(112ha) and

Lukalam(85ha) of Narasannapata Mandal. Five farmers from each village were selected randomly for the study. Thus the total sample size was 60. Ex-post-facto research design was used for the study as the variables already occurred. The interview schedule was prepared and data collected from the farmers on knowledge, adoption and constraints faced and as perceived by them.

This study was based on a descriptive analysis of the collected data from the selected sample of farmers. Hejase et al. [3] contend that informed objective decisions are based on facts and numbers, real, realistic, and timely information. Furthermore, according to Hejase and Hejase [4], “descriptive statistics deals with describing a collection of data by condensing the amounts of data into simple representative numerical quantities or plots that can provide a better understanding of the collected data” (p. 272). Therefore, this study used frequencies and percentages as primary data to conduct further analysis. The data were quantified and ranked based on frequency and percentage of respondents.

3. RESULTS AND DISCUSSION

Results were tabulated based on the cumulative frequency of variables and their corresponding percentages. Total cumulative frequency of responses 720 (i.e.,12question items for 60 farmers of the sample, N=60)

Table no.1: Knowledge level of respondent farmers:

S.No	Knowledge item	Cumulative Frequency of responses	Per centage
1	Not Awareness	302	41.94
2	Awareness	244	33.89
3	Knowledge	174	24.17
	Total	720	100

Table 1 and Figure 1 show that about 42 % per cent of the farmers not aware of the package of practices of pulses particularly black gram and green gram as per the recommendations of ANGRAU[5] followed by those farmers who are aware (33.89%) and those who have the knowledge by 24.17 %.

The above-mentioned results may be due to the low education (literacy) and less extension contact of farmers[6]. Item analysis revealed that the farmers not aware about the recommended package of practices particularly use of farm machinery, sowing method, foliar nutrient management and the existence of high- yielding varieties.




Figure 1. Respondent farmers' Knowledge level

Table No.2: Adoption of package of practices of Rice fallow pulses:

S No	Adoption items	Cumulative Frequency of responses	Per centage
1	Non Adoption	344	47.78
2	Partially adoption	249	34.58
3	Full Adoption	127	17.64
	Total	720	100

It can be observed from Table 2 and Figure that about 48 % of the farmers did not adopt the recommended package of practices of black gram and green gram.

About 35% of the farmers partially adopted the recommended package of practices of black gram and green gram. Finally, about 18% of the respondent

farmers adopted the package of practices as per the recommendations. A low adoption rate might be due to the lack of

knowledge of recommended technologies and situational constraints[6],[7].




Figure2. Adoption of Pulses production technologies

Constraints faced and perceived the respondent farmers:

Based on the literature [8], [9],[10] and on discussion with experts constraints are categorized into technical or situational and extension gaps perceived as constraints.

It can be revealed from Table 3 that low soil fertility was the major constraint in pulses productivity (100%).It might be due to the fact that the majority of the Srikakulam soils are red sandy loams, non-application of FYM and green manure years together, followed by the incidence of yellow vein mosaic virus (93.33%),

labour scarcity, and incidence of low temperatures and fog during the flowering period i.e., first fortnight of January (88.33%), Weed infestation particularly vicia, cuscuta. xanthium and Echinochloa (86.67%) and the lack of YVMV resistant & suitable varieties (83.33%) and the lack of a regulated market (96.67%) were the major constraints. There is a need to create awareness among the farmers about the availability of resistant varieties and the measures to mitigate fog incidence.

Table 3: Situational or technical constraints:

S.No	Constraints	Frequency	Percentage	Rank
1	Low soil fertility (low organic carbon)	60	100	I
2	Lack of YVMV resistant varieties & suitable varieties	50	83.33	VI
3	Incidence of Yellow Mosaic Virus Disease	56	93.33	II
4	Uncontrolled water flow at the time of sowings	28	46.67	XIII
5	Cropcan not withstand even little water stagnation	40	66.67	X
6	Prevalence of moisture stress at the time flowering and pod development	45	75.00	VIII

7	Incidence of Low temperatures and fog during flowering period i.e first fort night of January	53	88.33	IV
8	No possibility of inter cultivation	36	60.00	XI
9	Weed infestation particularly vicia, cuscuta. xanthium and Echinochloa	52	86.67	V
10	Not withstand to soil salinity which is increasing year by year	42	70.00	IX
11	Incidence of maruca & spodoptera infestation	48	80.00	VII
12	Pod dehiscence at the time of harvesting	30	50.00	XII
13	Labour scarcity	55	91.67	III
14	Incidence of powdery mildew	25	41.67	XIV
	Market related constraints			
15	Lack of regulated market	55	91.67	I
16	Price fluctuation	42	70.00	II

Table No.4 Extension gaps perceived as constraints

S.No	Constraint particulars	Frequency	Per centage	Rank
1	Due to lack of knowledge and ignorance	42	70.00	IX
2	Non- adoption of existing high yielding varieties	48	80.00	VI
3	Supply of seed by the Department not in time	35	58.33	X
4	Non- adoption of seed treatment	55	91.67	II
5	Non- adoption of proper sowing (way of broadcast) method and seed rate	54	90.00	III
6	Non- adoption of sowing window in time	45	75.00	VIII
7	Non- adoption of foliar nutrient management in rice fallow pulses	52	86.67	V
8	Non- adoption of weed management practices	53	88.33	IV
9	Non- adoption of pest management	46	76.67	VII
10	Non- adoption of irrigation management	56	93.33	I

Table 4 depicted that Non-adoption of irrigation management (93.33%) was perceived as a constraint, it might be due to two reasons one is due to the apprehension of the withering of plants due to excessive moisture and another one due to the non availability of water at that time. Non - adoption of seed treatment (91.67%) is another constraint, it might be due to a lack of awareness of its benefits as it reduces the incidence of sucking pests. Non- adoption of proper sowing (way of broadcast) method and

seed rate (90.00%) due to apprehension of poor germination and ignorance .Non adoption of weed management practices(88.33%) is another major constraint, it might be due to a lack of knowledge on weedicides suitable for different weeds, lack of moisture at that time, and apprehension about the injury to the crop. Non adoption of foliar nutrient management in rice fallow pulses (86.67%). In rice fallow pulses application of fertilizers is not possible and for better growth and yield it is very required to spray foliar nutrients but most of

the farmers did not follow .Non- adoption of existing high-yielding varieties (80.00%). There are certain YMV resistant varieties like LGB787,TBG104,GBG1 of black gram and TM96-2,IPM2-14, and LGG460 varieties of green gram are available but most of the farmers did not adopt. So there is a need to

Conclusions

Low soil fertility was the major constraint in pulses productivity, Non- application of FYM and green manure years together, followed by the incidence of yellow vein mosaic virus, laour scarcity, and incidence of low temperatures and fog during the flowering period *i.e.*, first fort night of January, Weed infestation particularly vicia, cuscuta. Xanthium and Echinochloa,

Acknowledgements

I am very thankful to my Associate Director of Research, RARS, Anakapalle, Dr M Bharata Lakshmi who gave me full cooperation in doing my work; I would also like to thank my university for timely instructions.

COMPETING INTEREST

Authors have declared that they no known competing financial interestsor Non-financial interests or personal relationship that could have appeared to influence the work reported in this paper.

AUTHORS Contributions

This work was carried out in collaboration among authors. Author P VenkataRao

conduct Front Line Demonstrations in the farmer's fields[11]. Non-adoption of pest management (76.67%), it might be due to a lack of knowledge on pests, ignorance and negligence. Non -adoption of sowing window in time (75.00%) were the major constraints.

Non-adoption of irrigation management, Non-adoption of seed treatment, Non-adoption of weed management practices were the major constraints in pulses production. There is a need to conduct more FLDs in the farmer fields and training programmes for the farmers and timely supply of inputs or make them available to the farmers prior to the season.

designed the study prepared the schedule and collection of data Author G Chitti babu performed the statistical analysis, Author S Neelaveni wrote the protocol and wrote the first draft of the manuscript. Author J. Jagannadham corrected the final manuscript. Authors read and approved the final manuscript.

References :

1. Reddy AA. Regional disparities in food habits and nutritional intake in Andhra Pradesh, India. Regional & Sectoral Econ Stud. 2010;10 (2):125-134.
2. NAARM. Annual Report (2021). ICAR-National Academy of Agricultural Research Management, Rajendranagar, Hyderabad-500030; 2022, pp. 1-192, ISSN: 2349-7548.
3. Hejase HJ, Hejase AJ, Hejase HANJ. Quantitative Methods for Decision

- Makers: Management Approach. Beirut, Dar Sader Publishers; 2012.
4. Hejase AJ, Hejase HJ. Research Methods: A Practical Approach for Business Students (2nd edition). Philadelphia, PA, USA: Masadir Incorporated; 2013.
 5. Ranga AN. ANGRAU Blackgram Outlook Report-January to December 2021. Available at: https://angrau.ac.in/downloads/AMIC/OutlookReports/2021/2-BLACKGRA_January%20to%20December%202021.pdf (Accessed: 2 March 2023).
 6. Choudhary AK. Technological and extension yield gaps in pulse crops in Mandi district of Himachal Pradesh, India. *Indian J of Soil Cons.* 2013;41:88-97.
 7. Saravanakumar S. Impact of Cluster Frontline Demonstration on Black gram in Western Zone of Tamilnadu. *Journal of Krishi Vigyan*;2018,7(1): 136-139
 8. Kumar N, Hazra KK, Singh S, Nadarajan N. Constraints and prospects of growing pulses in rice fallows of India. *Indian Farming*; 2016,66(6):13-16.
 9. Kumar N, Singh SS, Ghosh PK, Singh NP, et al. Issues and strategies for the promotion of pulses in untapped rice-fallow in India: A review. *Journal of Food Legumes*; 2020, 33(3): 139-150.
 10. NAAS. 2013. Improving productivity of rice fallows. Policy Paper No. 64, National Academy of Agricultural Sciences, New Delhi; 2013, 16 p.
 11. Venkata RP, Chittibabu G, Chinnam ND. Impact of Front Line Demonstrations on Integrated Crop Management in Rice Fallow Black Gram in Srikakulam District of Andhra Pradesh *Journal of KrishiVigyan*; 2020, 9(1):189-192