

With a new method measurement of Zn, Cu, Cu/Zn levels in third-trimester pregnant women

ABSTRACT

Objective: Our aim in this study; is to determine zinc, copper and copper/zinc ratios in third-trimester pregnant women with a new laboratory method.

Method: In this research conducted on 59 pregnant women in the last trimester, Those diagnosed with preeclampsia, preterm birth, postmaturity, and gestational diabetes were excluded from the study. In pregnant blood serums, zinc and copper levels were studied on fully automatic analyzers with a new method. Additionally, copper/zinc ratios were calculated. In addition, determined the APGAR (Activity-Pulse-Grimace-Appearance-Respiration) score of the babies. Statistical studies of the obtained data were performed at $p < 0.05$ significance level.

Results: Zinc and copper parameters; No statistically significant relationship was found in the analyzes performed with gestational age, gestational week, birth weight, APGAR score, gravida and the number of living children ($p > 0.05$). In addition, it was observed that there was no difference in zinc and copper/zinc ratios in the evaluations made according to the gender of the baby ($p > 0.05$). On the other hand, copper levels in pregnant women; were found to be significantly higher in female fetuses (200.72 ± 36.36) than in males (164.17 ± 42.47) ($p < 0.001$).

Conclusion

Although the copper values of pregnant women with female fetuses are significantly higher, these results require more comprehensive studies. On the other hand, there is not relationship between zinc, copper and copper/zinc values and ABGAR scores.

Keywords: Third trimester, pregnancy, zinc, copper, copper/zinc

INTRODUCTION

Although correct and healthy nutrition is critical during pregnancy, it is possible to detect this situation with test parameters. Among these tests is copper, zinc values and copper/zinc ratio in blood serum. Of course, the nutritional needs of individuals are different. Factors such as body size and the presence of pregnancy bring about energy consumption differences. Therefore, reference levels for the adequacy of diets are also set for pregnant women. The reference and recommended intakes for zinc and copper in pregnant women are shown in the

table (Table 1). However, these tables may differ slightly in European countries (David AB, Chapter 44, 2018).

Table1. Recommended dietary allowances and acceptable intakes for Zn and Cu

	Zn (mg)	Cu (mg)
Pregnant	11	1000
Lactating	12	1300

Source: Standing Committee on the Scientific Evaluation of Dietary Reference Intakes, Food and Nutrition Board, Institute of Medicine Dietary Reference (David AB, 2018).

When zinc and copper are not taken in sufficient amounts in the diet, metabolic function losses and various diseases occur as a result (Table 2). A higher-than-normal serum copper level is defined as hypercupremia, and a low level is defined as hypocupremia. Copper, which is effective in the mobilization of iron, participates in redox reactions as a trace element. In addition, copper is included in the structure of enzymes necessary for the use of iron in the formation of haemoglobin and is necessary for the activities of these enzymes. Copper deficiency is encountered in various malabsorption syndromes and causes cardiovascular disease symptoms. Again, copper deficiency is seen in Menkes disease, which is a rare, X-linked neurodegenerative disease. The daily Zn requirement is up to 15 mg. Most of the dietary Zn is absorbed from the small intestine by active transport. Zinc absorption requires a “zinc-binding exocrine ligand” secreted by the pancreas (David AB, Chapter 43, 2018). In its deficiency, regression in growth and skeletal development, hepatosplenomegaly, acrodermatitis enteropathica, skin rashes in infants, and decreased sense of taste and smell. Zinc; It participates in the structure of carbonic anhydrase, alcohol dehydrogenase, glutamate dehydrogenase, kidney phosphatase, carboxypeptidase, and uricase enzymes. It improves ischemic conditions by providing enlargements in cerebral vessels and coronary arteries. If you are in the zinc-deficient nutrition server, Acrodermatitis enteropathica and susceptibility to infections, in addition, various subclinical conditions arise (David AB, Chapter 44, 2018). In our study, the blood serum in the third trimester of pregnancy determined zinc, copper and copper/zinc ratio. Used a newly described method for zinc and copper measurements; caught the postnatal APGAR (Activity-Pulse-Grimace-Appearance-Respiration) score of the babies by scoring. Our study is a first in the literature considering the measurement techniques of the zinc and copper test.

Table2. Zn, Cu Functions and deficiency disease

	Zn (mg)	Cu (mg)
Functions		

	Function as prosthetic groups in Enzymes	Function as prosthetic groups in enzymes
Deficiency Disease	Acrodermatitis Enteropathica. Preteral Nutrition. Infectious Disease. Subclinical Effects of Deficiency	Menkes Syndrome. Malabsorption Syndromes. Cardiovascular Disease

Source: This table has been adapted from the Tietz section IV chapter (Burtis CA, 2006).

MATERIAL AND METHOD

The ethics committee of the study was recruited from Sakarya University (24.09.2018 date, 216 numbers). Fifty-nine pregnant women in the last trimester of pregnancy and who had a singleton pregnancy were included in the study. Accepted 37-40 weeks for the definition of the previous trimester. Pregnant women who were diagnosed with preeclampsia, preterm labour, postmaturity, and gestational diabetes during the study were excluded from the study. Exclusion criteria were gastrotaemia, small bowel resection, peptic ulcer, psychiatric disease, pregnancy toxemia, bronchial asthma, active hepatitis, cancer, chronic renal failure, and heart failure. In addition to the test parameters values examined in the research, it also looked at the baby's sex, ABGAR-1, ABGAR-5, gestational age, number of living children and fetal gender.

Laboratory process in the study

For zinc and copper detection, venous blood was drawn into special tubes with clot activator while patients were fasting (dark blue capped trace element tubes, BD vacutainer trace element testing). After the blood was drawn, it was inverted 8-10 times and the samples were transferred to the laboratory following the cold chain. Subsequently, the samples were subjected to refrigerated centrifugation (4000 rpm for 10 minutes). The obtained sera were stored in capped Eppendorf tubes (isolab centrifuge tubes 2.0 ml) at -80. Rel Assay Diagnostics kit was used on the fully automatic AU 680 analyzer (serial number: 2016024580, Tokyo, Made In Japan) on a working day. The zinc ion in the sample reacted with 5-Br-PAPS and gave absorbance at 548 nm wavelength in direct proportion to the total zinc level. The copper ion reacted with 3,5-DiBr-PAESA and showed absorbance at 572 nm wavelength in direct proportion to the total copper level. Zinc and copper measurements were made in this way by the colourimetric method (Table 3). In the study, the APGAR score of all babies was scored after delivery and determined.

Table 3. Performance of the tests used in the research

	Normal Range	Linearity	Accuracy	Precision %CV
Zinc (µg/dL)	60-120	4-1000 ^a	0.98 ^Z	2.32
Copper (µg/dL)	110-312 ^X	3-600 ^a	0.97 ^Y	1.85

^X: Pregnancy, ^Y: Correlation coefficient (r), ^Z: Correlation coefficient (r), ^a: Low-High

RESULTS


While there was no statistically significant difference between the zinc levels and zinc/copper ratios between the male and female genders in the evaluation made according to the fetus gender, the copper levels were found to be significantly higher in the pregnant women with a female fetus (Table 4, $p > 0.05$).

Table 4. Comparison of Cu, Zn and Cu/Zn values according to fetus gender

	Girl Fetus	Male Fetus	P
Zinc	86,64±23,81	80,38±24,40	0,346
Copper	200,72±36,36	164,17±42,47	0,001
Zinc/Copper	2,49±0,80	2,23±0,78	0,233
N	25	29	

Student's t test was performed. $P < 0.05$ was considered significant.

As a result of the correlation analysis of zinc, copper and Zinc/copper parameters with gestational age, the number of living children, and APGAR (1 and 5) parameters at the first and fifth minutes determined that there was no significant correlation between them (Table 5 and Graphic 1; $p < 0.05$).



Graphic 1. Comparison of copper levels by fetal sex ($p=0.001$)

Table 5. Relationship of gestational age, number of living children, APGAR-1 and 5 with test parameters

	Zinc		Copper		Copper/Zinc	
	r	p	r	p	r	p
Age	0,156	0,259	0,081	0,559	-0,115	0,407
Living children	0,010	0,941	-0,059	0,672	-0,062	0,656
APGAR-1	-0,067	0,628	0,190	0,170	0,202	0,143
APGAR-5	-0,051	0,714	0,178	0,198	0,189	0,170

Pearson korelasyon analizi yapılmıştır. $P<0.05$ was considered significant.

DISCUSSION

(Squitti R et al., 2002) It has long been known that elevated serum copper levels decrease the antioxidant system and correlate with poor neuropsychological performance and medial temporal lobe atrophy. (Lu CW et al., 2021) Indeed, examined serum levels of zinc, copper and iron in 1165 adults. This study showed that high serum zinc, copper and iron levels are associated with the risk of metabolic syndrome, BMI and the number of metabolic factors independent of insulin resistance. In our study, serum copper levels were found to be significantly higher in women who were pregnant with a girl. (Gao Y et al., 2020) In a study whose data were taken from the 2011-2016 National Health and Nutrition Examination surveys, it was stated that copper induces oxidative stress and zinc counteracts oxidative stress. On the other hand, the same study indicated that causality deserves more confirmation.

(Zhang H et al., 2021) In fertile women aged 18-44 years from a representative population in China, research was done on Zn, Cu and Cu/Zn ratios. A total of 191 women who gave birth to healthy children participated with a strict set of inclusion criteria. Determined baseline biological indicators and basic levels in the whole blood of the included women. After they evaluated the data they obtained, They concluded that “the Zn, Cu and Cu/Zn ratios in plasma and whole blood of fertile women can be used as an indicator to evaluate the reference range, element deficiency and overload status”. Our study, using a new method in zinc and copper measurements and with the results obtained, examined the relationship between ABGAR scores, number of living children, and fetus gender.

(Giddens JB, 2000) A study investigating dietary food intake in the second and third trimesters of pregnancy compared dietary standards in adolescent and adult pregnant women.

According to them, The diets of 59 pregnant adolescents and 97 pregnant adults were inadequate and below the recommended dietary intakes, including zinc. It is seen in our work and their work that; Continuous nutritional monitoring of pregnant adolescents and pregnant adults is required, including nutritional guidance highlighting food sources of zinc vitamins.

CONCLUSION

With zinc, copper and copper/zinc values in pregnant women in the third trimester, There does not appear to be a relationship regarding ABGAR scores. Similarly, there is no relationship between age and the number of living children. On the other hand, the fact that the copper values were significantly higher in pregnant women with female fetuses was seen as a result that requires further research.

REFERENCES

Blondin JH, LoGiudice JA. Pregnant women's knowledge and awareness of nutrition. *Appl Nurs Res.* 2018;39:167-174. doi: 10.1016/j.apnr.2017.11.020.

Cinemre H, Cinemre B, Cinemre S, MD, Çekdemir D, Aydemir B, Tamer A, Yazar H. Diagnosis of vitamin B12 deficiency in patients with myeloproliferative disorders. *J Investig Med.* 2015;63(4):636-40 doi: 10.1097/JIM.000000000000187

David AB, Peter AM. Harper's Illustrated Biochemistry. Lange. 30 th Edition, 2018. Sexion IX, Chapter 43 Nutrition, Digestion, & Absorption. Pp: 537-8.

David AB. Harper's Illustrated Biochemistry. Lange. 30 th Edition, 2018. Sexion IX, Chapter 44 Micronutrients: Vitamins & Minerals. Pp: 546-7.

Gao Y, Liu Y, Wang P, Meng X, Zhang W, Sun Y. Serum Copper and Zinc Levels and Urinary Incontinence in Adult Women. *Biol Trace Elem Res.* 2021 Mar;199(3):842-849. doi: 10.1007/s12011-020-02205-9. Epub 2020 May 28. PMID: 32468225.

Giddens JB, Krug SK, Tsang RC, Guo S, Miodovnik M, Prada JA. Pregnant adolescent and adult women have similarly low intakes of selected nutrients. *J Am Diet Assoc.* 2000;100:1334-40. doi: 10.1016/S0002-8223(00)00377-1.

Guggino A, Barbero S, Ponzo V, Viora E, Durazzo M, Bo S. Myths about nutrition in pregnancy. *J Obstet Gynaecol.* 2016;36(7):964-965. doi: 10.3109/01443615.2016.1168372

Lu CW, Lee YC, Kuo CS, Chiang CH, Chang HH, Huang KC. Association of Serum Levels of Zinc, Copper, and Iron with Risk of Metabolic Syndrome. *Nutrients.* 2021 Feb 7;13(2):548. doi: 10.3390/nu13020548. PMID: 33562398; PMCID: PMC7914992.

Said HM. Intestinal absorption of water-soluble vitamins in health and disease. *Biochem J.* 2011 Aug 1;437(3):357-72. doi: 10.1042/BJ20110326.

Squitti R, Lupoi D, Pasqualetti P, Dal Forno G, Vernieri F, Chiovenda P, Rossi L, Cortesi M, Cassetta E, Rossini PM. Elevation of serum copper levels in Alzheimer's disease. *Neurology.* 2002 Oct 22;59(8):1153-61. doi: 10.1212/wnl.59.8.1153. PMID: 12391342.

The Apgar Score. American Academy of Pediatrics Committee on Fetus and Newborn; American College of Obstetricians and Gynecologists Committee on Obstetric Practice. *Pediatrics.* 2015;136(4):819-22. doi: 10.1542/peds.2015-2651.

Tietz Textbook of Clinical Chemistry and Molecular Diagnostics. Edited by Burtis CA, Ashwood EA, Bruns DA. Fourth Edition. Elsevier Saunders. Printed in the United States of America. 2006. Pp: 1126-1143.

Zhang H, Cao Y, Man Q, Li Y, Lu J, Yang L. Study on Reference Range of Zinc, Copper and Copper/Zinc Ratio in Childbearing Women of China. *Nutrients.* 2021 Mar 15;13(3):946. doi: 10.3390/nu13030946. PMID: 33804217; PMCID: PMC7999022.