

Short Research Article

Nutritional and mineral compositions of Tiger nut (*Cyperus Esculentus*) tubers from different ecotypes in Niger.

Abstract

Nutritional characterization is an essential component for a better knowledge of the different ecotypes of Niger tiger nut. To do this, 20 samples of tiger nut tubers were taken for nutritional investigation, including 5 ecotypes representative of the Niger tiger nut and 15 others randomly selected from the ecotypes collected in the producing villages in the Maradi and Dosso regions. The content of mineral elements was analyzed by atomic absorption spectrophotometer and the content of macronutrients determined by different types of assay. The results showed that the tiger nut ecotypes are rich in lipids (12.00% to 25.20%), carbohydrates (24.50% to 47.70%), fiber (21.90% to 39.40%), protein (2.90% to 11.90%), and mineral elements such as calcium (14.100 ± 5.44 to 65.380 ± 19.940), iron (0.881 ± 0.017 to 15.809 ± 6.524), potassium (102.800 ± 2.970 to 308.410 ± 8.900), magnesium (366.800 ± 1.942 to 471.830 ± 8.927). The fat content is much higher in small tigernut ecotypes with an average of 21.60% than in large tigernut ecotypes with an average of 14.90%. The Copper (Cu), Sodium (Na), Calcium (Ca) and Magnesium (Mg) contents are significantly identical for all the ecotypes studied. On the other hand, the Zinc (Zn), Iron (Fe), Potassium(K) contents of ecotypes E1(Big tigernut), E2 (Small tiger nut), E3(Small tiger nut), E4(Small black tiger nut), E5(Small wild tiger nut), E7(Small wild tiger nut), E8(Small tiger nut), E9(Small tiger nut), E11(Small tiger nut), E12(Small tiger nut), et E15(Big tigernut) are significantly higher compared to the other ecotypes. The study showed that the tiger nut tubers are rich in iron and that the iron composition of the wild ecotype is much higher. Compared to other foods, the tigernut tubers of certain ecotypes studied (E1, E2, E3, E4, E5, E7, E8, E9, E11, E12 and E15) are richer in iron than the leaves of *Moringa oleifera* (3.810 ± 0.120 mg/100g), the pods of *Vigna unguiculata* (7.800 ± 0.150 mg /100g) and leaves of *Hibiscus sabdariffa* (8.270 ± 1.010 mg/100g).

Keywords : Characterization, macronutrients, micronutrients, ecotypes, tigernut, Niger

I. INTRODUCTION

Cyperus esculentus. L (Tigernut) is a perennial monocotyledous plant which has a tough erect fibrous root. The slender rhizomes of tigernut form weak runners above the ground level which develop small-sized tubers at the tip of the stem. Tigernut tubers can reach about six inches depth into the soil. The size of the tubers can be compared with that of peanut. The central erect stem of tigernut is usually covered by sheath of leaves [1].

In Niger, tiger nut is used in human food in different forms (biscuit, sweets, flour, milk, etc.) [2]. Also, the tiger nut, like most of these tubers, makes it possible to fight against malnutrition given their chemical composition and their medicinal properties [3]. Currently, malnutrition is a public health problem in the world. According to the report on global malnutrition 2017, published in November, all of the 140 countries studied are confronted with at least one of the main forms of this scourge: stunting, anemia in women of childbearing age and overweight in adults [4]. This malnutrition continues to plague the world in general

and in developing countries (DCs) in particular [5]. The number of chronically undernourished people has increased by almost 56 million people in Africa (especially West Africa) [6]. In Niger, the prevalence of global acute malnutrition (GAM) is 16% in rural areas compared to 10.4% in urban areas, corresponding to a serious situation according to the WHO (World Health Organization) classification scale [7]. And yet, tiger nut tubers are eaten all over the country. Despite a wide knowledge of tiger nut tubers, no study on its physico-chemical composition has been made in Niger. **The objective of this study is therefore to contribute to a better understanding of the mineral and nutritional compositions of cultivated and wild tiger nut tubers in Niger.**

II. MATERIALS AND METHODS

2.1. Material

a) Plant material

The plant material is made up of twenty (20) samples (small and large) of tiger nut tubers including five (5) ecotypes produced during the trial on the agro-morphological characterization of the five ecotypes representative of the Niger tiger nut and the fifteen (15) others randomly selected from the ecotypes collected at the level of the producing villages of tiger nut in the regions of Maradi and Dosso (Table 1).

Table 1: Different ecotypes analyzed

Sample No.	Kinds	Collection place (Regions
E1	Big tiger nut*	Trial(E1)/ FA	Niamey
E2	Small tiger nut*	Trial (E2)/FA	Niamey
E3	Small tigernut**	Trial (E3)/FA	Niamey
E4	Small Black Tigernut*	Trial (E4)/FA	Niamey
E5	Small Wild Tigernut***	Trial (E5)/FA	Niamey
E6	Small Black Tigernut	Maradi	Maradi
E7	Small Wild Tigernut	Rijiasamna	Dosso
E8	small tiger nut	Dommo	Dosso
E9	small tiger nut	Dalia	Maradi
E10	small tiger nut	Takalafia	Maradi
E11	small tiger nut	Dan Tsoutsou	Maradi
E12	small tiger nut	Gamozon	Dosso
E13	small tiger nut	Kimiakoara	Dosso
E14	small tiger nut	Dan Amina	Maradi
E15	big tiger nut	Dan Toudou	Maradi
E16	big tiger nut	RijiaSamna	Dosso
E17	big tiger nut	dan gamji	Maradi
E18	big tiger nut	Guidan Moussa	Maradi
E19	big tiger nut	Dan Sara	Maradi
E20	big tiger nut	Dan Amina	Maradi

*: Maradi market, **: Dosso market; *** : empty field , FA: faculty of agronomy

2.2. chemical analysis methods

2.2.1 Determination of moisture content and dry matter

The moisture content was determined into crucibles previously dried in an oven at 103° C for 30 min and cooled in a desiccator then weighed (P0), 5 g (TS) of the sample were introduced. These crucibles are placed in an oven at 103°C for three hours, and then weighed at regular time intervals of 15 min until a constant weight is obtained. Thus, the moisture level was determined by the following formula:

$$\% \text{ Moisture} = TS - (FW - EW) / TS \times 100$$

TS = test socket (5 g);

EW = empty weight of the crucibles;

FW = final weight (crucibles + EW).

The dry matter is obtained from this determination of the moisture content

$$\% \text{ DM} = 100 - \% \text{ moisture}$$

2.2.2 Determination of mineral matter (MM) and organic matter (OM)

Incineration was carried out to obtain all the cations (ammonium excluded) in the form of carbonates and other anhydrous mineral salts. To do this, crucibles are dried in an oven at 103°C for 30 min and cooled in a desiccator then weighed (TS). Next, 1 g (EW) of the sample are introduced into these crucibles and brought to incineration in a furnace at 550° C for 4 h. At the end of the incineration, the crucibles are removed and cooled in a desiccator for 30 min before being weighed (FW). The percentage of ash is given by the following relationship:

$$\% \text{ MM} = FW - TS / EW \times 100$$

TS: test socket;

FW: final weight (crucible + sample);

EW: empty weight of the crucibles.

The total Organic Matter (OM) is obtained from the results of the mineral matter or ash. The organic matter content is thus equal to:

$$\% \text{ OM} = \% (\text{DM}) - \% \text{ MM}$$

2.2.3 Dosage of fat

The determination of the fat content is carried out according to the Soxhlet extraction method using hexane as the solvent under reflux [8]. The balloon is first washed and dried. The empty balloon weight (P1) is noted 5g (P2) of the sample is introduced into the extraction cartridges, then plugged with cotton and placed in the soxhlet. The flask is filled with approximately 300 ml of hexane and then connected to the soxhlet. The latter connected to a refrigeration system is connected to a cryostat to condense the solvent vapors intended to entrain the lipids. The

extractions last 4 h. The hexane is recovered through a tap. The flask is dried in an oven at 105° C and cooled in a desiccator for 30 min then weighed. The fat content is obtained according to the following formula:

$$\% \text{ Fat} = (P3 - P0) / P2 \times 100$$

P1 = mass of the empty balloon;

P3 = mass of the balloon containing the fat;

P2= test socket

2.2.4. Determination of total nitrogenous matter

The content of total nitrogenous matter or proteins is determined by the Kjeldahl method. The organic nitrogen of the sample to be analyzed is transformed into mineral nitrogen in ammoniacal form $(\text{NH}_4)_2 \text{SO}_4$ thanks to the oxidizing action of concentrated sulfuric acid in the presence of a catalyst. After displacement with sodium hydroxide (strong base added in excess), the ammonia is distilled then titrated in the presence of a reagent (boric acid) by acidimetry. The total protein content is calculated by multiplying the amount of nitrogen by a conversion factor (6.25), or 16% in protein.

The total nitrogen content (protein) is calculated by the following formula:

$$\% \text{ protein} = ((C_s - C_w) / TS \times N \times 0.014 \times 100) \times 6.25$$

Cs = cruet drop (sample)

Cw = cruet drop (white)

N = normality of the sulfuric acid used for the assay = 0.1

TS= test socket (0.2 g)

6.25 = conversion factor multiplying the amount of nitrogen to get the total protein content

0.014= molar mass of nitrogen \times 0.001

2.2.5 Determination of crude fiber

The method makes it possible to measure by a double hydrolysis in animal feed the materials free of fat and insoluble in acid medium and in alkaline medium called crude cellulose. The assay was carried out according to the Weende method [9]. Indeed, the sample is successively treated with boiling solutions of sulfuric acid $\text{H}_2 \text{SO}_4$ 0.255 N and sodium hydroxide NaOH 0.313 N and washed, dried and then calcined. The weight loss resulting from the calcination corresponds to the crude cellulose of the test portion (1 g). A concentrated solution corresponds to 35.64 N To obtain a 0.255 N solution, 14.3 ml of concentrated sulfuric acid must be taken and made up to 2 l with distilled water. For 2l weigh 25.04g and top up with distilled water.

$$\% \text{ Crude fiber} = ((P2 - P3) / P1) \times 100$$

P1: test socket (g)

P2: weight of the crucible + cellulose + mineral matter (g)

P3: weight of the crucible + mineral matter (g)

2.2.6 Carbohydrate levels

The rate of total carbohydrates is obtained from the following formula:

$$C(\%) = 100 - (\text{moisture} + \text{fat} + \text{protein} + \text{ash} + \text{fiber}) [9].$$

2.2.7 Dosage of mineral elements

The mineral elements (Ca, Na, K, Mg, Zn, Cu and Fe) were assayed from the ashes obtained by the incineration of flours from ground tubers. For each sample the contents of mineral elements are determined. In practice, the samples are mineralized, the extracts are prepared and the reading is made by atomic absorption and the calculation of the different values using a calibration curve.

2.2.7.1 Digestion and preparation of sample extracts

A quantity of 1 g of sample is weighed in a porcelain crucible then placed in the oven at 500° C for 3 h. After cooling, 10ml of 1 molar nitric acid is added to the ash obtained and then evaporated completely on a hot plate at 100°C. To the residue are added 5 ml of 0.1 molar hydrochloric acid. Evaporate to dryness on a hot plate, then study at 105°C for about an hour to insolubilize the silica. Take up with 1 ml of HCl at ½ filter, the crucible is rinsed twice with 10 ml of hydrochloric acid. The filtrate is brought to 50 ml in a volumetric flask [10, 11].

III. RESULTS

3.1 Macronutrient, dry matter and moisture content of the tubers of the ecotypes of the tiger nut studied

The results of the table 2 show that the moisture (M) content of tiger nut tubers is relatively low, around 6%. The ash content (mineral matter) is on average 1.7% for the big tiger nut and 1.8% for the small tiger nut. The classification according to ecotypes shows that the ash content of ecotype E2 is highest (3%) and that ecotype E4 recorded the lowest ash content (0.2%).

The fat content is higher in all the ecotypes of the small tiger nut with an average of 21.6% than in the ecotypes of the large tiger nut whose average is 14.9%. The E11 and E9 ecotypes have a higher fat content (25.2%) compared to the other ecotypes. The low-fat content is recorded by ecotype E17 (12.0%).

The average protein (P) content is 4.6% for the big sedge ecotypes and 5.5% for the small shoveler ecotypes. The E7 ecotype has a higher protein content (11.9%). For both types of tiger nut, the lowest protein content was recorded for ecotype E3 (3.1%).

The tiger nut is rich in dietary fiber (F) with an average of 29.5% for the big tiger nut and 28.5% for the small tiger nut. The highest fiber contents were recorded for ecotypes E15 (37%) and E11 (39.4%). Ecotype E7 (small wild tigernut from RijiaSamna) is less rich in fiber (19.8%) compared to the other ecotypes (small and large tiger nuts). However, tiger nut ecotypes have carbohydrate (C) levels ranging from 24.5% to 49.6%. The classification by region of the macronutrient compositions shows that the E11, E9 and E18 ecotypes of the Maradi region are respectively richer in fiber and fat (25.20% and 39.40%), fat (25.20%) and carbohydrate (49.60%). And the wild ecotype E7 from the Dosso region is richer in moisture and protein (7.00% and 11.9%).

Table 2: Macronutrient, dry matter and humidity contents of the tubers of Small and Big tigernut ecotypes (%)

Kinds	Ecotypes	M	DM	MM	OM	FAT	P	F	C
Small tiger nut*	E2	6.40	93.56	3.00	90.56	20.60	6.4	28.80	34.80
Small tigernut**	E3	6.20	93.82	1.90	91.92	22.60	3.1	29.90	36.30
Small Black Tigernut*	E4	6.30	93.70	0.20	93.50	18.50	4.5	27.60	42.90
Small Black Tigernut	E6	6.00	94.00	1.10	92.90	23.70	2.9	27.70	38.60
Small Wild Tigernut	E7	7.00	92.98	2.30	90.68	17.10	11.9	19.80	41.90
small tiger nut	E8	6.50	93.50	1.50	92.00	18.80	8.6	32.50	32.10
small tiger nut	E9	6.00	93.98	2.00	91.98	25.20	5.4	24.60	36.80
small tiger nut	E10	6.40	93.56	1.20	92.36	23.20	9.3	25.20	34.70
small tiger nut	E11	6.10	93.92	1.20	92.72	25.20	3.6	39.40	24.50
Small Wild Tigernut***	E5	6.70	93.34	2.90	90.44	18.50	4.4	25.50	42.00
small tiger nut	E12	5.50	94.48	1.70	92.78	23.60	3.4	30.00	35.80
small tiger nut	E13	5.80	94.24	1.80	92.44	22.50	4.8	29.90	35.20
small tiger nut	E14	5.30	94.66	1.90	92.76	21.10	3.2	28.90	39.60
Mean of smalls tiger nuts		6.20	93.80	1.80	92.00	21.60	5.5	28.50	36.40
Big tigernut	E15	6.50	93.50	0.80	92.70	14.30	4.6	37.00	36.80
Big tigernut	E1	7.00	93.00	1.90	91.10	16.50	4.6	27.40	42.60
Big tigernut	E16	6.90	93.10	2.60	90.50	15.80	5.1	21.90	47.70
Big tigernut	E17	6.50	93.50	0.50	93.00	12.00	4.1	30.50	46.40
Big tigernut	E18	5.80	94.20	2.50	91.70	13.90	4.1	24.10	49.60
Big tigernut	E19	6.00	94.00	1.80	92.20	16.00	6	33.00	37.20
Big tigernut	E20	5.80	94.20	1.90	92.30	15.80	3.4	32.70	40.40
Mean of big tiger nuts		6.30	93.70	1.70	92.00	14.90	4.6	29.50	43.00

Legend: F: fibers, MM: mineral matter, fat (lipids), P: protein, OM: organic matter, C: carbohydrates, M: Moisture, DM: dry matter

3.2 Mineral element content of the tubers of the tiger nut ecotypes studied

The composition of the tiger nut in mineral elements (Table 3) is very variable at the level of the two types of ecotypes (small and large tiger nut) but also within the same types. The results of Table 3 show that most of the mineral elements have higher contents for the ecotypes of the small tiger nut than for the ecotypes of the large tiger nut.

The Copper (Cu), Sodium (Na), Calcium (Ca) and Magnesium (Mg) contents are significantly identical for all the ecotypes studied. The Zn contents of the E2 (2.4361±0.7261a mg/100g) and E3 (2.4349±0.6123amg/100g) ecotypes of tiger nut are significantly higher compared to the ecotypes E19 and E20. The Zn contents of the ecotypes of the big tiger nut E19 (0.6757±0.4510bmg/100g) and E20 (0.8386±0.3169bmg/100g) are significantly lower. The iron content of ecotype E7 (15.809±6.524 a mg/100g) is highly significant compared to the ecotypes studied. However, the E1, E2, E3, E4, E5, E8, E9, E11, E12 and E15 ecotypes are distinguished with respectively relatively high iron contents of : 5.969 ± 3.112; 11.406±5.780ab; 10.877±5.134; 10.961±5.612; 5.338±5.159; 2.982±1.758; 4.029±2.469; 3.157±2.108; 5.063±2.600 and 3.139±2.134mg/100g. Potassium content is statistically significant for ecotypes E2 (308.41±8.90amg/100g), E3 (288.24±8.32abmg/100g) and E15 (300.25±8.67amg/100g). the classification by region of the composition of mineral elements shows that the ecotypes E4, E2 of the Maradi region are respectively richer in sodium (Na)(8,157±3,418) and potassium (K)(308,410±8,900). And the wild E7 ecotype and the E3 ecotype of the Dosso region are richer in iron (Fe) (15,809±6,524) and zinc (Zn)(2,435±0,612)

Table 3: Contents of mineral elements in the tubers of the tiger nut ecotypes (mg/100g)

Kinds	Ecotypes	Cu	Na	Ca	Zn	Fe	K	Mg
Big tiger nut	E1	0.233±0.030a	4,909±0,748a	28,850±15,400a	1,211±0,465ab	5,969±3,112ab	112,890±3,260ij	372,150±3,707a
Small tiger nut*	E2	0.502±0.061a	3,776±1,282a	65,380±19,940a	2,436±0,726a	11,406±5,780ab	308,410±8,900a	411,730±7,338a
Small tiger nut*	E3	0.397±0.036a	5,514±0,748a	41,670±13,600a	2,435±0,612a	10,877±5,134ab	288,240±8,320ab	456,490±6,052a
Small Black Tiger nut*	E4	0.242±0.028a	8,157±3,418a	42,310±25,380a	1,993±0,436ab	10,961±5,612ab	112,890±3,260ij	406,210±1,967a
Small Black Tiger nut	E5	0.378±0.085a	5,363±0,320a	20,510±10,880a	1,171±0,492ab	5,338±5,159ab	243,560±7,030def	408,700±2,724a
Small Wild Tiger nut	E6	0.120±0.026a	5,211±0,107a	34,620±19,940a	1,029±0,307ab	2,451±1,314b	154,690±4,460gh	439,550±3,404a
small tiger nut	E7	0.311±0.081a	5,211±0,320a	52,560±38,070a	1,424±0,317ab	15,809±6,524a	102,800±2,970d	471,830±8,927a
small tiger nut	E8	0.258±0.036a	5,363±0,320a	43,590±16,320a	1,363±0,316ab	2,982±1,758ab	259,890±7,500cd	433,130±4,312a
small tiger nut	E9	0.178±0.036a	4,683±0,427a	24,360±10,880a	1,146±0,219ab	4,029±2,469ab	232,030±6,700ef	387,480±4,363a
small tiger nut	E10	0.153±0.024a	4,909±0,320a	21,150±20,850a	1,927±0,482ab	0,881±0,017b	162,370±4,690g	370,360±6,985a
Small Wild Tiger nut**	E11	0.482±0.502a	5,211±0,320a	29,490±14,500a	1,111±0,365ab	3,157±2,108ab	251,250±7,250cde	382,490±4,665a
small tiger nut	E12	0.266±0.036a	6,344±1,068a	20,510±7,250a	1,326±0,474ab	5,063±2,600ab	224,820±6,490f	366,800±1,942a
small tiger nut	E13	0.25±	6,495±	19,230±7,250a	1,299±0,407ab	2,399±	273,820±7,900bc	438,480±3,556a

rnut		0.041a	1,495a			1,189b		
smalltiger nut	E14	0.499± 0.530a	5,136± 0,214a	33,970± 17,220a	1,025± 0,399ab	2,365± 1,038b	222,900± 6,430f	443,120± 4,413a
Big tiger nut	E15	0.541± 0.574a	4,305± 1,175a	28,21± 19,940a	0,9392± 0,4733ab	3,139± 2,134ab	300,25± 8,670a	440,44± 3,278a
Big tiger nut	E16	0.235± 0.069a	5,060± 0,320a	14,10± 5,440a	0,8686± 0,3033ab	2,560± 1,160b	161,89± 4,670g	449,18± 7,086a
Big tiger nut	E17	0.400± 0.077a	6,042± 1,068a	16,67± 5,440a	0,7732± 0,3226ab	1,085± 0,458b	168,14± 4,850g	399,79± 3,984a
Big tiger nut	E18	0.159± 0.071a	5,211± 0,107a	16,03± 4,530a	0,6618± 0,2491ab	2,133± 0,967b	177,26± 5,120g	470,22± 9,154a
Big tiger nut	E19	0.004± 0.000a	7,175± 1,389a	15,38± 5,440a	0,6757± 0,4510b	1,932± 1,451b	134,51± 3,880hi	413,87± 1,992a
Big tiger nut	E20	0.004± 0.004a	6,722± 1,816a	23,08± 5,440a	0,8386± 0,3169b	1,807± 1,428b	242,60± 7,00def	454,89± 6,279a

Values are means (n=2) ±SD. Values with the same letter in the same column are not significantly different ($p < 0.05$)

VI. DISCUSSION

Tiger nut tubers, whether small or large, have an average moisture (M) rate of about 6%. The results support those of [12, 13, 14] in their respective studies on the physico-chemical characterization of the three morphotypes of tiger nuts with humidity levels of $5.190 \pm 0.180\%$; $4.560 \pm 0.220\%$, $4.990 \pm 0.780\%$ and $8,300 \pm 0,100 \text{ à } 8,600 \pm 0,170\%$. Also, on the characterization of four tiger nut ecotypes with $8.660 \pm 0.040\%$, $7.750 \pm 0.270\%$, $6.380 \pm 0.450\%$ and $7.450 \pm 0.310\%$. Higher humidity levels were recorded by [15] (9.23%) and by [16] (9.73%). The variation in moisture content could depend on the degree of drying after harvest.

The average fat contents obtained are 14.90% for the ecotypes of the big tiger nut and 21.60% for the ecotypes of the small tiger nut. These results are corroborated by those of [17] who obtained a fat content in tiger nut between $18.23 \pm 0.01\%$ and $22.15 \pm 0.03\%$ depending on the “varieties” studied. However, the E11 and E9 ecotypes have much higher fat contents (25.2%) but close to the contents obtained by the [18] (25.6%); [19] (25.40%); [20], (24.45%) and [12] (26.14%). Moreover, these contents are lower than those of [21] who obtained contents ranging from $26.88 \pm 3.32\%$ to $44.92 \pm 0.76b\%$; from [22] on the yellow and brown varieties whose respective contents are 32.13 and 35.43% and from [23], which is 32.8%. The tubers of the ecotypes of small tiger nut are richer in fat and would therefore be interesting for oil extraction [30]

However, the fat contents of the ecotypes of the tiger nut analyzed remain higher than those contained in certain tubers such as cassava (0.1 to 0.8%) [24] cited by [15], taro (0.33 to 1.17%) [25] and sweet potato (0.9%) [26]. Cereals such as millet and sorghum also contain respectively lower contents (5.14 to 5.96%) [27] and 2.40% to 10.54% [28,29].

The protein contents obtained vary from 2.90% to 11.90% for the ecotypes of two types of tiger nut (large tiger nut and small tiger nut) studied. These results show a fairly good protein profile. Also, these contents are consistent with those of [13] and [30] with protein contents ranging from $3.28 \pm 0.10\%$ to $8.45 \pm 0.20\%$ and $3,43 \pm 0,05$ to $8,35 \pm 0,22$; [31] (5.04%); that of the [18] which is $4.8 \pm 1.1\%$; those of Nigerian varieties which vary from 7.15 to 10.50% [22,

19] and those of the samples from Côte d'Ivoire studied by [3]. This observed variability in protein content may be due to climatic and soil conditions [15].

The average dietary fiber contents of the ecotypes are 29.50% and 28.50% for those of the big tiger nut and the small tiger nut respectively. These results are close to those of [20, 32]. Our results are higher than the fiber content obtained by [18] (11.7%) and very different from those of [17] and [31] who obtained grades ranging from $5.40 \pm 0.01\%$ to $8.63 \pm 0.03\%$ and 8.91%. Indeed, the ecotypes of Niger tiger nut contain a higher fiber content than yam and cassava, which contain 3.14% and 3.66% respectively [15]. On the other hand, this content is much higher than those of cereal flours such as wheat, millet, sorghum and maize, the contents of which vary respectively between 1.5 and 2%; 1.5 to 2.3% and 1.5 to 1.8% [15, 27, 28]. The difference in the levels observed can be explained by the fact that the present study considered whole tubers (skin with flesh) whereas in the studies mentioned, the authors only used tuber flour. These fiber contents of the ecotypes studied give tiger nut tubers interesting properties in terms of digestion [15].

Tigernut tubers contain a high rate of carbohydrate content with an average of 43.00% for large tigernut ecotypes and 36.40% for small tigernut ecotypes. These contents are similar to those of [21] which are between $52.2 \pm 0.95\%$ and $42.14 \pm 0.04\%$ and are higher than that obtained by [33] which is $19.40 \pm 0.11\%$.

The mineral element composition of the studied ecotypes is significantly similar for copper (Cu), sodium (Na), calcium (Ca) and magnesium (Mg). However, they differ for Zinc (Zn), Iron (Fe) and Potassium (K). The Cu, Ca, Zn, Fe and K contents obtained for most of the ecotypes studied are consistent with those obtained by the [18]. These results are consistent with those of [12] with Cu (0.430 ± 0.01 to 0.71 ± 0.03), Ca (22.13 ± 1.64 to 32.27 ± 5.66), Zn (1.88 ± 0.22 to 2.7 ± 0.03), and in Fe ($3.570.17 \pm 11.44 \pm 0.48$). The Na contents (3.776 ± 1.282 to 7.175 ± 1.389) obtained for all the ecotypes are lower than the content obtained by the [18] but with higher Mg contents (366.80 to 471.83 ± 8.927) than those obtained by [18, 15] ($102, 104.9 \pm 0.67$). The Mg ($895,79 \pm 41,94$ to $1344,98 \pm 69,30$) and K ($418,73 \pm 41,46$ to $944,60 \pm 48,57$) contents obtained by [32] in their study on 9 ecotypes of tigernut of Burkina Faso are superior. The differences in the contents of mineral elements observed could be linked to the composition of the soil which can influence the absorption and storage of these elements in the tubers [12]. Ecotypes E1, E2, E3, E4, E5, E7, E8, E9, E11, E12 and E15 differ from other ecotypes due to their high Zn, Fe and K contents.

The tiger nut tubers of certain ecotypes studied are richer in iron than other foods, in particular the leaves of *Moringa oleifera* ($3.81 \pm 0.12\text{mg}/100\text{g}$), *vigna unguiculate L.* ($7.80 \pm 0.15\text{cmg}/100\text{g}$) and the leaves of *Hubicus sabdariffa* ($8.27 \pm 1.01\text{bmg}/100\text{g}$) mentioned by [34]. The composition of mineral elements such as copper, iron, zinc, calcium, and the fiber and protein compositions of the potato obtained by [18] are lower than that of the tubers of the studied tiger nut ecotypes.

V.CONCLUSION

The chemical composition of the tubers of the studied nutsedge ecotypes is rich and very varied. The averages of the fat, protein and iron contents obtained are higher for the ecotypes of the small tiger nut than for the ecotypes of the large tiger nut from Niger. The high fat content of cultivated and wild small tiger nut ecotypes may be favorable for oil extraction. The E1, E2, E3, E4, E5, E7, E8, E9, E11, E12 and E15 ecotypes are distinguished from the

other ecotypes by their high iron content. these ecotypes are richer in iron than the leaves of *Moringa oleifera* ($3.81 \pm 0.12\text{mg}/100\text{g}$), the pods of *Vigna unguiculata* ($7.80 \pm 0.15\text{cmg}/100\text{g}$) and the leaves of *Hibiscus sabdariffa* ($8.27 \pm 1.01\text{bm}/100\text{g}$). the study showed that the compositions of mineral elements such as copper, iron, zinc, calcium and the fiber and protein compositions of tiger nut are superior to those of apple earthen. this study resulted in a better nutritional profile of Niger tiger nut tubers.

Declarations

Competing interest

The authors have no relevant financial or non-financial interests to disclose. The authors have no competing interests to declare that are relevant to the content of this article. All authors certify that they have no affiliations with or involvement in any organization or entity with any financial interest or non-financial interest in the subject matter or materials discussed in this manuscript. The authors have no financial or proprietary interests in any material discussed in this article.

REFERENCES

1. Bamishaiye EI, Bamishaiye OM. Tigernut. As a plant, its derivatives and benefits. *Afri. J. Food, Agric. Nutri. Dev.* (2011). 11(5):5157,5170. <https://www.ajol.info/index.php/ajfand/article/view/70443> consulted 05/09/2022
2. Bori H, Toudou D A K, Moumouni DMA, Adam T. Cultivation of tiger nuts (*Cyperus esculentus*) in Niger : Origin, assets and constraints *Journal of Animal & Plant Sciences.* (2018). Vol.37, Issue 1: 5997-6007: 11. Français.
3. Ban-Koffi G, Nemlin S, Le Fevre, K A. "Physicochemical characterization and therapeutic potential of sweet pea (*Cyperus esculentus* L. _ (Cyperaceae))", *Agron . Afr.* (2005). vol. 17, No. 1: 63-71: DOI:10.4314/aga.v17i1.1658. Français.
4. Development Initiatives (DI). *Global Nutrition Report. Nutrition at the service of the SDGs.* Bristol UK. (2017). 115. Français.
5. Houndji BV, Romaric O, Serge B L, Kou SSE, Boniface Y, Cohovi BA. Microbiological and physico-chemical characterizations of *Moringa* leaf powder *oleifera* (Lam.), a traditional leaf vegetable in Benin. *Int. J. Biol. Chem. Sci.* (2013). 75-85:11.
6. Materne M. Des chiffres et des faits sur la malnutrition dans le monde. *FAO.* (2018).8. Français.
7. INS. National Institute of Statistics, Niger. Final report of the Assessment of the nutritional situation by the SMART methodology in Niger. (2016). 104. Français.
8. AOAC. *Official Methods of Analysis. Association of Official Analytical Chemists.* 14th édition, Washington (USA). (1990). 150.
9. Adamou OB. Feeding practices and microbiological and nutritional quality of food for children aged 6 to 24 months suffering from moderate acute malnutrition in the Mayahi department in Niger. Doctoral thesis from Abdou Moumouni University of Niamey. (2019). 132. Français.
10. Tahri M, Benya`Ich F, Bounakhla M, Bilal E, Gruffat JJ, Moutte J, Garcia D. Multivariate analysis of heavy metal contents in soils, sediments and water in the region of Meknes (central Morocco). *Environmental Monitoring and Assessment.* 2005. 13.
11. Didier. Assay of major mineral elements in plants: Methods used in the ORSTOM foliar diagnostic laboratory. 1966. 56. Français.

12. Bado S, Patrice B, Gouyahali S, Moe T, Brian P, Stephan N, et al. Physicochemical Characteristics and Composition of Three Morphotypes of *Cyperus esculentus* Tubers and Tuber Oils. *Journal of Analytical Methods in Chemistry*. 2015.9.Français.
13. Codina-Torrella I, Buenaventura G, Antonio JT. Characterization and comparison of tiger nuts (*Cyperus esculentus* L.) from different geographical origin. Physico -chemical characteristics and protein fractionation. *Ind. Crops Prod*.2015.<http://dx.doi.org/10.1016/j.indcrop.2014.11.007>
14. Yapi J C, Deffan Z A B, Koko A C, Diabagate J R, Kouame Kan K B, Kouame L P. Influence de la granulométrie sur les caractéristiques physicochimiques et techno fonctionnelles des farines de souchet (*Cyperus esculentus* L.). *Agronomie Africaine Sp*. 2021.33 (2): 239 – 250.Français.
15. NDIAYE B. Contribution to the development of tiger nut tubers (*Cyperus esculentus*): nutritional intake, technological routes and galactogenic properties (Doctoral dissertation, Cheikh Anta University Diop).2021.Français.
16. Monago C, Uwakwe C. “Proximate composition and in vitro anti-sickling property of Nigeria *Cyperus esculentus* (tiger nut sedge)”. *Trees Life J*. 2009. vol. 4, no. 2: 1–6.
17. Chinma, CE, Olufemi A, Joseph OA .Effect of germination on the chemical, functional and pasting properties of flour from brown and yellow varieties of tigernut(*Cyperus esculentus*). *Food Research International*. 2009. 42:1004–1009.
18. FAO. Food Composition Table for Western Africa (2019) User Guide & Condensed Food Composition Table. 2020.556p.
19. Adebayo S, and Arinola S. “Effect of Germination on the Nutrient and Antioxidant Properties of Tigernut (*Cyperus esculentus*) » *Journal of Biology, Agriculture and Healthcare*. 2017. Vol.7, n°1888-94.
20. Okoye, Ene. “Effects of processing on the nutrient and anti-nutrient contents of tiger nut (*Cyperus esculentus*)”. *J Food Tech Food Chem*. 2018, vol. 1, no. 1: 1-7.
21. Richard A, Djomdi, Robert N. Characteristics of tigernut (*Cyperus esculentus*) Tubers and their performance in the production of a milky drink. *Journal of Food Processing and Preservation*.2006. 20: 145–163.
22. Oladele A K, Aina OJ. “Chemical composition and functional properties of flour produced from two varieties of tigernut (*Cyperus esculentus*)”, *Afr. J. Biotechnol*.2007. vol. 6, no. 21: 2473-2476.
23. El-Naggar EA. Physicochemical Characteristics of Tiger Nut Tuber (*Cyperus esculentus* Lam) Oil. *Middle East Journal of Applied Sciences*. 2016. 2077-4613: 1003-1011.8
24. Charles AL, Ssiroth K, Huang T. "Proximate composition, mineral contents, hydrogen cyanide and phytic acid of 5 cassava genotypes" *Food Chem*. 2005. vol. 92, No. 4: 615-620.
25. Aboubakar. "Optimization of the production and conservation parameters of taro flour (*Colocasia esculenta*)” Nancy-university, national polytechnic institute of Lorraine / university of Ngaoundere. 2009. 211.Français.
26. NdanguiChancelle B. Production and characterization of sweet potato flour (*Ipomoea batatas*.Lam): optimization of bread-making technology. *Human medicine and pathology*. University of Lorraine. 2015
27. Kulthe AA, Thorat SS, Lande SB. "Characterization of Pearl Millet Cultivars for Proximate Composition, Minerals and Anti-Nutritional Contents", *Adv. Life Sci*. 2016. vol. 5, no. 11, 4672–4675.
28. Mohammed NA, Ahmad IAM, Babiker EE. “Nutritional evaluation of sorghum flour (*Sorghum bicolor* L. Moench) during processing of injera”, *World Acad. Science. Eng. Technol*.2011. vol. 51,72-76.
29. Jimoh W, Abdullahi M. “Proximate analysis of selected sorghum cultivars”, *Bayero J. Pure Appl. Science*. 2017. vol. 10, No. 1. 285-288.

30. Semdé Z, Parkouda C, Tamboura D, Sawadogo L H, Semdé Z, Parkouda C, et al. Variability of physical and nutritional characteristics of tiger nut (*Cyperus esculentus* L.) tubers grown in Burkina Faso. *Science and technology, Natural and applied sciences*. 2019. Vol. 38: 21-33
31. Maduka, I. Tigernut Plant and Useful Application of Tigernut Tubers (*Cyperus esculentus*). *Current Journal of Applied Science and Technology* Article no.2018. CJUST.43551. ISSN: 2457-1024. 24.
32. Ekeanyanwu, Ononogbu C . “Nutritive value of Nigerian tigernut (*Cyperus esculentus*)” *Agric. J.*2010. vol. 5: 297-302.
33. Aka BA, Konan N'Guessan Y, Coulibaly A, Biegho GH. Nutritional and Sensory Analysis of Milk Processed from Seeds of Sweet Pea (*Cyperus esculentus* L.) Consumed in Ivory Coast. *JALSI*. 2016. 8(2): 1-12(12). Français.
34. Tchiégang C, Kitikil A. Ethno-nutritional data and physico-chemical characteristics of leafy vegetables consumed in the Adamaoua savannah (Cameroon). *TROPICULTURA*. 2004. 22, 1, 11-18pp. Français.