

Original Research Article

Phytochemical profile and in vitro antioxidant capacity of *Petiveria alliacea* plant parts extract

ABSTRACT

Plants have significantly contributed to the discovery of modern drugs that have therapeutic potential. This study was carried out in order to evaluate the phytochemical constituents and in vitro antioxidant potential of the leaf and root of *Petiveria alliacea* extracts. The aqueous and ethanolic leaf and root extracts of *Petiveria alliacea* were screened for qualitative and quantitative analysis. Different concentrations of the aqueous and ethanolic extracts were scanned within the wavelength range of 378–654 nm. The in vitro antioxidant (catalase and hydrogen peroxide scavenging) potential of the extracts was determined spectrophotometrically. The study revealed the presence of tannins, phlobatannins, saponins, flavonoids, terpenoids, and cardiac glycosides in aqueous and ethanolic extracts of the leaves and root. High levels of total alkaloids and flavonoids were observed in the ethanolic and aqueous leaf extracts. The ultraviolet-visible profile of the extracts revealed high peaks at 378 nm for the leaf and root at different concentrations, confirming the presence of flavonoids and alkaloids in the plant. The high hydrogen peroxide scavenging activity was observed in aqueous extracts of the leaf and root and in the ethanolic leaf extract but not in the ethanolic root extract. The phosphate buffer leaf extract, as well as the soaked ethanolic and aqueous leaf extracts, had the highest catalase activity, according to the study. These findings suggest that both the leaf and root of *Petiveria alliacea* may be good sources of natural antioxidants and might be useful in treating diseases associated with oxidative stress. patients. These predictors, however, need further work to validate reliability.

Keywords: *Petiveria alliacea*, catalase, antioxidants, phytochemical

1. INTRODUCTION

Reactive oxygen species (ROS) include free radicals such as superoxide anion radicals (O_2^-), hydroxyl radicals (OH) and non-free radical species such as hydrogen peroxide (H_2O_2) and singlet oxygen, that can damage cell structures such as carbohydrates, nucleic acids, lipids, and proteins and alter their functions [1-3]. Moreover, the antioxidant systems can be overwhelmed by oxidative stress that contributes to many pathological conditions, including cancer, neurological disorders, atherosclerosis, hypertension, ischemia/perfusion, diabetes, acute respiratory distress syndrome, idiopathic pulmonary fibrosis, chronic obstructive pulmonary disease, and asthma [4-6].

Harmful ROS in animals and humans can be scavenged by phytochemical compounds, such as alkaloids and flavonoids which are antioxidants that have potential health effects [7,8].

For decades, modern drugs have been isolated from medicinal plants, and recently much attention has been dedicated to antioxidants and their association with health benefits [9,10]. The presence of phytochemical constituents in plants is necessary for their biological activities [11] and these can be derived from the different parts of the plant like the leaf, stem, seed, fruit, and root that work with the nutrients and fibers to act as a defense system or to protect against disease [12].

Petiveria alliacea, a Phytolaccaceae family medicinal plant, is a perennial shrub that had traditionally been used as analgesic, anti-inflammatory, anti-microbial, anticancer, anti-rheumatic, antispasmodic, diuretic, sedative, abortive, and hypoglycaemic agents [13-16]. To establish its medicinal effect, the study was designed to assess the phytochemical profiles and evaluate the in vitro antioxidant capacities of aqueous and ethanolic leaf and root extracts of *Petiveria alliacea*.

2. METHODOLOGY

2.1 Collection of plant materials and extraction procedures

Petiveria alliacea leaves and roots were collected from a farm at Ado-Ota Local Government Area, Ogun State, Nigeria, and authenticated in the Department of Botany, Lagos State University, Ojo, Nigeria. The leaf and roots were cleaned in running water to remove impurities and air-dried under ambient conditions ($30 \pm 2^{\circ}\text{C}$) for seven (7) days. The dried samples were ground into a fine powder in a pulveriser then stored in an airtight container at room temperature. The aqueous and ethanolic leaf and root extraction was carried out according to the modified methods of Ogunrinola *et al.* [17]. Briefly, 200 g of *Petiveria alliacea* leaves and roots were boiled with 500 mL of water and ethanol, respectively, for 10 minutes. The resulting solutions were then soaked for 48 hours in 500 mL of water and ethanol, respectively. The obtained extracts were filtered over Whatman No. 1 paper, and the filtrate was concentrated by a rotary evaporator at 50°C and kept for further analysis.

2.2 Phytochemical profile

2.2.1 Qualitative analysis for phytochemical constituents

The following phytoconstituents: tannins phlobatannins saponins flavonoids terpenoids, cardiac glycosides, phenolics, alkaloids in the aqueous and ethanolic extracts of *Petiveria alliacea* leaves and roots respectively using modified standard procedures described by Kalaichelvi and Dhivya [18], Edeoga *et al.*, [19], Ayoola *et al.*, [20], Trease and Evans [21], Ogunrinola *et al.*, [17], and Harborne, [22].

2.2.2 Quantitative analysis for phytochemical constituents

The amounts of alkaloids and flavonoids present in the crude powdered samples of *Petiveria alliacea* leaves and roots were evaluated using standard procedures based on the modified methods of Ayodele *et al.* [23], Boham *et al.* [24], and Harborne [22].

2.3 Ultraviolet-visible Spectroscopic analysis

The modified method of Karpagasundari and Kulothungan [25] was used to identify the chemical constituents of aqueous and ethanolic *Petiveria alliacea* leaf and root extracts. Briefly, the soaked and unsoaked extracts were centrifuged at 3000 rpm for 10 minutes and filtered through Whatman No. 1 filter paper. The sample was diluted to 1:10 with the same solvent. Different concentrations of the extracts were mixed with 50 mM Tris-HCl buffer (pH 7.0), and scanned at a wavelength ranging from 378-654 nm using Thermo Scientific Spectroscan Ultraviolet-Visible (UV) 2700 double beam Spectrophotometer, and the characteristic peaks were detected. The UV peak values were recorded. For spectrum confirmation, each and every analysis was repeated twice.

2.4 Evaluation of In Vitro Antioxidant

2.4.1 Hydrogen peroxide scavenging effects

The scavenging activity of *Petivera alliacea* aqueous and ethanolic extracts of leaves and roots was evaluated by the modified method of Ruch *et al.* [26]. Briefly, a solution of hydrogen peroxide (40 mM) was prepared in phosphate buffer (0.1M, pH 7.4). Concentrations of *Petiveria alliacea* leaf and root extracts ranging from 0.1 - 0.6 mg/ml were added to the hydrogen peroxide buffer solution, mixed, and absorbance at 230 nm against the phosphate buffer as blank after 10 minutes. The percentage of hydrogen peroxide (H₂O₂) inhibition was calculated as:

% Scavenging of H₂O₂ = [(AC - AS)/AC] x 100; where AC was the absorbance of the control, and AS was the absorbance of the extracts and standard.

2.4.2 Evaluation of catalase activity

The modified methods of Aebi [27] and Luck [28] were employed to assay for catalase activity of *Petiveria alliacea* leaf and root extracts. In brief, a 20% homogenate of *Petiveria alliacea* leaf and root extracts was prepared in phosphate buffer (0.067M, pH 7.0). 2 ml of homogenate was mixed with 1 ml of hydrogen peroxide solution (2mM) in a quart cuvette. Also, 1 ml of soaked (for 24 hours), and unsoaked aqueous and ethanolic extracts were mixed with of hydrogen peroxide. Catalase activity was assayed by measuring hydrogen peroxide consumption at 240 nm for 180 seconds with a 30 second interval, as indicated by a decrease in absorbance. The control and blank were assay solution without the extract. Enzyme activity was expressed as catalase unit/g protein.

3. RESULTS AND DISCUSSION

3.1 RESULTS

3.1.1 Qualitative analysis of ethanolic and aqueous leaf and root *Petivera alliacea* extracts

Table 1 shows the phytochemical constituents of the ethanolic and aqueous *Petivera alliacea* leaf and root extracts. Alkaloids, terpenoids, tannins, saponins, phlobatannins, flavonoids, and cardiac glycosides were present in both extracts.

Table 1. Qualitative analysis of ethanolic and aqueous leaf and root *Petivera alliacea* extracts

Phytochemical constituents	Ethanolic Extract		Aqueous Extract	
	Leaves	Root	Leaves	Root
Tannins	+ve	+ve	+ve	+ve
Phlobatannins	+ve	+ve	+ve	+ve
Saponins	+ve	+ve	+ve	+ve
Flavonoids	+ve	+ve	+ve	+ve
Alkaloids	+ve	+ve	+ve	+ve
Terpenoids	+ve	+ve	+ve	+ve

Cardiac Glycosides	+ve	+ve	+ve	+ve
--------------------	-----	-----	-----	-----

+ve = positive

3.1.2 Quantitative analysis of ethanolic and aqueous leaf and root *Petivera alliacea* extracts

The quantitative estimation of ethanolic and aqueous leaf and root *Petivera alliacea* extracts is depicted in Table 2. While the ethanolic extract of *Petivera alliacea* had a high level of alkaloids (38.3 w/w) in the leaves and flavonoids in the root (17.7 w/w), the aqueous leaves had a high level of flavonoids (28.3 w/w), but a low level of alkaloids and flavonoids in the root extracts.

Table 2. Quantitative analysis of ethanolic and aqueous leaf and root *Petivera alliacea* extracts

phytochemical constituents	Ethanolic Extract		Aqueous Extract	
	Leaf (w/w)	Root (w/w)	Leaf (w/w)	Root (w/w)
Alkaloids	38.3	2.1	6.8	0.5
Flavonoids	5.3	17.7	28.3	0.3

3.1.3 Ultraviolet-visible Spectroscopic of ethanolic and aqueous leaf and root *Petivera alliacea* extracts

The qualitative UV spectrophotometric profiles of different concentrations of aqueous and ethanolic extracts of *Petiveria alliacea* leaves and roots are shown in Figures 1-5. The absorption of 0.01 g/ml of aqueous and ethanolic extracts of *Petiveria alliacea* at different wavelengths is depicted in Figures 1a and 1b. The leaves and root aqueous extracts showed a high absorptivity at 378 nm which further decreased as the wavelength increased. While the root ethanolic extract showed the same pattern, the leaf ethanolic extract had increased absorptivity at 378 nm, which decreased at a wavelength of 608 nm and increased again at a wavelength of 654 nm.




Figure 1. UV-Visible spectrophotometric profile of 0.01g/ml aqueous (a) and ethanolic (b) extracts of *Petiveria alliacea* leaves and root

The UV absorption spectrophotometric profile of 0.025 g/ml aqueous and ethanolic extracts of *Petiveria alliacea* is shown in Figures 2a and 2b. All the extracts showed the same pattern of decreasing intensity as the wavelength increased.




Figure 2. UV-Visible spectrophotometric profile of 0.025g/ml aqueous (a) and ethanolic (b) extracts of *Petiveria alliacea* leaves and root

The UV absorption spectrophotometric profile of 0.075 g/ml aqueous and ethanolic extracts of *Petiveria alliacea* is depicted in Figures 3a and 3b, in which the leaves and root aqueous extracts (Fig. 3a) showed a high absorptivity at wavelength 378 nm, which decreased as the wavelength increased. In Fig. 3b, there was an up-and-down absorptivity of the leaf and root ethanolic extracts.




Figure 3. UV-Visible spectrophotometric profile of 0.075g/ml aqueous (a) and ethanolic (b) extracts of *Petiveria alliacea* leaves and root

The UV absorption spectrophotometric profile of 0.1 g/ml aqueous and ethanolic extracts of *Petiveria alliacea* shown in Figures 4a and 4b revealed a decrease in absorptivity as the wavelength increased for both leaves and roots aqueous extracts (Fig. 4a) and an up-and-down absorptivity of the leaves and roots ethanolic extracts (Fig. 4b).




Figure 4. UV-Visible spectrophotometric profile of 0.1g/ml aqueous (a) and ethanolic (b) extracts of *Petiveria alliacea* leaves and root

Figures 5a and 5b indicated that the absorptivity of the root aqueous extract was higher than that of the leaves, but both extracts followed the pattern of increased absorptivity as wavelength increased (Fig. 5a), while the ethanolic leaf extract (Fig. 5b) showed the highest absorptivity in an up-and-down manner.




Figure 5. UV-Visible spectrophotometric profile of 0.15g/ml aqueous (a) and ethanolic (b) extracts of *Petiveria alliacea* leaves and root

3.1.4 Hydrogen peroxide scavenging effects of ethanolic and aqueous leaf and root *Petiveria alliacea* extracts

As shown in Figure 6, the aqueous extract of *Petiveria alliacea* demonstrated increased hydrogen peroxide decomposition activity of 10.4 % (leaves) and 11.7 %, 11.8 % (root), respectively, in a concentration dependent manner.




Figure 6. The % hydrogen peroxide scavenging activity against different concentrations of aqueous root and leaf *Petiveria alliacea* extracts

Figure 7 shows that the ethanolic leaf extract had a high level of hydrogen peroxide decomposition at a 0.1 g/ml concentration compared to the root extract.




Figure 7. The % hydrogen peroxide scavenging activity against different concentrations of ethanolic root and leaf *Petiveria alliacea* extracts

Phosphate buffered leaf extracts (Fig. 8) exhibited higher catalase activity (0.612 Unit/g) compared to the root extract (0.198 Unit/g). Figure 9 depicts the catalase activity of soaked and unsoaked aqueous and ethanolic extracts of *Petiveria alliacea* leaves and root respectively. There was a high level of catalase activity in the ethanol-soaked leaf and root extracts, representing 0.198 Unit/g and 0.076 Unit/g, respectively, compared to the unsoaked extract. The aqueous soaked leaves extract had a high catalase activity of 0.102

Unit/g compared to the root extract, and the unsoaked aqueous root extract has a high activity (0.07 Unit/g) compared to the leaves extract.




Figure 8. Catalase activity (unit/g) of phosphate buffer *Petiveria alliacea* leaf and root extract

UNDER PEER REVIEW




Figure 9. Catalase activity (unit/g) of soaked and unsoaked aqueous and ethanolic extracts *Petiveria alliacea* leaves and root
 ASRE = aqueous soaked root extract, ESRE = ethanolic soaked root extract, ASLE = aqueous soaked leaf extract, ELSE = ethanolic leaf soaked extract, UALE = unsoaked aqueous leaf extract, UERE = unsoaked ethanolic root extract, UELE = unsoaked ethanolic leaf extract, UARE = unsoaked aqueous root extract

3.2 DISCUSSION

Screening for potential phytochemicals and antioxidant agents in natural plant products is still the main focus of many therapeutic scientists because antioxidants have the capacity to stop the chain reaction of oxidative species and the deleterious health hazards they pose to the body. The pharmacological activities of any medicinal plant are due to the presence of the secondary metabolites: flavonoids, glycosides, triperinoids, alkaloids, tannins, saponins, carbohydrates, glycosides, flavonoids, and steroids [7,8,19]. These metabolites have been reported as scavengers of ROS and capable therapeutic drugs for free radical mediated pathologies, including vascular diseases [29]. The present study shows that these metabolites are present in both the aqueous and ethanolic extracts of *Petiveria alliacea* leaves and roots, respectively. The results obtained are consistent with the findings of Sathiyabalan *et al.* [30] and Ayodele *et al.*[23]. The presence of these secondary metabolites explains the industrial and medicinal importance of the plant. This study observed the presence of high levels of alkaloids and flavonoids in ethanolic leaf, root, and aqueous leaf extracts of *Petiveria alliacea*.

Investigation of medicinal plant composition and activity is very important to promote their therapeutic purposes. The ultraviolet-visible spectroscopy profile of aqueous and ethanolic leaf and root *Petiveria alliacea* extracts of different concentrations revealed the highest peaks in the region from 378 to 470 nm. This is a clear indication of the presence of unsaturated groups and heteroatoms such as sulphur, nitrogen, and oxygen, which confirms the presence of organic chromophores such as phenolic, alkaloid, flavonoids, and glycosides compounds within the *Petiveria alliacea* extracts [31]. This is in accordance with the research of Syed and Johnson [32]; Mamta *et al.* [33]; and Neha and Jyoti [34].

Flavonoid and alkaloid compounds are known to exhibit biological activities including antipyretic, analgesic, anti-inflammatory, anti-arthritic, anti-carcinogenic, immunomodulatory and anti-atherosclerotic properties [35,36]. These activities might be related to their antioxidant activities [7], whereby they act in a structure-dependent manner; scavenge reactive oxygen species, and chelate transition metals, which play vital roles in the initiation of deleterious free radical reactions [37]. Antioxidants can be clarified as reductants, and in-activators of oxidants, whose reducing power may serve as a significant indicator of potential antioxidant activity [38]. Therefore, the in vitro antioxidant potential of different extracts of *Petiveria alliacea* leaves and roots was estimated. Hydrogen peroxide is a non-radical oxidizing specie formed in tissues by oxidative processes, which generate hydroxyl radicals and thus initiate and propagate lipid peroxidation [3]. The scavenging of hydroxyl radicals is a significant antioxidant activity for the protection of living systems from deleterious reactions between hydroxyl radicals and a wide range of molecules in living cells, like sugars, amino acids, lipids, and nucleotides [38]. The present study shows that the percentage hydrogen peroxide scavenging activity of aqueous *Petiveria alliacea* leaf and root extracts but not the ethanolic extracts increased in a concentration-dependent manner. This is consistent with the research of Sowndhararajan and Kang,[38]. This implies that the aqueous extract will be able to scavenge hydroxyl radicals.

Catalase is a vital antioxidant enzyme that converts hydrogen peroxide into water and molecular oxygen using either an iron or manganese as cofactor, thereby protecting the tissues from hydroxyl radicals [39,40]. The study revealed the highest catalase activity in the phosphate buffer of *Petiveria alliacea* leaf extract and in the soaked and unsoaked leaves and root of *Petiveria alliacea* aqueous and ethanolic extracts. The ability of *Petiveria alliacea* extract to suppress oxidative stress in vitro through the antioxidant defense enzymes as observed in this study is similar to the previous reports of Ighodaro *et al.* [41] and Valyova *et al.*[42]. The presence of catalase, a potential scavenger of hydrogen peroxide, prevented the uncontrolled export of toxic species from organelles to the cytosol and competed to remove hydrogen peroxide. The phytoconstituents, especially alkaloids and flavonoids, found in *Petiveria alliacea* may be one of the reasons it can act as an antioxidant by releasing electrons to quaiacol-type peroxidases for scavenging hydrogen peroxide, suppressing the formation of reactive oxygen species by inhibiting some enzymes or chelating trace metals involved in free radical production, and stimulating antioxidant defense [43,44].

4. CONCLUSION

The results obtained in the current study revealed that *Petiveria alliacea* leaves and roots are rich sources of phytoconstituents, which was confirmed by the spectroscopic profile. The antioxidant activity exhibited by the plant parts via free radical scavenging and catalase-reducing power might be attributed to its alkaloid and flavonoid contents. These findings therefore suggest that *Petiveria alliacea* leaves and roots might act as a source of natural antioxidants that could have enormous importance as therapeutic agents. However, additional research will be required to fully determine its bioactivity, and toxicity profile for proper use in the pharmaceutical industries.

REFERENCES

1. Karim M, Islam M, Rahman M, Sultana S, Biswas S, Hosen M J, *et al.* Evaluation of antioxidant, anti-hemolytic, cytotoxic effects and anti-bacterial activity of selected mangrove plants (*Bruguiera gymnorrhiza* and *Heritiera littoralis*) in Bangladesh. *Clinical Phytoscience*. 2020: 6(1), 1-12.

2. Oyinloye BE, Adenowo AF, Kappo AP. Reactive oxygen species, apoptosis, antimicrobial peptides and human inflammatory diseases. *Pharmaceuticals*. 2015; 8(2): 151-175.
3. Bhattacharya S, Reactive oxygen species and cellular defense system, in *Free radicals in human health and disease*. 2015: Springer. 17-29.
4. Rdwan HA, ElAreefy AA. Evaluation of oxidative stress and antioxidative defense systems in Infants and Young Children with Iron-Deficiency Anemia of Sharkia Governorate, Egypt. *Journal of Home Economics*. 2016: 26(3).
5. Adeniran OY. Computational study on the molecular interactions of tramadol with selected antioxidant and detoxification enzymes in *Drosophila melanogaster*. *GSC Biological and Pharmaceutical Sciences*. 2021: 15(3), 281-288.
6. Iqbal Z, Javed M, Gull S, Mahmood MHR, Hai Z. Total phenolic contents of two varieties of *Crocus sativus* and their antioxidant activity. *International Journal of Biosciences*. 2019: 14(3), 128-132.
7. Bajpai VK, Agrawal P. Studies on phytochemicals, antioxidant, free radical scavenging and lipid peroxidation inhibitory effects of *Trachyspermum ammi* seeds. *Indian J Pharm Educ Res*, 2015. 49: 58-65.
8. Swargiary A, Brahma, D. Phytochemical analysis and antioxidant activity of *Hodgsonia heteroclita* (Roxb). *Indian Journal of Pharmaceutical Sciences*. 2017: 79(2), 212-219.
9. Dutra RC, Campos MM, Santos A R, Calixto JB. Medicinal plants in Brazil: Pharmacological studies, drug discovery, challenges and perspectives. *Pharmacological research*. 2016: 112, 4-29.
10. Saranraj P, Sivasakthi S, Deepa M. Phytochemistry of pharmacologically important medicinal plants—a review. *Int. J. Curr. Res. Chem. Pharm. Sci*. 2016: 3(11): 56-66.
11. Hussein RA, El-Anssary AA. Plants secondary metabolites: the key drivers of the pharmacological actions of medicinal plants. *Herbal medicine*. 2019: 1, 13.
12. Tekuri SK, Pasupuleti SK, Konidala KK, Amuru SR, Bassaiahgari P, Pabbaraju N. Phytochemical and pharmacological activities of *Solanum surattense* Burm. f.—A review. *Journal of Applied Pharmaceutical Science*. 2019: 9(3), 126-136.
13. Zavala-Ocampo LM, Aguirre-Hernandez E, Perez-Hernandez N, Rivera G, Marchat LA, Ramirez-Moreno E. Antiamoebic activity of *Petiveria alliacea* leaves and their main component, isoarborinol. *Journal of Microbiology and Biotechnology*. 2017: 27(8), 1401-1408.
14. Luz DA, Pinheiro AM, Silva ML, Monteiro MC, Prediger RD, Maia CSF, *et al*. Ethnobotany, phytochemistry and neuropharmacological effects of *Petiveria alliacea* L. (Phytolaccaceae): A review. *Journal of ethnopharmacology*. 2016: 185, 182-201.
15. Gunawan VA, Soetjipto H, Mustika A. Hypoglycemic and Antioxidant Activity of *Petiveria alliacea* in Diabetic Rat Models. *Biomolecular and Health Science Journal*. 2020: 3(1), 19-23.
16. Garcia-Perez ME, Alfonso-Castillo A, Lores OF, Batista-Duharte A, Lemus-Rodriguez Z. Toxicological evaluation of an aqueous suspension from leaves and stems of *Petiveria alliacea* L. (Phytolaccaceae). *Journal of Ethnopharmacology*. 2018: 211, 29-37.
17. Ogunrinola OO, Odulate JE, Elemo BO. Proximate analysis and pharmacognostical investigation of some medicinal plants: *Naudea latifolia*; *Morinda lucida*; *Alstonia congensis* and *Anchornea cordifolia*. 2004: 3, 190-193.
18. Kalaichelvi K, Dhivya S. Phytochemical screening and antibacterial activity of leaf extract of *Martynia annua*, L. and *Premna latifolia*, Roxb. *Journal of Medicinal Plants Studies*. 2016: 4(4), 84-87.
19. Edeoga HO, Okwu D, Mbaebie B. Phytochemical constituents of some Nigerian medicinal plants. *African journal of biotechnology*. 2005: 4(7), 685-688.
20. Ayoola P, Adeyeye A, Onawumi O. Trace elements and major minerals evaluation of *Spondias mombin*, *Vernonia amygdalina* and *Momordica charantia* leaves. *Pakistan journal of nutrition*. 2010: 9(8), 755-758.
21. Trease G, Evans W. *Pharmacognosy*. 11th edn. Brailliar Tiridel Can. 1989. Macmillian publishers.

22. Harborne JB. Phytochemical Methods. 1973. 33-41. Phytochemical Methods. A Guide to modern Techniques of Plant Analysis. Chapman and Hall London. 1973: 33-41.
23. Ayodele OD, Oyegbade O, Oseni SR. Phytochemical analysis and antioxidant activities of dry and fresh leaves of *Petivera alliacea* and *Ocimum gratissimum*. International Journal of Sciences: Basic and Applied Research. 2015: 24(3), 1-13.
24. Boham B, Kocipai-Abyazan R. Flavonoids and condensed tannins from leaves of *Hawaiian vaccinium vaticulatum* and *V. calycinium*. Pacific Sci. 1974: 48(4), 458-463.
25. Karpagasundari C, Kulothungan S. Analysis of bioactive compounds in *Physalis minima* leaves using GC MS, HPLC, UV-VIS and FTIR techniques. Journal of Pharmacognosy and Phytochemistry. 2014: 3(4), 196-201.
26. Ruch RJ, Cheng SJ, Klaunig JE. Prevention of cytotoxicity and inhibition of intercellular communication by antioxidant catechins isolated from Chinese green tea. Carcinogenesis. 1989: 10(6), 1003-1008.
27. Aebi H. Catalase in vitro Methods in enzymology. 1984. Elsevier. 105, 121-126.
28. Luck H. Estimation of catalase, methods in enzymatic analysis. Bergmeyer, HU, & Gawhn, K. (Eds.), 1974: 885-894.
29. Velavan S. Free radicals in health and diseases-A Mini Review. Pharmacologyonline Newsletter, 2011: 1, 1062-1077.
30. Sathiyabalan GPK, Tresina PS, Muthukumarasamy S, Mohan VR. Pharmacognostical, Physicochemical and Phytochemical Standardization of *Petiveria alliacea* L. . International Journal of Pharmacognosy and Phytochemical Research. 2017: 9(2), 233.
31. Jain PK, Soni A, Jain P, Bhawsar J. Phytochemical analysis of *Mentha spicata* plant extract using UV-VIS, FTIR and GC/MS technique. *J Chem Pharm Res*. 2016: 8(2),1-6.
32. Johnson M, Fathima MSA. Spectroscopic studies on *Pouzolzia wightii* benn. International Journal of Pharmacy and Pharmaceutical Sciences. 2018: 10(3), 124-132.
33. Saxena M, Saxena J. Evaluation of phytoconstituents of *Acorus calamus* by FTIR and UV-VIS spectroscopic analysis. International Journal of Biological & Pharmaceutical Research. 2012: 3(3), 498-501.
34. Sahu N, Saxena J. Phytochemical analysis of *Bougainvillea glabra Choisy* by FTIR and UV-VIS spectroscopic analysis. Int. J. Pharm. Sci. Rev. Res. 2013: 21(1), 196-198.
35. Naidoo CM, Naidoo Y, Dewir YH, Murthy HN, El-Hendawy S, Al-Suhaibani N. Major bioactive alkaloids and biological activities of *Tabernaemontana* species (Apocynaceae). Plants. 2021: 10(2), 313.
36. Choi KC, Son YO, Hwang JM, Kim BT, Chae M, Lee JC. Antioxidant, anti-inflammatory and anti-septic potential of phenolic acids and flavonoid fractions isolated from *Lolium multiflorum*. Pharmaceutical biology. 2017: 55(1), 611-619.
37. Hacisevki A, Baba B. An overview of melatonin as an antioxidant molecule: a biochemical approach. Melatonin molecular biology, clinical and pharmaceutical approaches. 2018: 59-85.
38. Sowndhararajan K, Kang SC. Free radical scavenging activity from different extracts of leaves of *Bauhinia vahlii Wight & Arn*. Saudi journal of biological sciences. 2013: 20(4), 319-325.
39. Glorieux CZM, Sandoval JM, Verrax J, Calderon PB. Regulation of catalase expression in healthy and cancerous cells. Free Radical Biology and Medicine. 2015: 1(87), 84-97.
40. Mehta SK, Gowder SJT. Members of antioxidant machinery and their functions. Basic principles and clinical significance of oxidative stress. 2015: 11, 59-85.
41. Ighodaro OM, Adeosun AM, Adeoye BA, Gabriel O. Total Antioxidant Capacity of *Sapium ellipticum* (Hochst) Pax Leaf Extract, and Its Antioxidant Enzymes Potential in Free-Radical Systems in vitro. Journal of American Science. 2017: 13(7), 113-118.
42. Valyova M, Stoyanov S, Markovska Y, Ganeva Y. Original Research Evaluation of in vitro antioxidant activity and free radical scavenging potential of variety of *Tagetes erecta* L.

flowers growing in Bulgaria. International Journal of Applied Research in Natural Products. 2012: 5 (2), 19-25.

43. Maity S, Chatterjee S, Variyar PS, Sharma A, Adhikari S, Mazumder S. Evaluation of antioxidant activity and characterization of phenolic constituents of *Phyllanthus amarus* root. Journal of agricultural and food chemistry. 2013: 61(14), 3443-3450.

44. Ghasemzadeh A, Ghasemzadeh N. Flavonoids and phenolic acids: Role and biochemical activity in plants and human. Journal of medicinal plants research. 2011: 5(31), 6697-6703.

UNDER PEER REVIEW