

Original Research Article

Effect of *Terminalia Catappa* Seed Extract on Liver Enzymes and Lipid profile of Male Wistar Rat

ABSTRACT

The traditional use of plant-derived natural substances has continued to gain traction in the last few decades and has gradually become the mainstay of human pharmacotherapy. The present evaluated the effect of *T. catappa* seed extract on the lipid profile and serum liver enzymes of male Wistar rats. Fresh, ripe fruits of *Terminalia Catappa* were sourced locally from the University of Port Harcourt and the dried seeds were extracted using methanol and used for the study. Fifteen (15) male Wistar rats (160 – 180g) were grouped into three (3) of five (5) animals each and used for the study. Group I served as the control and received distilled water while groups II and III received 500 and 1000mg/kg of ethanolic seed extract of *T. catappa* (TEC) by oral gavage for twenty-eight (28) days. Animals were sacrificed by cardiac puncture and the blood collected was analyzed for serum liver enzymes (aspartate transaminase (AST), alanine Transaminase (ALT), and alkaline phosphatase (ALP) and lipid profile (total cholesterol (TC), triglycerides (TG), high-density lipoproteins (HDL) and low-density lipoproteins (LDL) using standard laboratory methods. Results obtained from the study indicate that serum ALT and LDL were significantly increased among rats that received TEC compared to the control ($p < 0.05$). The study concludes that TEC exhibited possible hepatoprotective properties and modulation of lipid profile and hence may be effective against liver injury and protection against cardiovascular and neurodegenerative diseases.

Key words: *Terminalia Catappa*, liver enzyme, lipid profile, hepatoprotection, lipid peroxidation

INTRODUCTION

Terminalia catappa is among the most common trees found in Nigeria. The generic name *terminalia* originates from “terminalis”, a Latin word that refers to the way the leaves team at the end of the shoots. It belongs to the family of Combretaceae. The tree can grow up to 35 meters tall and has a broad, spreading canopy. Its leaves are large, glossy, and oval-shaped, and can range in colour from green to red to copper. The tree produces small, white flowers that grow in clusters, which are followed by oval-shaped fruits that contain a hard, woody shell surrounding a seed, which can be eaten raw [1, 2]. They are grown for ornamental and medicinal purposes. In Nigeria, the almond tree is found in foremost towns and villages where the trees serve as shades with the seed highly sought by children who consume them as a foraged snack and by some rural dwellers as a protein supplement in local delicacies and the preparation of local herbs [3-5].

Several cultivated almond varieties display a different chemical profile in their leaves, barks, fruits and seeds due to genetic and ecological factors, as well as processing conditions [1, 6, 7].

They typically contain lipids, carbohydrates, proteins and other bioactive compounds. The leaves of *T. catappa* contain glucose, phenols, tannins, flavones, xanthenes, alkaloids, flavonoids and triterpene steroids [8-11]. The fruit contains proteins, carbohydrates, β -carotene, glucose, tannin, oil and vitamin C [2, 11] while the bark is rich in glycoside, cardiac tannins, volatile oil, saponin and phenols while the fruit contains protein, carbohydrate, β -carotene, glucose, tannin, oil and vitamin C [11]. The seed is rich in fixed oils, olein, stearin, crude fibre, protein, fat and carbohydrate and phytochemicals such as terpenoids, alkaloids, tannins, steroids, flavonoids, saponin, glycosides, phenols, anthraquinones and phlobaphenins [11-13] and in other minerals such as calcium, iron, phosphorus, potassium, magnesium, copper and zinc [14]. These diverse phytochemical constituents confer on various parts, several applications in folk remedies. The leaves, barks and roots are used in traditional medicine. Fallen leaves of *T. catappa* used to treat headaches, liver diseases, colic and as a diuretic [15]. In Ayurveda, the fruit is referred to as the “king of medicine” and used in the treatment of inflammation, asthma, ulcer, diarrhoea and allergies [16]. The seeds are thought to possess aphrodisiac activity, sexual malfunction and premature ejaculation [17, 18]. Documented research evidence has shown the seeds of *T. catappa* to possess anthelmintic [19], antioxidant [20, 21], aphrodisiac [18], antidiabetic and hypolipidaemic potentials [22].

The leaves, bark and fruit of *T. catappa* have been evaluated for their pharmacological actions [23-27], however, the seed (or nut/kernel) has remained minimally explored for potential effects on biochemical parameters. The present study, therefore, is aimed at evaluating the effect of *T. catappa* seed extract on the lipid profile and serum liver enzymes of Wistar rats to plug the knowledge gap on the safety of the seeds and shed more light on their effect on lipid metabolism.

MATERIAL AND METHODS

Source and Preparation of Plant Material

Fresh, ripe fruits of *T. catappa* were collected from host trees at the University of Port Harcourt, Port Harcourt, Nigeria from February through April 2022. The fruits were meticulously cracked open to expose the seed within. The fresh fruit of *T. catappa* was identified at the Department of Plant Science and Biotechnology, Faculty of Science, University of Port Harcourt and deposited with a herbarium number UPH/P/267. Seeds of *T. catappa* were shed dried for 4 weeks and then ground into a coarse powder. The ground seeds were then extracted by the method of Soxhlet [28]. *T. catappa* extract (TCE) was refrigerated at 5°C pending administration.

Research Animals

Fifteen (15) male Wistar rats (160 – 180g) were locally sourced from the Animal house of the Department of Human Physiology and used for the study. The animals were housed in standard rat cages under hygienic animal husbandry conditions: temperature, 25 - 28°C; humidity 40 – 60% while maintaining a 12hr light/dark cycle. The animals were allowed to acclimatise for 4 weeks to their new environment during which they were allowed a standard rat chow and water *ad libitum*.

Experimental Design

The fifteen (15) male Wistar rats were randomly grouped into three (3) groups of five (5) animals each. Group I served as the control and received distilled water, while groups II and III served as the experimental animals and received a methanolic extract of *T. catappa* at 500 and 1000mg/kg respectively by oral gavage. Daily administration of distilled water and extracts of *T. catappa* lasted for 28 days between 8 – 10 am. The animals were weighed before and after the experiment using a digital weighing scale (SF400 Shenzhen Jufengcai Electronic Technology Co., LTD, China). The animals were then anaesthetized with sodium thiopental (50 mg/kg, i.p.) and blood was collected through a cardiac puncture into plain sample bottles for laboratory analysis.

Laboratory Analysis

Blood samples were centrifuged at 3000rpm (Xiangtian 800-1, China). The supernatant serum was carefully collected using an automatic micropipette (Rong Tai Biochemical Engineering Co., Ltd, China). Lipid profile: total cholesterol (TC), triglycerides (TG) and high-density lipoproteins (were determined using standard laboratory kits (Randox, UK) while low-density lipoproteins were estimated using Friedewald's equation [29]. Liver enzymes: aspartate transaminase (AST), alanine Transaminase (ALT), and alkaline phosphatase (ALP) were determined using colourimetric methods (Randox, UK).

Ethical Considerations

Animals used for the study were housed and handled in compliance with standard guidelines and care of the use of laboratory animals [30, 31]. The research design and protocol were approved by the intuitional research ethic committee.

Statistical Analysis

Data obtained from laboratory investigations in the study were analyzed using IBM Statistical Product and Service Solutions (SPSS version 25). The mean and standard error of the mean were calculated for each parameter. The means were compared using the analysis of variance (ANOVA) followed by a least significant difference (LSD) posthoc analysis (ANOVA). A p-value less than 0.05 ($p < 0.05$) was considered statistically significant.

RESULTS

Table 1: The effect of *T. catappa* on serum liver enzymes of male Wistar rats

Groups	Treatment	AST (iu/L)	ALT (iu/L)	ALP (iu/L)
I	Distilled water	36.20 ± 1.66	18.00 ± 0.77	93.60 ± 6.30
II	500 mg/kg TCE	35.80 ± 8.22	19.40 ± 1.60	84.20 ± 5.68
III	1000 mg/kg TCE	31.00 ± 4.23	17.80 ± 0.80	50.20 ± 3.25*

values expressed as mean ± standard error of mean; *significant difference compared to control ($P < 0.05$)

Table 1 shows the effect of 28 days of administration of methanolic extract of *T. catappa* on the serum liver enzyme of male Wistar rats. The results indicate that there was a significant reduction in the mean values of serum ALP among the animals that received 1000mg/kg of *T. catappa* compared to the control ($p < 0.05$). Though the mean serum values of AST and ALT showed a dose-dependent reduction among the 500 and 1000mg/kg when compared with the control, it was not statistically significant.

Table 2: The effect of *T. catappa* on the serum lipid profile of male Wistar rats

Groups	Treatment	TC (mmol/L)	TG (mmol/L)	HDL (mmol/L)	LDL (mmol/L)
I	Distilled water	3.61 ± 0.23	0.43 ± 0.13	2.47 ± 0.21	0.92 ± 0.10
II	500 mg/kg TCE	3.24 ± 0.19	0.34 ± 0.34	2.89 ± 0.76	0.20 ± 0.14*
III	1000 mg/kg TCE	3.12 ± 0.51	0.31 ± 0.25	2.84 ± 0.55	0.14 ± 0.10*

values expressed as mean ± standard error of mean; *significant difference compared to control ($P < 0.05$)

Table 2 shows the effect of 28 days of administration of methanolic extract of *T. catappa* on the liver profile of male Wistar rats. Our results indicate that there was a dose-dependent reduction in the mean values of serum LDL among animals that received 500 and 1000mg/kg of *T. catappa* compared to the control ($p < 0.05$). Though the mean serum values of TC and TG showed a gradual dose-dependent reduction among the 500 and 1000mg/kg when compared with the control, it was not statistically significant. Similarly, the mean values of serum HDL were observed to show a dose-dependent non-significant increase among the experimental groups compared to the control.




Fig 1. The effect of *T. catappa* on percentage weight changes of male Wistar rats.

Data on the percentage weight gain of weight of male Wistar rats following a 28 days administration of methanolic *T. catappa* seed extract is shown in the bar charts above. Data obtained showed that group II animals (500mg/kg) had the highest percentage increase in their weight gain (46.98%) followed by group I (control) (46.68%) and group III (1000mg/kg) (42.9). However, there was no significant changes among the groups and the experimental groups were compared with the control.

DISCUSSION

The traditional use of plant-derived natural substances has continued to gain traction in the last few decades and has gradually become the mainstay of human pharmacotherapy [32, 33]. The present investigated the effect of the oral administration of Terminalia catappa seed extract (TCE) on liver enzymes and lipid profile and male Wistar rats.

Effect on Liver Enzymes

Results obtained from this study that the oral administration of TCE caused a significant in the mean values of serum ALP among the animals that received 1000mg/kg of *T. catappa* compared to the control ($p < 0.05$) (Table 1). Data also indicate that the mean serum values of AST and ALT showed a dose-dependent reduction among the 500 and 1000mg/kg when compared with the control, however, it was not statistically significant. The liver functions in many metabolic processes in the body and hence it is the primary target of most toxicants. It plays a role in clearing and transforming toxicants and other chemicals [34]. Routine measurement of liver function includes the measurement of serum AST, ALT and ALP often used as indicators of liver function and can be affected by liver damage or disease [35-37]. Though they are all found in the kidney, muscle and liver, they are predominately used to access liver function. The observed significant reduction in ALP and the marginal reduction in AST and ALT could be attributed to the hepatoprotection potential of TCE. The seed extract of TEC has been shown to contain significant quantities of flavonoids and phenols [11-13] which possess potent hepatoprotective ability [38-40]. Also, TCE has been demonstrated to have antioxidant activity [20, 21] which has to ability to attenuate liver enzymes [38]. Similarly, the leaf extract of *T. catappa* has been shown to reverse chemically induced liver injury in experimental models [11, 41, 42]. The present, therefore, hypothesizes that TCE has potent hepatoprotective potential and could be useful in the treatment of the toxin-induced liver injury.

Effect on Lipid Profile

The result of the present study shows that there was a dose-dependent reduction in the mean values of serum LDL among animals that received 500 and 1000mg/kg of TEC compared to the control ($p < 0.05$) (Table 2). Also, serum TC and TG showed a marginal dose-dependent reduction among the 500 and 1000mg/kg when compared with the control, it was not statistically significant. On the other hand, serum HDL was observed to a show dose-dependent increase among the experimental groups (II and III) compared to the control. The possible TEC-attenuating effect on lipid profile is attributable to its antioxidant property [20, 21] and its ability to prevent lipid peroxidation. Lipid peroxidation is a process that occurs when free radicals react with lipids in cell membranes, resulting in damage to the cell membrane and the release of harmful byproducts [43, 44]. This process is implicated in the aetiology of cardiovascular and neurodegenerative diseases, and cancer [45, 46]. Flavonoids and other antioxidant phytochemicals contained in TEC can help to prevent lipid peroxidation by neutralizing free radicals and protecting cell membranes from damage and inhibiting the formation of reactive oxygen species (ROS) [47-49]. They also enhance the activity of antioxidant enzymes, such as superoxide dismutase (SOD), glutathione peroxidase (GPx) and catalase (CAT) which can

further protect cells from oxidative damage [38, 49]. Also, Flavonoids are known to impact lipid metabolism by reducing cholesterol synthesis or increasing LDL receptor expression [50, 51]. Also, saponins have been shown to suppress cholesterol absorption, increase lipid peroxidation and depress the rates of intestinal hepatic and intestinal cholesterol synthesis [51-54]. Leaf extracts of TEC have been shown to possess anti-hyperlipidaemic properties in experimentally induced hyperlipidaemia and diabetes [22, 55, 56]. Besides the significant reduction in LDL, the present study hypothesizes that TEC can attenuate lipid metabolism by marginally reducing TG and TC and increasing HDL.

Effect on Body Weight

There was no significant change in the body weight of animals treated with TEC as the data obtained showed that group II animals (500mg/kg) had the highest percentage increase in their weight gain (46.98%) followed by group I (control) (46.68%) and group III (1000mg/kg) (42.9). This could be due to the short duration of the administration of TEC. However, the leaves of *T. catappa* has shown potent anti-obesity activity in experimentally induced obesity animal models [55, 56].

CONCLUSION

In light of the paucity of data regarding the pharmacological effects of seed extract of *T. catappa*, the present study evaluated the effects of TEC on liver enzymes and lipid profile using Wistar rat models. The study concludes that TEC exhibited possible hepatoprotective properties and modulation of lipid profile. This study shows that *T. catappa* may be effective against liver injury and protection against cardiovascular and neurodegenerative diseases due to the reduction of LDL. Hence the consumption of *T. catappa* seeds as a dietary supplement is recommended.

REFERENCES

1. Thomson LAJ, Evans B. *Terminalia catappa* (tropical almond). Species Profiles for Pacific Island Agroforestry. 2006;2(2):1-20.
2. Anand A, Divya N, Kotti P. An updated review of *Terminalia catappa*. Pharmacognosy reviews. 2015;9(18):93.
3. Omodamiro OD, Omekara I. Evaluation of in-vitro antioxidant and in-vivo diuretic activities of ethanol leaves extract of *terminalia catappa* leaves. Journal of International Research in Medical and Pharmaceutical Sciences. 2017;12(13):94-104.
4. Mbah BO, Eme PE, Eze CN. Nutrient potential of Almond seed (*Terminalia catappa*) sourced from three states of Eastern Nigeria. African Journal of Agricultural Research. 2013;8(7):629-33.
5. Oboh B, Ogunkanmi B, Olasan L. Phenotypic diversity in *Terminalia catappa* from South Western Nigeria. Pakistan Journal of Biological Sciences: PJBS. 2008;11(1):135-8.
6. Gilman EF, Watson DG. *Terminalia catappa* tropical-almond. Gainesville: Institute of Food and Agricultural Sciences, University of Florida. 1994.
7. Ezeokonkwo CA, Dodson WL. The potential of *Terminalia catappa* (tropical almond) seed as a source of dietary protein. Journal of Food Quality. 2004;27(3):207-19.

8. Terças AG, Monteiro AdS, Moffa EB, Santos JRd, Sousa EMd, Pinto AR, et al. Phytochemical characterization of *Terminalia catappa* linn. Extracts and their antifungal activities against candida spp. *Frontiers in Microbiology*. 2017;8:595.
9. Ihuma JO, Noel DO, Adogo LY. Antimicrobial Activities of *Aerminalia catappa* linn. Leaves Extract on Some Selected Salmonella Species. *Journal of Biology and Nature*. 2021;13(1):35-41.
10. Zainudin R, Nanding A, Mohamad H, Sinsuat DRR, Abo S, Usman S, et al. Phytochemical and Antimicrobial Analyses of *Terminalia Catappa* Linn. Ethanol Leaf Extract and Its Effect on Treating Pimple. *Psychology and Education: A Multidisciplinary Journal*. 2022;4(6):571-80.
11. Gao J, Tang X, Dou H, Fan Y, Zhao X, Xu Q. Hepatoprotective activity of *Terminalia catappa* L. leaves and its two triterpenoids. *Journal of Pharmacy and Pharmacology*. 2004;56(11):1449-55.
12. Lawal AR, Olayinka BU, Abdulkareem KA, Abdulra'uf LB, Murthadah RA, Kayode OV. Proximate, Minerals And Phytochemical Evaluation Of Pericarp And Seed Of *Terminalia Catappa* Fruit. *The Bioscientist Journal*. 2022;10(2):191-203.
13. Salawu RA, Onyegbula AF, Lawal IO, Akande SA, Oladipo AK. Comparative study of the nutritional, phytochemical and mineral compositions of the nuts of tropical almond (*Terminalia catappa*) and sweet almond (*Prunus amygdalus*). *Ruhuna Journal of Science*. 2018;9(1):70-7.
14. Weerawatanakorn M, Janporn S, Ho C-T, Chavasit V. *Terminalia catappa* Linn seeds as a new food source. *Songklanakarinn Journal of Science & Technology*. 2015;37(5).
15. Owolabi MS, Lawal OA, Ogunwande IA, Hauser RM, Setzer WN. Chemical composition of the leaf essential oil of *Terminalia catappa* L. growing in Southwestern Nigeria. *American Journal of Essential Oils and Natural Products*. 2013;1(1):51-4.
16. Ravi L. A review on medicinal potential of *Terminalia catappa*. *International Journal of Green Pharmacy (IJGP)*. 2020;14(03):229-33.
17. Ladele B, Kpoviessi S, Ahissou H, Gbenou J, Kpadonou-Kpoviessi B, Mignolet E, et al. Chemical composition and nutritional properties of *Terminalia catappa* L. oil and kernels from Benin. *Comptes Rendus Chimie*. 2016;19(7):876-83.
18. Ratnasooriya W, Dharmasiri M. Effects of *Terminalia catappa* seeds on sexual behaviour and fertility of male rats. *Asian Journal of Andrology*. 2000;2(3):213-9.
19. Katiki LM, Gomes ACP, Barbieri AME, Pacheco PA, Rodrigues L, Veríssimo CJ, et al. *Terminalia catappa*: chemical composition, in vitro and in vivo effects on *Haemonchus contortus*. *Veterinary Parasitology*. 2017;246:118-23.
20. Adu OB, Omojufehinsi M, Esanboro M, Abe D, Shofolahan A, Uzodinma E, et al. Effect of processing on the quality, composition and antioxidant properties of *Terminalia catappa* (indian almond) seed oil. *African Journal of Food, Agriculture, Nutrition and Development*. 2013;13(3).
21. Halilu EM, Ugwah-Oguejiofor CJ, Oduncuoğlu G, Matthias SG. Physicochemical, Toxicity and Antioxidant Activity of *Terminalia catappa* Kernel Oil in Mice. *Pharmacognosy Research*. 2023;15(1):199-27.
22. Luka CD, Istifanus G, Olatunde A, Eli Z. Effect of Aqueous Seed Extract of *Terminalia catappa* Linn on Some Biochemical Parameters in Alloxan-induced Diabetic Rats. *Journal of Applied Life Sciences International*. 2017;13(3):1-8.

23. Venkatalakshmi P, Brindha P, Induja K. In-vitro anti oxidant and antitumor studies on Terminalia catappa bark. *Aging*. 2001;18:685-716.
24. Neelavathi P, Venkatalakshmi P, Brindha P. Antibacterial activities of aqueous and ethanolic extracts of Terminalia catappa leaves and bark against some pathogenic bacteria. *Int J Pharm Pharm Sci*. 2013;5(1):114-20.
25. Chyau C-C, Tsai S-Y, Ko P-T, Mau J-L. Antioxidant properties of solvent extracts from Terminalia catappa leaves. *Food chemistry*. 2002;78(4):483-8.
26. Akharaiyi F, Ilori R, Adesida J. Antibacterial effect of Terminalia catappa on some selected pathogenic bacteria. *Int J Pharm Biomed Res*. 2011;2(2):64-7.
27. Nagappa AN, Thakurdesai PA, Rao NV, Singh J. Antidiabetic activity of Terminalia catappa Linn fruits. *Journal of ethnopharmacology*. 2003;88(1):45-50.
28. Evans WC. *Trease and Evans' Pharmacognosy*. 16th ed. New York: Elsevier Health Sciences; 2009.
29. Friedewald WT, Levy RI, Fredrickson DS. Estimation of the concentration of low-density lipoprotein cholesterol in plasma, without use of the preparative ultracentrifuge. *Clinical Chemistry*. 1972;18(6):499-502.
30. Albus U. *Guide for the care and use of laboratory animals* (8th edn). SAGE Publications Sage UK: London, England; 2012.
31. Benjamin B. Overview of laboratory animal lifestyle, care, and management: a case study of albino rats. *Journal of Applied Sciences and Environmental Management*. 2019;23(8):1431-5.
32. Atanasov AG, Waltenberger B, Pferschy-Wenzig E-M, Linder T, Wawrosch C, Uhrin P, et al. Discovery and resupply of pharmacologically active plant-derived natural products: A review. *Biotechnology advances*. 2015;33(8):1582-614.
33. Ugoeze KC, Oluigbo KE, Chinko BC. Phytochemical and Nutritional Benefits of the GC-FID Quantified Phytochemicals of the Aqueous Extract of Azadirachta indica leaves. *Journal of Pharmacy and Pharmacology Research*. 2020;4(4):149-63.
34. García-Niño WR, Zazueta C. Ellagic acid: Pharmacological activities and molecular mechanisms involved in liver protection. *Pharmacological Research*. 2015;97:84-103.
35. Gowda S, Desai PB, Hull VV, Avinash AK, Vernekar SN, Kulkarni SS. A review on laboratory liver function tests. *The Pan african medical journal*. 2009;3.
36. Macpherson I, Abeysekera KWM, Harris R, Mansour D, McPherson S, Rowe I, et al. Identification of liver disease: why and how. *Frontline Gastroenterology*. 2022;3(5).
37. Chinko BC, Pughikumo DT. Haematological and Hepatorenal Alterations Induced by Potash (Akanwu) on Male Wistar Rats. *International Blood Research & Reviews*. 2023;14(1):38-46.
38. Roy A, Bhoumik D, Sahu RK, Dwivedi J. Medicinal plants used in liver protection-a review. *Pharmaceutical and Biosciences Journal*. 2014;2(1):23-33.
39. Pan X, Ma X, Jiang Y, Wen J, Yang L, Chen D, et al. A comprehensive review of natural products against liver fibrosis: flavonoids, quinones, lignans, phenols, and acids. *Evidence-based complementary and alternative medicine*. 2020;2020.
40. Wang Y, Zhang N, Zhou J, Sun P, Zhao L, Zhou F. Protective Effects of Several Common Amino Acids, Vitamins, Organic Acids, Flavonoids and Phenolic Acids against Hepatocyte Damage Caused by Alcohol. *Foods*. 2022;11(19):3014.
41. Tang X-h, Gao J, Wang Y-p, Xu L-z, Zhao X-n, Xu Q. Hepatoprotective effects of chloroform extract from leaf of Terminalia catappa in relation to the inhibition of liver

- IL-6 expression. *Zhongguo Zhong yao za zhi= Zhongguo Zhongyao Zazhi= China Journal of Chinese Materia Medica*. 2003;28(12):1170-4.
42. Tang XH, Gao J, Dou H, Wang YP, Xu LZ, Zhu ZR, et al. Protective effect of the extract of *Terminalia catappa* leaves on acute liver injury induced by D-GalN in mice. *Zhongguo Zhong yao za zhi= Zhongguo Zhongyao Zazhi= China Journal of Chinese Materia Medica*. 2004;29(11):1069-73.
 43. Gaschler MM, Stockwell BR. Lipid peroxidation in cell death. *Biochemical and biophysical research communications*. 2017;482(3):419-25.
 44. Repetto M, Semprine J, Boveris A. Lipid peroxidation: chemical mechanism, biological implications and analytical determination. *Lipid peroxidation*. 2012;1:3-30.
 45. Singh R, Devi S, Gollen R. Role of free radical in atherosclerosis, diabetes and dyslipidaemia: larger \square than \square life. *Diabetes/metabolism research and reviews*. 2015;31(2):113-26.
 46. Valko M, Leibfritz D, Moncol J, Cronin MTD, Mazur M, Telser J. Free radicals and antioxidants in normal physiological functions and human disease. *The international journal of biochemistry & cell biology*. 2007;39(1):44-84.
 47. Banjarnahor SD, Artanti N. Antioxidant properties of flavonoids. *Medical Journal of Indonesia*. 2014;23(4):239-44.
 48. Karak P. Biological activities of flavonoids: an overview. *Int. J. Pharm. Sci. Res*. 2019;10(4):1567-74.
 49. Terao J. Dietary flavonoids as antioxidants. *Food factors for health promotion*. 2009;61:87-94.
 50. Zeka K, Ruparelia K, Arroo RR, Budriesi R, Micucci M. Flavonoids And Their Metabolites: Prevention In Cardiovascular Diseases And Diabetes. *Diseases*. 2017;5(3):1-18.
 51. You C-L, Su P-Q, Zhou X-X. Study On Effect And Mechanism Of *Scutellaria Baicalensis* Stem-Leaf Total Flavonoid In Regulating Lipid Metabolism. *China Journal of Chinese Materia Medica*. 2008;33(9):1064-6.
 52. Barbalho SM, Damasceno DC, Spada APM, Silva VSd, Martuchi KA, Oshiiwa M, et al. Metabolic profile of offspring from diabetic Wistar rats treated with *Mentha piperita* (peppermint). *Evidence-Based Complementary and Alternative Medicine*. 2011;2011.
 53. Alor PC, Chinko BC. Flat Tummy Water Attenuates Lipid Profile and Serum Glucose of High-Fat Diet-Induced Obese Female Wistar Rats. *Journal of Complementary and Alternative Medical Research*. 2022;20(4):19-28.
 54. Ikete PW, Chinko BC. Inhibition of Serum Lipase as a Mechanism of Action for Anti-Obesity Potentials of *Dioscorea bulbifera* Extracts on Wistar Rats. *Asian Journal of Health Sciences*. 2022;8(1):ID29-ID.
 55. Mukundan U, Sharma A. Anti-Obesity And Anti-Hyperlipidemic Activity Of *Terminalia Catappa* Linn. In High-Fat-Diet Induced Obese Rats. *Asian Journal of Pharmaceutical Research and Development*. 2013;1(6):114-20.
 56. Ambidevi S. Anti-Obesity And Anti-Hyperlipidemic Activity Of *Terminalia Catappa* Linn. High-Fat-Diet Induced Obese Rats. *Asian Journal of Pharmaceutical Research and Development, Mumbai*. 2014;1(6).