

## Review Article

### “Prevalence of Anxiety During COVID-19 Pandemic: A Comprehensive Review on Scientific Literatures”

#### **ABSTRACT:**

The global severity of COVID-19 remains high which results anxiety and other mental health problems, also it altered people's everyday lives, affected human connections and economic operations. The goal of this comprehensive review was to identify the effects of the linkage COVID-19 pandemic on the mental health of different groups and communities. This study compiled evidences of a link between anxiety rates and the COVID-19 pandemic. The evaluation period started in June' 2022 and ended on August'2022, during this time, total four databases such as PubMed, ScienceDirect, Tailor & Francis Online, and Springer were used to search scientific literatures. A total 616 studies were identified from all four databases and 63 scientific literatures were selected based of predetermined criteria for review which were published in between 2020 to 2022. Three groups of population such as general population, students and healthcare professionals were taken for review the findings from the selected literatures. Gender, physical disorders, psychiatric disorders, COVID infection, infection rates in colleagues or family members, experience of frontline work & non-frontline work, close contact with infected patients, high exposure risk, quarantine experience, etc. were highly considered as factors associated with increased prevalence of anxiety among all three groups. During the COVID-19 pandemic, the general population, healthcare professionals, and students experienced an increase in the prevalence of mental diseases, whereas infected individuals had a decrease. Females were highly prevalent to anxiety than male. Our comprehensive review concluded significant correlation between anxiety and COVID-19 but long-term study is needed to better understand which may define the population's mental condition in future.

Keyword: COVID-19, Coronavirus Disease, Anxiety, Mental Health Disorders

#### **Introduction:**


Pandemic potential infectious illnesses like plague, cholera, flu, SARS-CoV, and MERS-CoV have regularly emerged and spread throughout history, and now the world faces COVID-19, declared a global pandemic by WHO on 11th March 2020 [1-6]. In December 2019, health workers identified a novel coronavirus 2019 (2019-nCoV) after discovering many cases admitted to hospitals with symptoms similar to viral pneumonia, including fever, cough, chest pain, dyspnea, and lung infiltration [7, 8] and the spread of the new variant of coronavirus began at the Huanan Seafood Wholesale Market in Wuhan, Hubei Province [3, 5, 6]. In order to minimize the spread of viral outbreaks in communities, nations worldwide took precautionary measures such as imposing nationwide lockdowns, shutting down institutions, and isolating infected individuals, which were essential, but they also stimulated various psychological stressors such as experiencing fear and panic, feeling frustrated and bored, facing a scarcity of basic supplies,

lacking authentic and reliable information, being overwhelmed with stigma, losing jobs, and facing financial recession [9, 10]. The possibility of contracting the virus, healthcare system overload, lack of effective treatment options, and vaccine hesitancy were significant stressors that have led to a range of mental health disorders in the population, including anxiety, depression, psychological distress, stress, post-traumatic disorders, and suicidal tendencies [11-14].

The main objective of this study is to review the correlation between anxiety and COVID-19 across demographics to inform future investigations.

**Materials and Method:**

The review period began in June 2022 and updated in July and August. Popular databases, such as PubMed, ScienceDirect, Springer, and Tailor & Francis Online, were selected. Bibliographic referencing was done using EndNote 20.3, and literature screening was done with MS Excel. Information was analyzed by publication year, authors, sample, country, and study type. The records were categorized by demographic types in an MS Excel sheet. Sample sizes, study parameters, design, outcomes, and key findings were gathered for the review.



**Figure 1: Literature Screening Flowchart**

**Results:**

Anxiety can arise from changes in homeostasis, sometimes due to infectious disorders [15, 16]. However, exaggerated perceptions can lead to significant issue [7, 17]. COVID-19 has increased anxiety in recent years, especially with lockdown and isolation measures causing frustration and fear [17, 18]. Exposure to conflicting information on social media during the pandemic may cause psychological difficulties [19, 20] Pregnant women and students may mistake COVID-19 symptoms for anxiety, leading to self-isolation and medical attention-seeking [21-23]. Excessive stress can lead to abnormal behaviors, such as hand washing, social withdrawal, and panic over trivial issues, with negative consequences for individuals, families, and society, including financial resource shortages [15, 24]. Health workers, the frontline warriors in the fight against

COVID-19, face anxiety over the risk of infection and transmission to others, especially as many become sick themselves [25-28].

The largest group of studies in the present study focused on mental health in the general population, with anxiety being the most prevalent issue at almost 35%. Four studies [29-32] reported a three-fold increase in anxiety during the COVID-19 pandemic compared to previous years, with rates increasing from 6% in 2017 to 19% during the pandemic. While another study conducted by Peter G van der Velden, Carlo Contino, Marcel Das, Peter van Loon and Mark W G Bosmans found no significant change in anxiety rates between 2019 and 2020, authors noted changes in risk factors since the pandemic, particularly affecting students, job seekers, and housekeepers [32, 33]. Anxiety rates vary based on the location's COVID-19 risk level. Red zone areas had higher anxiety rates than others [30, 34-40], while areas with higher testing rates had lower anxiety rates [41]. Fang Tang's online survey of 1160 quarantined people in Wuhan, China, found that 70.78% experienced anxiety disorder due to the COVID-19 lockdown [41, 42]. Another study found that COVID-19-related anxiety was linked to the lockdown's effect on family finances, as people were worried about the expense of protective measures [43]. Many studies have identified job loss or financial difficulties resulting from COVID-19 as common predictors or causes of increased anxiety [21, 29, 39, 44-47]. A cross-sectional study by Xi-Ru Zhang's team on 123,768 Chinese workers found that 26.2% of the population had mental health disorders, of which 3.4% had anxiety during the pandemic [38]. Similarly, Constanza Jacques-Aviñó's team reported anxiety and other mental disorders in 31.2% of women and 17.7% of men among 7,053 individuals in Spain during the pandemic [45]. Several studies supported these findings [35, 38, 48-50], but males with dependents were more anxious during the pandemic [45, 51]. Longitudinal studies in Australia and Germany reported a 23% rise in anxiety during a 12-week quarantine period, which decreased when restrictions were eased [25, 35, 52]. People who had more contact with COVID-19 infected individuals, particularly healthcare workers in hospitals, were found to be more anxious than those who stayed at home during lockdown [53-57]. Furthermore, many studies during the pandemic found that individuals infected with COVID-19 were more anxious than those who were not infected [45, 46, 56, 58].

Data on the impact of COVID-19 on college and university students' mental health is limited, but 23 reports showed anxiety levels as mild, moderate, or severe. Akhtarul et al.'s 2020 web-based survey found that more than 85% of students (n=476) experienced mild to severe anxiety symptoms, with females (26.1%) reporting severe anxiety more frequently than males (17%) [41]. Female students reported higher levels of severe anxiety compared to male students during the COVID-19 pandemic in four different studies [39, 40, 46, 47]. A study in Bangladesh found that anxiety levels among university students increased during the pandemic, with anxiety related to academic and career concerns [59]. Chinese students reported increased anxiety due to daily life restrictions during lockdowns, according to a study by Wenjun Cao and colleagues [21, 60]. Lockdowns during the pandemic led to increased loneliness and lack of social support, contributing to a rise in clinical anxiety levels among university and college students [21, 46]. In a web-based survey conducted by Iqbal and his team in Bangladesh, nearly 80% of university

students and workers (n=2,350) reported moderate to severe mental health disorders such as depression, anxiety, and stress during the pandemic, with undergraduate students showing more anxiety than postgraduates or job holders [61]. Similarly, Odriozola-Gonzalez found a significant number of undergraduate students experiencing severe anxiety in a study of Spanish university students and workers (n=3,707) [60], while two other studies reported high anxiety levels for postgraduates and job holders [46, 59]. Two studies on school teachers with a combined sample size of 90,244 found that 27% to 49% of respondents had COVID-19-associated anxiety, with female teachers experiencing higher anxiety levels [62, 63]. Students' anxiety during the COVID-19 pandemic was linked to fear of infection and exposure to COVID-19 news, reported by Sun et al. and Wathélet et al. in web-based surveys on Chinese and European students [39, 47]. Anxiety was also reported by over one-third of students from various countries including Ethiopia [64], France [39, 65], India [66], Israel [67], Jordan [68], Nigeria [69], the Philippines [70], Saudi Arabia [71], and the United Arab Emirates [72].

The COVID-19 outbreak has led to high psychological stress among healthcare workers, who are considered a high-risk group with long-term psychological repercussions [58, 73]. Factors such as challenging working conditions, heavy workloads, inadequate protective equipment, risk of infection, and difficult ethical decisions have led to mental health issues like anxiety [9, 73-75]. In Spain, a survey of 1,422 healthcare workers found that 58.6%, including doctors and nurses, experienced severe anxiety [76]. Some comprehensive reviews found that anxiety and depression rates during COVID-19 were 31-34% in the general population and community sectors, including healthcare professionals, children, and students which suggests potential mental health impacts of the pandemic [77-80]. Researchers from Imperial College London evaluated anxiety in 33,062 respondents, with a prevalence rate of 23.2% where, female medical staffs had higher anxiety rates [81-83]. Frontline healthcare workers had more anxiety due to COVID-19 exposure [73, 74, 81, 83-85]. However, some studies found higher anxiety rates in non-clinical staff and younger trainees [83, 86-90]. Non-clinical workers had decreased anxiety rates after safety training [88]. Chinese research found high rates of depression (50.4%), anxiety (44.6%), sleeplessness (34%), and psychological discomfort (71.5%) in frontline healthcare workers [90]. Argentine healthcare workers experienced high rates of sleep issues (73.7%), anxiety (76.5%), and depression (81.0%) [86]. Indonesian researchers found that lower resilience was associated with greater mental disorder with 33% having high state anxiety and 26.9% having high trait anxiety among healthcare workers during COVID-19 [91]. Gainer et. al. found a significant link between treating COVID-19 for more days and higher anxiety risk in US physician [70, 89].

### **Discussion:**

Several recent systematic reviews have focused on the relationship between mental health and COVID-19, but many studies have only examined specific populations. Our study is the most comprehensive analysis to date, focusing on three large groups: the general population, students, and healthcare professionals. The authors reviewed numerous studies on mental health problems, particularly anxiety. The authors found that mental health concerns were common and varied among different populations, including the general community, healthcare workers, students,

older adults, infected patients, and survivors. The authors identified unique mental health concerns for various demographics and proposed a preventative plan based on risk factor analysis.

Lockdowns were implemented worldwide to reduce COVID-19 transmission, but caused significant emotional distress globally and strong mental health at the beginning of lockdowns deteriorated as they became longer and more-strict [92]. Quarantine and lockdown orders increased loneliness, isolation, and anxiety in the general population, healthcare professionals, and students [59, 86, 93-95]. A group of systematic reviews showed that anxiety was more prevalent in 2020 than in 2019 due to COVID-19 restrictions, such as home quarantine, physical and social isolation, and job loss [21, 29, 39, 41, 44, 46, 96-99]. While some studies found little effect of lockdowns on anxiety, these studies had limited sample size and variability in the data [11, 19, 21, 26, 28]. However, some studies suggesting that lockdown and quarantine orders have little or no impact on anxiety [32], the limited sample size in some of these studies, which lacked a wide range of socioeconomic diversity and exhibited data variability, could account for these results [100].

The COVID-19 pandemic has increased anxiety among healthcare workers, especially in clinical roles due to direct patient contact and greater susceptibility to illness. Non-clinical workers have also reported anxiety due to inadequate crisis training, though this improved after receiving such training. Fear of infecting family predicted health and job-related anxiety in healthcare workers in which younger practitioners reported more anxiety, possibly due to lack of experience/training. Mental health issues were equally prevalent among infected patients and the general population, which contradicts expectations given that most patients were middle-aged adults. This observation may be explained by the lower odds ratios for mental symptoms in older individuals. Current study showed that populations residing in areas with higher COVID-19 infection rates had greater rates of moderate to severe anxiety compared to those in low-epidemic regions. Increased testing within epicenters significantly reduced public worry, lowering overall health anxiety and indicating successful epidemic containment. COVID-19 exposure, whether through chance contacts, work settings, or confirmed infection, was associated with elevated anxiety levels. Contact with infected family members was especially linked to higher anxiety risk. Job loss due to COVID-19 was identified as a significant contributor to increased anxiety, alongside financial instability and other mental health problems. The study also found that individuals with pre-existing mental health problems were at higher risk of deterioration due to the disruption of daily routines caused by quarantine and limitations. In lockdown period during COVID-19, the university students become more addicted to social media such as Facebook, Instagram, twitter etc. where COVID-19 related news may make them more anxious than general population.

The COVID-19 pandemic has negatively impacted the mental health of university students by causing disruption to their daily lives and predisposing them to stress, potentially leading to a public health crisis. Similar rates of anxiety were observed among university students and the general population, with suicidal ideation also being reported [101]. The transition to online

schooling during the COVID-19 pandemic has caused academic anxiety, loneliness, and decreased academic self-efficacy among students, particularly those who struggle with self-directed learning and this has been identified as a significant source of stress and may increase the risk of depression and suicidal tendencies. However, lockdown measures have provided opportunities for students to engage in hobbies and improve sleep patterns, the increased use of social media during lockdowns may contribute to anxiety related to COVID-19 news. The literature reports a mixed impact of online schooling and lockdowns on students' mental health [102]. On-campus students have higher anxiety and loneliness scores. Financial uncertainty of living on campus without work has contributed to mental deterioration among many university students [40].

The comprehensive review of literatures has limitations that need to be acknowledged. Firstly, the inclusion of only periodic literature may not fully capture the evolving nature of mental health during the COVID-19 pandemic. Secondly, more research is needed to better understand the link between COVID-19 and anxiety, including studies that focus on different populations and collect longitudinal data. Finally, more long-term studies are needed to understand the underlying causes of pandemic fear and to better prepare for future outbreaks and emergency situations. Follow-up research can help to improve our understanding of the mental health of the population.

#### **Conclusion:**

The study demonstrated that COVID-19 significantly increased anxiety across all groups, particularly among high-risk individuals such as healthcare workers. Maintaining good mental health is essential for societal progress, and thus it is crucial to identify and provide appropriate psychological interventions for susceptible individuals from various population layers during the current crisis.

#### References:

1. Piret J, Boivin G. Pandemics Throughout History. *Front Microbiol.* 2020;11. Epub 20210115. doi: 10.3389/fmicb.2020.631736. PubMed PMID: 33584597; PubMed Central PMCID: PMC7874133.
2. Khan M, Adil SF, Alkathlan HZ, Tahir MN, Saif S, Khan M, et al. COVID-19: A Global Challenge with Old History, Epidemiology and Progress So Far. *Molecules.* 2020;26(1). Epub 20201223. doi: 10.3390/molecules26010039. PubMed PMID: 33374759; PubMed Central PMCID: PMC7795815.
3. Batista P, Duque V, Luzio-Vaz A, Pereira A. Anxiety impact during COVID-19: a systematic review. *J Infect Dev Ctries.* 2021;15(3):320-5. Epub 20210331. doi: 10.3855/jidc.12730. PubMed PMID: 33839704.

4. Gralinski LE, Menachery VD. Return of the Coronavirus: 2019-nCoV. *Viruses*. 2020;12(2). Epub 20200124. doi: 10.3390/v12020135. PubMed PMID: 31991541; PubMed Central PMCID: PMC7077245.
5. Mitu MM, Nayan M, Hossain I, Nandi T. A review of Covid-19's impact on pregnancy. *Journal of Advances in Medicine and Medical Research*. 2022;34(23):350-60.
6. Priyanka RA, Nayan M, Hossain I, Alam M, Nandi T. The Influences of COVID-19 Pandemic on Patient with Eating Disorders-A Literature Review. *Journal of Pharmaceutical Research International*. 2022;34(56):62-76.
7. Hu B, Guo H, Zhou P, Shi ZL. Author Correction: Characteristics of SARS-CoV-2 and COVID-19. *Nat Rev Microbiol*. 2022;20(5):315. doi: 10.1038/s41579-022-00711-2. PubMed PMID: 35197601; PubMed Central PMCID: PMC8864972.
8. Fagbule OF. 2019 Novel Coronavirus. *Ann Ib Postgrad Med*. 2019;17(2):108-10. PubMed PMID: 32669985; PubMed Central PMCID: PMC7358814.
9. Sadigov R. Rapid Growth of the World Population and Its Socioeconomic Results. *ScientificWorldJournal*. 2022;2022:8110229. Epub 20220323. doi: 10.1155/2022/8110229. PubMed PMID: 35370481; PubMed Central PMCID: PMC8967589.
10. Cucinotta D, Vanelli M. WHO Declares COVID-19 a Pandemic. *Acta Biomed*. 2020;91(1):157-60. Epub 20200319. doi: 10.23750/abm.v91i1.9397. PubMed PMID: 32191675; PubMed Central PMCID: PMC7569573.
11. Lesin Gacina D, Jandrokovic S, Marcinko D, Skegro I, Vidas Pauk S, Tomic M, et al. Anxiety and Treatment Adherence among Glaucoma Patients during COVID-19 Pandemic and Earthquakes in Croatia. *Psychiatr Danub*. 2022;34(2):348-55. doi: 10.24869/psyd.2022.348. PubMed PMID: 35772157.
12. Wang C, Tee M, Roy AE, Fardin MA, Srichokchatchawan W, Habib HA, et al. The impact of COVID-19 pandemic on physical and mental health of Asians: A study of seven middle-income countries in Asia. *PLoS One*. 2021;16(2):e0246824. Epub 20210211. doi: 10.1371/journal.pone.0246824. PubMed PMID: 33571297; PubMed Central PMCID: PMC7877638.
13. Wang C, Chudzicka-Czupala A, Tee ML, Nunez MIL, Tripp C, Fardin MA, et al. A chain mediation model on COVID-19 symptoms and mental health outcomes in Americans, Asians and Europeans. *Sci Rep*. 2021;11(1):6481. Epub 20210319. doi: 10.1038/s41598-021-85943-7. PubMed PMID: 33742072; PubMed Central PMCID: PMC7979938.
14. Xiong J, Lipsitz O, Nasri F, Lui LMW, Gill H, Phan L, et al. Impact of COVID-19 pandemic on mental health in the general population: A systematic review. *J Affect Disord*. 2020;277:55-64. Epub 20200808. doi: 10.1016/j.jad.2020.08.001. PubMed PMID: 32799105; PubMed Central PMCID: PMC7413844.

15. Asmundson GJG, Rachor G, Drakes DH, Boehme BAE, Paluszek MM, Taylor S. How does COVID stress vary across the anxiety-related disorders? Assessing factorial invariance and changes in COVID Stress Scale scores during the pandemic. *J Anxiety Disord.* 2022;87:102554. Epub 20220303. doi: 10.1016/j.janxdis.2022.102554. PubMed PMID: 35278802; PubMed Central PMCID: PMC8893927.
16. Xiao H, Zhang Y, Kong D, Li S, Yang N. Social Capital and Sleep Quality in Individuals Who Self-Isolated for 14 Days During the Coronavirus Disease 2019 (COVID-19) Outbreak in January 2020 in China. *Med Sci Monit.* 2020;26:e923921. Epub 20200320. doi: 10.12659/MSM.923921. PubMed PMID: 32194290; PubMed Central PMCID: PMC7111105.
17. Zhu N, Zhang D, Wang W, Li X, Yang B, Song J, et al. A Novel Coronavirus from Patients with Pneumonia in China, 2019. *N Engl J Med.* 2020;382(8):727-33. Epub 20200124. doi: 10.1056/NEJMoa2001017. PubMed PMID: 31978945; PubMed Central PMCID: PMC7092803.
18. Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet.* 2020;395(10223):497-506. Epub 20200124. doi: 10.1016/S0140-6736(20)30183-5. PubMed PMID: 31986264; PubMed Central PMCID: PMC7159299.
19. Wang Y, Zhao X, Feng Q, Liu L, Yao Y, Shi J. Psychological assistance during the coronavirus disease 2019 outbreak in China. *J Health Psychol.* 2020;25(6):733-7. Epub 20200417. doi: 10.1177/1359105320919177. PubMed PMID: 32301628.
20. Gao J, Zheng P, Jia Y, Chen H, Mao Y, Chen S, et al. Mental health problems and social media exposure during COVID-19 outbreak. *PLoS One.* 2020;15(4):e0231924. Epub 20200416. doi: 10.1371/journal.pone.0231924. PubMed PMID: 32298385; PubMed Central PMCID: PMC7162477.
21. Cao W, Fang Z, Hou G, Han M, Xu X, Dong J, et al. The psychological impact of the COVID-19 epidemic on college students in China. *Psychiatry Res.* 2020;287:112934. Epub 20200320. doi: 10.1016/j.psychres.2020.112934. PubMed PMID: 32229390; PubMed Central PMCID: PMC7102633.
22. Adalja AA, Toner E, Inglesby TV. Priorities for the US Health Community Responding to COVID-19. *JAMA.* 2020;323(14):1343-4. doi: 10.1001/jama.2020.3413. PubMed PMID: 32125355.
23. Milne SJ, Corbett GA, Hehir MP, Lindow SW, Mohan S, Reagu S, et al. Effects of isolation on mood and relationships in pregnant women during the covid-19 pandemic. *Eur J Obstet Gynecol Reprod Biol.* 2020;252:610-1. Epub 20200608. doi: 10.1016/j.ejogrb.2020.06.009. PubMed PMID: 32616415; PubMed Central PMCID: PMC7278652.
24. Corbett GA, Milne SJ, Hehir MP, Lindow SW, O'Connell M P. Health anxiety and behavioural changes of pregnant women during the COVID-19 pandemic. *Eur J Obstet Gynecol*

Reprod Biol. 2020;249:96-7. Epub 20200413. doi: 10.1016/j.ejogrb.2020.04.022. PubMed PMID: 32317197; PubMed Central PMCID: PMC7194619.

25. Lu W, Wang H, Lin Y, Li L. Psychological status of medical workforce during the COVID-19 pandemic: A cross-sectional study. *Psychiatry Res.* 2020;288:112936. Epub 20200404. doi: 10.1016/j.psychres.2020.112936. PubMed PMID: 32276196; PubMed Central PMCID: PMC7195354.
26. Naser Moghadasi A. Evaluation of the Level of Anxiety among Iranian Multiple Sclerosis Fellowships During the Outbreak of COVID-19. *Arch Iran Med.* 2020;23(4):283. Epub 20200401. doi: 10.34172/aim.2020.13. PubMed PMID: 32271605.
27. Yuan S, Liao Z, Huang H, Jiang B, Zhang X, Wang Y, et al. Comparison of the Indicators of Psychological Stress in the Population of Hubei Province and Non-Endemic Provinces in China During Two Weeks During the Coronavirus Disease 2019 (COVID-19) Outbreak in February 2020. *Med Sci Monit.* 2020;26:e923767. Epub 20200415. doi: 10.12659/MSM.923767. PubMed PMID: 32294078; PubMed Central PMCID: PMC7177041.
28. da Silva ML, Rocha RSB, Buheji M, Jahrami H, Cunha KDC. A systematic review of the prevalence of anxiety symptoms during coronavirus epidemics. *J Health Psychol.* 2021;26(1):115-25. Epub 20200823. doi: 10.1177/1359105320951620. PubMed PMID: 32830577.
29. Dawel A, Shou Y, Smithson M, Cherbuin N, Banfield M, Calcar AL, et al. The Effect of COVID-19 on Mental Health and Wellbeing in a Representative Sample of Australian Adults. *Front Psychiatry.* 2020;11:579985. Epub 20201006. doi: 10.3389/fpsy.2020.579985. PubMed PMID: 33132940; PubMed Central PMCID: PMC7573356.
30. Amit Aharon A, Dubovi I, Ruban A. Differences in mental health and health-related quality of life between the Israeli and Italian population during a COVID-19 quarantine. *Qual Life Res.* 2021;30(6):1675-84. Epub 20210114. doi: 10.1007/s11136-020-02746-5. PubMed PMID: 33447961; PubMed Central PMCID: PMC7808402.
31. Pieh C, Budimir S, Probst T. The effect of age, gender, income, work, and physical activity on mental health during coronavirus disease (COVID-19) lockdown in Austria. *J Psychosom Res.* 2020;136:110186. Epub 20200703. doi: 10.1016/j.jpsychores.2020.110186. PubMed PMID: 32682159; PubMed Central PMCID: PMC7832650.
32. van der Velden PG, Contino C, Das M, van Loon P, Bosmans MWG. Anxiety and depression symptoms, and lack of emotional support among the general population before and during the COVID-19 pandemic. A prospective national study on prevalence and risk factors. *J Affect Disord.* 2020;277:540-8. Epub 20200820. doi: 10.1016/j.jad.2020.08.026. PubMed PMID: 32889378; PubMed Central PMCID: PMC7438386.
33. Wang C, Pan R, Wan X, Tan Y, Xu L, Ho CS, et al. Immediate Psychological Responses and Associated Factors during the Initial Stage of the 2019 Coronavirus Disease (COVID-19) Epidemic among the General Population in China. *Int J Environ Res Public Health.* 2020;17(5).

Epub 20200306. doi: 10.3390/ijerph17051729. PubMed PMID: 32155789; PubMed Central PMCID: PMC7084952.

34. Albagmi FM, AlNujaidi HY, Al Shawan DS. Anxiety Levels Amid the COVID-19 Lockdown in Saudi Arabia. *Int J Gen Med.* 2021;14:2161-70. Epub 20210531. doi: 10.2147/IJGM.S312465. PubMed PMID: 34103971; PubMed Central PMCID: PMC8180301.
35. Canet-Juric L, Andres ML, Del Valle M, Lopez-Morales H, Poo F, Galli JI, et al. A Longitudinal Study on the Emotional Impact Cause by the COVID-19 Pandemic Quarantine on General Population. *Front Psychol.* 2020;11:565688. Epub 20200918. doi: 10.3389/fpsyg.2020.565688. PubMed PMID: 33071893; PubMed Central PMCID: PMC7531077.
36. Di Giuseppe M, Zilcha-Mano S, Prout TA, Perry JC, Orru G, Conversano C. Psychological Impact of Coronavirus Disease 2019 Among Italians During the First Week of Lockdown. *Front Psychiatry.* 2020;11:576597. Epub 20200930. doi: 10.3389/fpsyg.2020.576597. PubMed PMID: 33192713; PubMed Central PMCID: PMC7554332.
37. Li WW, Li Y, Yu H, Miller DJ, Rouen C, Yang F. Mental Health of Chinese People During the COVID-19 Pandemic: Associations With Infection Severity of Region of Residence and Filial Piety. *Front Psychol.* 2021;12:633452. Epub 20210528. doi: 10.3389/fpsyg.2021.633452. PubMed PMID: 34122223; PubMed Central PMCID: PMC8192824.
38. Zhang XR, Huang QM, Wang XM, Cheng X, Li ZH, Wang ZH, et al. Prevalence of anxiety and depression symptoms, and association with epidemic-related factors during the epidemic period of COVID-19 among 123,768 workers in China: A large cross-sectional study. *J Affect Disord.* 2020;277:495-502. Epub 20200826. doi: 10.1016/j.jad.2020.08.041. PubMed PMID: 32882506; PubMed Central PMCID: PMC7448744.
39. Wathélet M, Duhem S, Vaiva G, Baubet T, Habran E, Veerapa E, et al. Factors Associated With Mental Health Disorders Among University Students in France Confined During the COVID-19 Pandemic. *JAMA Netw Open.* 2020;3(10):e2025591. Epub 20201001. doi: 10.1001/jamanetworkopen.2020.25591. PubMed PMID: 33095252; PubMed Central PMCID: PMC7584927.
40. Halperin SJ, Henderson MN, Prenner S, Grauer JN. Prevalence of Anxiety and Depression Among Medical Students During the Covid-19 Pandemic: A Cross-Sectional Study. *J Med Educ Curric Dev.* 2021;8:2382120521991150. Epub 20210215. doi: 10.1177/2382120521991150. PubMed PMID: 33644399; PubMed Central PMCID: PMC7890732.
41. Tang F, Liang J, Zhang H, Kelifa MM, He Q, Wang P. COVID-19 related depression and anxiety among quarantined respondents. *Psychol Health.* 2021;36(2):164-78. Epub 20200622. doi: 10.1080/08870446.2020.1782410. PubMed PMID: 32567952.

42. Ni MY, Yao XI, Leung KSM, Yau C, Leung CMC, Lun P, et al. Depression and post-traumatic stress during major social unrest in Hong Kong: a 10-year prospective cohort study. *Lancet*. 2020;395(10220):273-84. Epub 20200109. doi: 10.1016/S0140-6736(19)33160-5. PubMed PMID: 31928765.
43. Kawohl W, Nordt C. COVID-19, unemployment, and suicide. *Lancet Psychiatry*. 2020;7(5):389-90. doi: 10.1016/S2215-0366(20)30141-3. PubMed PMID: 32353269; PubMed Central PMCID: PMC7185950.
44. Ruengorn C, Awiphan R, Wongpakaran N, Wongpakaran T, Nochaiwong S, Health O, et al. Association of job loss, income loss, and financial burden with adverse mental health outcomes during coronavirus disease 2019 pandemic in Thailand: A nationwide cross-sectional study. *Depress Anxiety*. 2021;38(6):648-60. Epub 20210401. doi: 10.1002/da.23155. PubMed PMID: 33793028; PubMed Central PMCID: PMC8251094.
45. Jacques-Avino C, Lopez-Jimenez T, Medina-Perucha L, de Bont J, Goncalves AQ, Duarte-Salles T, et al. Gender-based approach on the social impact and mental health in Spain during COVID-19 lockdown: a cross-sectional study. *BMJ Open*. 2020;10(11):e044617. Epub 20201124. doi: 10.1136/bmjopen-2020-044617. PubMed PMID: 33234664; PubMed Central PMCID: PMC7688440.
46. Fu W, Yan S, Zong Q, Anderson-Luxford D, Song X, Lv Z, et al. Mental health of college students during the COVID-19 epidemic in China. *J Affect Disord*. 2021;280(Pt A):7-10. Epub 20201111. doi: 10.1016/j.jad.2020.11.032. PubMed PMID: 33197782; PubMed Central PMCID: PMC7656159.
47. Sun S, Goldberg SB, Lin D, Qiao S, Operario D. Psychiatric symptoms, risk, and protective factors among university students in quarantine during the COVID-19 pandemic in China. *Global Health*. 2021;17(1):15. Epub 20210125. doi: 10.1186/s12992-021-00663-x. PubMed PMID: 33494769; PubMed Central PMCID: PMC7829620.
48. Sinawi HA, Al Balushi N, Al-Mahrouqi T, Al Ghailani A, McCall RK, Sultan A, et al. Predictors of psychological distress among the public in Oman amid coronavirus disease 2019 pandemic: a cross-sectional analytical study. *Psychol Health Med*. 2021;26(1):131-44. Epub 20201105. doi: 10.1080/13548506.2020.1842473. PubMed PMID: 33151748.
49. Warren AM, Zolfaghari K, Fresnedo M, Bennett M, Pogue J, Waddimba A, et al. Anxiety sensitivity, COVID-19 fear, and mental health: results from a United States population sample. *Cogn Behav Ther*. 2021;50(3):204-16. Epub 20210217. doi: 10.1080/16506073.2021.1874505. PubMed PMID: 33595414.
50. Wang X, Tao J, Zhu Q, Wu X, Li T, Zhao C, et al. Depression and anxiety symptoms to COVID-19 outbreak among the public, medical staff and patients during the initial phase of the pandemic: an online questionnaire survey by a WeChat Mini Program. *BMJ Open*. 2021;11(6):e046350. Epub 20210624. doi: 10.1136/bmjopen-2020-046350. PubMed PMID: 34168028; PubMed Central PMCID: PMC8228573.

51. Garcia-Fernandez L, Romero-Ferreiro V, Padilla S, David Lopez-Roldan P, Monzo-Garcia M, Rodriguez-Jimenez R. Gender differences in emotional response to the COVID-19 outbreak in Spain. *Brain Behav.* 2021;11(1):e01934. Epub 20201211. doi: 10.1002/brb3.1934. PubMed PMID: 33305891; PubMed Central PMCID: PMC7821629.
52. Bendau A, Plag J, Kunas S, Wyka S, Strohle A, Petzold MB. Longitudinal changes in anxiety and psychological distress, and associated risk and protective factors during the first three months of the COVID-19 pandemic in Germany. *Brain Behav.* 2021;11(2):e01964. Epub 20201123. doi: 10.1002/brb3.1964. PubMed PMID: 33230969; PubMed Central PMCID: PMC7744907.
53. Passavanti M, Argentieri A, Barbieri DM, Lou B, Wijayarathna K, Foroutan Mirhosseini AS, et al. The psychological impact of COVID-19 and restrictive measures in the world. *J Affect Disord.* 2021;283:36-51. Epub 20210115. doi: 10.1016/j.jad.2021.01.020. PubMed PMID: 33516085; PubMed Central PMCID: PMC7833558.
54. Fisher JR, Tran TD, Hammarberg K, Sastry J, Nguyen H, Rowe H, et al. Mental health of people in Australia in the first month of COVID-19 restrictions: a national survey. *Med J Aust.* 2020;213(10):458-64. Epub 20201026. doi: 10.5694/mja2.50831. PubMed PMID: 33107063; PubMed Central PMCID: PMC7756394.
55. Khubchandani J, Sharma S, Webb FJ, Wiblishauser MJ, Bowman SL. Post-lockdown depression and anxiety in the USA during the COVID-19 pandemic. *J Public Health (Oxf).* 2021;43(2):246-53. doi: 10.1093/pubmed/fdaa250. PubMed PMID: 33426559; PubMed Central PMCID: PMC7928742.
56. Hassannia L, Taghizadeh F, Moosazadeh M, Zarghami M, Taghizadeh H, Dooki AF, et al. Anxiety and Depression in Health Workers and General Population During COVID-19 in IRAN: A Cross-Sectional Study. *Neuropsychopharmacol Rep.* 2021;41(1):40-9. Epub 20201225. doi: 10.1002/npr2.12153. PubMed PMID: 33369264; PubMed Central PMCID: PMC8182959.
57. Serafim AP, Duraes RSS, Rocca CCA, Goncalves PD, Saffi F, Cappelozza A, et al. Exploratory study on the psychological impact of COVID-19 on the general Brazilian population. *PLoS One.* 2021;16(2):e0245868. Epub 20210203. doi: 10.1371/journal.pone.0245868. PubMed PMID: 33534820; PubMed Central PMCID: PMC7857630.
58. Fiorillo A, Sampogna G, Giallonardo V, Del Vecchio V, Luciano M, Albert U, et al. Effects of the lockdown on the mental health of the general population during the COVID-19 pandemic in Italy: Results from the COMET collaborative network. *Eur Psychiatry.* 2020;63(1):e87. Epub 20200928. doi: 10.1192/j.eurpsy.2020.89. PubMed PMID: 32981568; PubMed Central PMCID: PMC7556907.
59. Islam MS, Sujon MSH, Tasnim R, Sikder MT, Potenza MN, van Os J. Psychological responses during the COVID-19 outbreak among university students in Bangladesh. *PLoS One.* 2020;15(12):e0245083. Epub 20201231. doi: 10.1371/journal.pone.0245083. PubMed PMID: 33382862; PubMed Central PMCID: PMC7775049.

60. Odriozola-Gonzalez P, Planchuelo-Gomez A, Irurtia MJ, de Luis-Garcia R. Psychological effects of the COVID-19 outbreak and lockdown among students and workers of a Spanish university. *Psychiatry Res.* 2020;290:113108. Epub 20200519. doi: 10.1016/j.psychres.2020.113108. PubMed PMID: 32450409; PubMed Central PMCID: PMC7236679.
61. Nayan M, Uddin M, Hossain M, Alam M, Zinnia M, Haq I, et al. Comparison of the performance of machine learning-based algorithms for predicting depression and anxiety among University Students in Bangladesh: A result of the first wave of the COVID-19 pandemic. *Asian Journal of Social Health and Behavior.* 2022;5(2):75-84. doi: 10.4103/shb.shb\_38\_22.
62. Li Q, Miao Y, Zeng X, Tarimo CS, Wu C, Wu J. Prevalence and factors for anxiety during the coronavirus disease 2019 (COVID-19) epidemic among the teachers in China. *J Affect Disord.* 2020;277:153-8. Epub 20200813. doi: 10.1016/j.jad.2020.08.017. PubMed PMID: 32828002; PubMed Central PMCID: PMC7425543.
63. Ozamiz-Etxebarria N, Berasategi Santxo N, Idoiaga Mondragon N, Dosil Santamaria M. The Psychological State of Teachers During the COVID-19 Crisis: The Challenge of Returning to Face-to-Face Teaching. *Front Psychol.* 2020;11:620718. Epub 20210112. doi: 10.3389/fpsyg.2020.620718. PubMed PMID: 33510694; PubMed Central PMCID: PMC7835279.
64. Assefa ZM, Haile TG, Wazema DH, Tafese WT, Berrie FW, Beketie ED, et al. Mental Health Disorders During COVID-19 Pandemic Among Southwest Ethiopia University Students: An Institutional-Based Cross-Sectional Study. *SAGE Open Nurs.* 2021;7:23779608211064374. Epub 20211215. doi: 10.1177/23779608211064374. PubMed PMID: 34950769; PubMed Central PMCID: PMC8689431.
65. Husky MM, Kovess-Masfety V, Swendsen JD. Stress and anxiety among university students in France during Covid-19 mandatory confinement. *Compr Psychiatry.* 2020;102:152191. Epub 20200712. doi: 10.1016/j.comppsy.2020.152191. PubMed PMID: 32688023; PubMed Central PMCID: PMC7354849.
66. Biswas S, Biswas A. Anxiety level among students of different college and universities in India during lock down in connection to the COVID-19 pandemic. *Z Gesundh Wiss.* 2021:1-7. Epub 20210107. doi: 10.1007/s10389-020-01431-8. PubMed PMID: 33432293; PubMed Central PMCID: PMC7788277.
67. Savitsky B, Findling Y, Erel A, Hendel T. Anxiety and coping strategies among nursing students during the covid-19 pandemic. *Nurse Educ Pract.* 2020;46:102809. Epub 20200602. doi: 10.1016/j.nepr.2020.102809. PubMed PMID: 32679465; PubMed Central PMCID: PMC7264940.
68. Hamaideh SH, Al-Modallal H, Tanash M, Hamdan-Mansour A. Depression, anxiety and stress among undergraduate students during COVID-19 outbreak and "home-quarantine". *Nurs Open.* 2022;9(2):1423-31. Epub 20210514. doi: 10.1002/nop2.918. PubMed PMID: 33988913; PubMed Central PMCID: PMC8242644.

69. Najjuka SM, Checkwech G, Olum R, Ashaba S, Kaggwa MM. Depression, anxiety, and stress among Ugandan university students during the COVID-19 lockdown: an online survey. *Afr Health Sci.* 2021;21(4):1533-43. doi: 10.4314/ahs.v21i4.6. PubMed PMID: 35283951; PubMed Central PMCID: PMC8889827.
70. Cleofas JV, Rocha ICN. Demographic, gadget and internet profiles as determinants of disease and consequence related COVID-19 anxiety among Filipino college students. *Educ Inf Technol (Dordr).* 2021;26(6):6771-86. Epub 20210404. doi: 10.1007/s10639-021-10529-9. PubMed PMID: 33841028; PubMed Central PMCID: PMC8019527.
71. Hakami Z, Khanagar SB, Vishwanathaiah S, Hakami A, Bokhari AM, Jabali AH, et al. Psychological impact of the coronavirus disease 2019 (COVID-19) pandemic on dental students: A nationwide study. *Journal of Dental Education.* 2021;85(4):494-503. doi: <https://doi.org/10.1002/jdd.12470>.
72. Saddik B, Hussein A, Sharif-Askari FS, Kheder W, Temsah MH, Koutaich RA, et al. Increased Levels of Anxiety Among Medical and Non-Medical University Students During the COVID-19 Pandemic in the United Arab Emirates. *Risk Manag Healthc Policy.* 2020;13:2395-406. Epub 20201103. doi: 10.2147/RMHP.S273333. PubMed PMID: 33177898; PubMed Central PMCID: PMC7652570.
73. Haddad JM, Macenski C, Mosier-Mills A, Hibara A, Kester K, Schneider M, et al. The Impact of Social Media on College Mental Health During the COVID-19 Pandemic: a Multinational Review of the Existing Literature. *Curr Psychiatry Rep.* 2021;23(11):70. Epub 20211006. doi: 10.1007/s11920-021-01288-y. PubMed PMID: 34613542; PubMed Central PMCID: PMC8493361.
74. Rajkumar RP. COVID-19 and mental health: A review of the existing literature. *Asian J Psychiatr.* 2020;52:102066. Epub 20200410. doi: 10.1016/j.ajp.2020.102066. PubMed PMID: 32302935; PubMed Central PMCID: PMC7151415.
75. Mokhtari R, Moayedi S, Golitaleb M. COVID-19 pandemic and health anxiety among nurses of intensive care units. *International journal of mental health nursing.* 2020;29(6):1275.
76. Luceno-Moreno L, Talavera-Velasco B, Garcia-Albuerne Y, Martin-Garcia J. Symptoms of Posttraumatic Stress, Anxiety, Depression, Levels of Resilience and Burnout in Spanish Health Personnel during the COVID-19 Pandemic. *Int J Environ Res Public Health.* 2020;17(15). Epub 20200730. doi: 10.3390/ijerph17155514. PubMed PMID: 32751624; PubMed Central PMCID: PMC7432016.
77. Bao Y, Sun Y, Meng S, Shi J, Lu L. 2019-nCoV epidemic: address mental health care to empower society. *Lancet.* 2020;395(10224):e37-e8. Epub 20200207. doi: 10.1016/S0140-6736(20)30309-3. PubMed PMID: 32043982; PubMed Central PMCID: PMC7133594.
78. Salari N, Hosseini-Far A, Jalali R, Vaisi-Raygani A, Rasoulpoor S, Mohammadi M, et al. Prevalence of stress, anxiety, depression among the general population during the COVID-19 pandemic: a systematic review and meta-analysis. *Globalization and health.* 2020;16(1):1-11.

79. Ryu S, Chun BC, Korean Society of Epidemiology -nCo VTFT. An interim review of the epidemiological characteristics of 2019 novel coronavirus. *Epidemiol Health*. 2020;42:e2020006. Epub 20200206. doi: 10.4178/epih.e2020006. PubMed PMID: 32023775; PubMed Central PMCID: PMC7011107.
80. Chen Q, Liang M, Li Y, Guo J, Fei D, Wang L, et al. Mental health care for medical staff in China during the COVID-19 outbreak. *Lancet Psychiatry*. 2020;7(4):e15-e6. Epub 20200219. doi: 10.1016/S2215-0366(20)30078-X. PubMed PMID: 32085839; PubMed Central PMCID: PMC7129426.
81. Pappa S, Ntella V, Giannakas T, Giannakoulis VG, Papoutsis E, Katsaounou P. Prevalence of depression, anxiety, and insomnia among healthcare workers during the COVID-19 pandemic: A systematic review and meta-analysis. *Brain Behav Immun*. 2020;88:901-7. Epub 20200508. doi: 10.1016/j.bbi.2020.05.026. PubMed PMID: 32437915; PubMed Central PMCID: PMC7206431.
82. Shen M, Xu H, Fu J, Wang T, Fu Z, Zhao X, et al. Investigation of anxiety levels of 1637 healthcare workers during the epidemic of COVID-19. *PLoS One*. 2020;15(12):e0243890. Epub 20201222. doi: 10.1371/journal.pone.0243890. PubMed PMID: 33351814; PubMed Central PMCID: PMC7755185.
83. Lai J, Ma S, Wang Y, Cai Z, Hu J, Wei N, et al. Factors Associated With Mental Health Outcomes Among Health Care Workers Exposed to Coronavirus Disease 2019. *JAMA Netw Open*. 2020;3(3):e203976. Epub 20200302. doi: 10.1001/jamanetworkopen.2020.3976. PubMed PMID: 32202646; PubMed Central PMCID: PMC7090843.
84. Antonijevic J, Binic I, Zikic O, Manojlovic S, Tosic-Golubovic S, Popovic N. Mental health of medical personnel during the COVID-19 pandemic. *Brain Behav*. 2020;10(12):e01881. Epub 20201017. doi: 10.1002/brb3.1881. PubMed PMID: 33070475; PubMed Central PMCID: PMC7645856.
85. Hacimusalar Y, Kahve AC, Yasar AB, Aydin MS. Anxiety and hopelessness levels in COVID-19 pandemic: A comparative study of healthcare professionals and other community sample in Turkey. *J Psychiatr Res*. 2020;129:181-8. Epub 20200721. doi: 10.1016/j.jpsychires.2020.07.024. PubMed PMID: 32758711; PubMed Central PMCID: PMC7372275.
86. Giardino DL, Huck-Iriart C, Riddick M, Garay A. The endless quarantine: the impact of the COVID-19 outbreak on healthcare workers after three months of mandatory social isolation in Argentina. *Sleep Med*. 2020;76:16-25. Epub 20200925. doi: 10.1016/j.sleep.2020.09.022. PubMed PMID: 33059247; PubMed Central PMCID: PMC7518855.
87. Chew NWS, Ngiam JN, Tan BY, Tham SM, Tan CY, Jing M, et al. Asian-Pacific perspective on the psychological well-being of healthcare workers during the evolution of the COVID-19 pandemic. *BJPsych Open*. 2020;6(6):e116. Epub 20201008. doi: 10.1192/bjo.2020.98. PubMed PMID: 33028449; PubMed Central PMCID: PMC7542327.

88. Hennein R, Mew EJ, Lowe SR. Socio-ecological predictors of mental health outcomes among healthcare workers during the COVID-19 pandemic in the United States. *PLoS One*. 2021;16(2):e0246602. Epub 20210205. doi: 10.1371/journal.pone.0246602. PubMed PMID: 33544761; PubMed Central PMCID: PMC7864435.
89. Gainer DM, Nahhas RW, Bhatt NV, Merrill A, McCormack J. Association Between Proportion of Workday Treating COVID-19 and Depression, Anxiety, and PTSD Outcomes in US Physicians. *J Occup Environ Med*. 2021;63(2):89-97. doi: 10.1097/JOM.0000000000002086. PubMed PMID: 33201021; PubMed Central PMCID: PMC7864594.
90. Lai J, Ma S, Wang Y, Cai Z, Hu J, Wei N, et al. Factors associated with mental health outcomes among health care workers exposed to coronavirus disease 2019. *JAMA network open*. 2020;3(3):e203976-e.
91. Setiawati Y, Wahyuhadi J, Joestandari F, Maramis MM, Atika A. Anxiety and Resilience of Healthcare Workers During COVID-19 Pandemic in Indonesia. *J Multidiscip Healthc*. 2021;14:1-8. Epub 20210106. doi: 10.2147/JMDH.S276655. PubMed PMID: 33442258; PubMed Central PMCID: PMC7797347.
92. Santabarbara J, Lasheras I, Lipnicki DM, Bueno-Notivol J, Perez-Moreno M, Lopez-Anton R, et al. Prevalence of anxiety in the COVID-19 pandemic: An updated meta-analysis of community-based studies. *Prog Neuropsychopharmacol Biol Psychiatry*. 2021;109:110207. Epub 20201215. doi: 10.1016/j.pnpbp.2020.110207. PubMed PMID: 33338558; PubMed Central PMCID: PMC7834650.
93. Denning M, Goh ET, Tan B, Kanneganti A, Almonte M, Scott A, et al. Determinants of burnout and other aspects of psychological well-being in healthcare workers during the Covid-19 pandemic: A multinational cross-sectional study. *PLoS One*. 2021;16(4):e0238666. Epub 20210416. doi: 10.1371/journal.pone.0238666. PubMed PMID: 33861739; PubMed Central PMCID: PMC8051812.
94. Jia R, Ayling K, Chalder T, Massey A, Broadbent E, Coupland C, et al. Mental health in the UK during the COVID-19 pandemic: cross-sectional analyses from a community cohort study. *BMJ Open*. 2020;10(9):e040620. Epub 20200915. doi: 10.1136/bmjopen-2020-040620. PubMed PMID: 32933965; PubMed Central PMCID: PMC7493070.
95. Planchuelo-Gomez A, Odriozola-Gonzalez P, Iurtia MJ, de Luis-Garcia R. Longitudinal evaluation of the psychological impact of the COVID-19 crisis in Spain. *J Affect Disord*. 2020;277:842-9. Epub 20200907. doi: 10.1016/j.jad.2020.09.018. PubMed PMID: 33065825; PubMed Central PMCID: PMC7476580.
96. Cai Q, Feng H, Huang J, Wang M, Wang Q, Lu X, et al. The mental health of frontline and non-frontline medical workers during the coronavirus disease 2019 (COVID-19) outbreak in China: A case-control study. *J Affect Disord*. 2020;275:210-5. Epub 20200702. doi: 10.1016/j.jad.2020.06.031. PubMed PMID: 32734910; PubMed Central PMCID: PMC7329671.

97. Chen L, Zhao H, Razin D, Song T, Wu Y, Ma X, et al. Anxiety levels during a second local COVID-19 pandemic breakout among quarantined people: A cross sectional survey in China. *J Psychiatr Res.* 2021;135:37-46. Epub 20210105. doi: 10.1016/j.jpsychires.2020.12.067. PubMed PMID: 33445059; PubMed Central PMCID: PMC7783475.
98. Nandi T. Importance of sufficient time-points for efficient pharmacokinetic (PK) compartmental modeling. *International Journal of Applied Pharmaceutics.* 2023;15(1):87-92.
99. Ghosh R, Nayan MIH, Mitu MM, Nandi T. Common Approaches of Cytochrome P450 (CYP) Induction Assays. *International Blood Research & Reviews.* 2023;14(1):6-14.
100. Qian M, Wu Q, Wu P, Hou Z, Liang Y, Cowling BJ, et al. Anxiety levels, precautionary behaviours and public perceptions during the early phase of the COVID-19 outbreak in China: a population-based cross-sectional survey. *BMJ Open.* 2020;10(10):e040910. Epub 20201008. doi: 10.1136/bmjopen-2020-040910. PubMed PMID: 33033099; PubMed Central PMCID: PMC7545627.
101. Tasnim R, Islam MS, Sujon MSH, Sikder MT, Potenza MN. Suicidal ideation among Bangladeshi university students early during the COVID-19 pandemic: Prevalence estimates and correlates. *Child Youth Serv Rev.* 2020;119:105703. Epub 20201110. doi: 10.1016/j.chilyouth.2020.105703. PubMed PMID: 33204046; PubMed Central PMCID: PMC7654299.
102. Rizun M, Strzelecki A. Students' Acceptance of the COVID-19 Impact on Shifting Higher Education to Distance Learning in Poland. *Int J Environ Res Public Health.* 2020;17(18). Epub 20200905. doi: 10.3390/ijerph17186468. PubMed PMID: 32899478; PubMed Central PMCID: PMC7558862.