

Original Research Article

Effects of Methanolic Leaves Extracts of Botanicals on the Toxicity and Enzymatic Parameters of *Callosobruchus maculatus* Fab. Larvae (Coleoptera: Bruchidae)

ABSTRACT

A study was conducted in the laboratory to evaluate the effect of leaves extract of three botanicals on the biochemical parameters of insect pest of stored beans, *Callosobruchus maculatus*. The results of the qualitative phytochemicals revealed the presence of different metabolites such as tannins, saponins, cardiac glycosides, phenolic compounds, flavonoids, alkaloids etc. The effect of methanolic leaves extracts of *Lasianthera africana*, *Hippocratea africana* and *Uvaria chamae* on the toxicity and biochemical parameters after treatment against *C. maculatus* was evaluated. The result shows that treatment of the insects with these extracts revealed a significant mortality in *C. maculatus*. The enzymatic activities of the digestive enzymes such as amylase and invertase revealed a significant inhibition in the treated larvae. The transaminase enzymes (AST and ALT) was found to be reduced in the insect after treatment (33.2 and 42.6) while the result of the phosphatase enzyme activity (ACP and ALP) showed a potent inhibitory effect on both enzymes which was more pronounced in ALP than ACP with percentage inhibition of 75.8 and 31.8 respectively.

Keywords: *Callosobruchus maculatus*, *Lasianthera africana*, *Hippocratea africana*, *Uvaria chamae*, Enzymatic, Toxicity

1. Introduction

Indiscriminate use of synthetic chemical pesticides to control pests has led to the development of insect's resistance and also affected non target organisms. Hence, an environmental-friendly alternative is needed [1]. Recently, an intensive research has been carried out to control agricultural pests by using natural insecticides of plant origin to decrease hazards in the environment. Plant based products had been used to control different pests by farmers atleast for two millennia [2]. Because of the abundance of phytochemicals in plant, the extracts of various plant species with different secondary metabolite have been used to control pests of various order including the Coleopterans [3].

Uvaria chamae leaves are alternately arranged, the leaf has simple structures, lanceolate in shape with entire lamina and net veined. Leaves are stipulate, leaf apex acuminate and the leaf vestiture is glabrous [4]. *Lasianthera africana* is monospecific genus. It is a perennial glabrous shrub that reaches a height of 61 – 136 cm. [5]. *Hippocratea africana* is a green forest perennial climber without hairs (glabrous), reproducing from seeds [6].

It is a cosmopolitan post-harvest pest that causes quantitative and qualitative losses of stored grains in West Africa and its infestation begins in the field at low levels and increases in stored population [7]. It is used as a model organism for research and education due to its rapid development in storage [8]. The adult weevils are short, stout-bodied beetles about 4.76 mm long with the wing covers shortened and not covering the tip of the abdomen [9]. Their antennae are

usually conspicuous and the body is narrow toward the front [10]. This weevil cowpea lacks the 'snout' of a real weevil. It is reddish-brown overall, with two central black spots marked by black and grey elytra. The last abdomen segment extends below the short elytra, as well as having two black spots [11].

2. Materials and Methods

2.1 Collections and Identification of Plant Materials

The fresh leaves of *Uvaria chamae*, *Lasianthera africana* and *Hippocratea africana* were obtained from Faculty of Pharmacy Medicinal Farm of University of Uyo, Akwa Ibom State and validated by a taxonomist in the Department of Botany and Ecological Studies, University of Uyo. Voucher specimens with number: UUH/3687, UUH/3688 and UUH/3689 was deposited in their herbarium for further referencing.

2.2 Rearing of Test Organisms

To provide comparable age weevils for the experiment, *C. maculatus* cultures were established. A total of five (5 kg) bean seeds were purchased and cleaned to remove any seeds with visible damage. To prevent potential field infestation, the clean seeds were kept in a sealed container in the fridge at 4°C for a month. Seeds were placed in soft bags and stored at room temperature for two weeks. The beetle is sexually dimorphic, and it easily distinguishes males from females. Sometimes the females are larger than the males, overall the females are darker while the males are brunette. The plate which covers the end of the abdomen is large and dark in female on the sides and smaller in male without the dark areas [12]. The insects were cultured on clean seeds, with 50 weevils per 200 g of seeds in each jar. To allow airing and prevent weevil escape, the jar was covered with muslin cloth and secured with a rubber band and kept at room temperature. All parent weevils were removed from each jar seven days after oviposition. The jars were placed in an insect rearing cage kept in the Entomology Laboratory, Department of Animal and Environmental Biology, University of Uyo, Uyo. Newly emerged, two day-old insects was used for the experiment.

2.3 Preparation of plant powder and extract

After collection, the plant leaves were washed and chopped into pieces and room dried to a constant weight. Using a power-driven blender (Braun Multiquick Immersion Hand Blender, B White Mixer MR 5550CA, Germany), the dry plants were melded into fine powder and then kept in an airtight container pending use. The crude leaf extracts were then prepared using standard procedures as outlined by Santana [13]; Mukhtar and Huda [14] and Fatope [15]. This involved soaking 50 g of the powder for 48 – 72 hours at room temperature in 95 percent methanol. This was followed by filtrate evaporation using a rotatory evaporator to obtain the crude extract.

2.4 Phytochemical Analysis of the Plants

The initial phytochemical screening of the different plants was carried out in Pharmacognosy Laboratory of University of Uyo, Akwa Ibom State using the standard procedures as described by Prashant [16]; Kokate [17]; Evans [18] and Harbone [19].

2.5 Determination of LC₅₀

The acute toxicity (LC₅₀) of the extract of the extract types used in this study were established using the method of Ousman [20] and Abbott [21] where LC₅₀ of the extract types were obtained

during preliminary studies until a concentration that will have effects 50% of the tests after 24 hours was obtained.

2.6 Toxicity Experiment

Healthy grains were kept in a freezer for one week to control hidden infestation, then left to equilibrate to room temperature. Twenty (20 g) of the grains was measured into 200 ml plastic cups and treated with different unitary and binary formulations of the botanicals at 5 and 10 %. A conventional insecticide at 0.25 g concentration was used as standard, while the experimental control was set up without any treatment. An hour after the addition of the botanicals, 10 pairs of sexed adult *C. maculatus* of two days old were introduced into treated and untreated grains. The plastic cups were covered with white muslin cloth held in place with rubber bands. The experiment was laid out using a completely randomized design and replicated four times. Mortality was recorded after 24, 48 and 72 hours post treatment. Insects were considered dead on failure to respond to three probes using a blunt dissecting probe [22, 23,].

2.7 Determination of the Digestive Enzyme: Amylase and invertase activities were assayed calorimetrically according to the methods described by Ishayaa and Swirski [24]. The activities of the enzymes were based on the digestion of starch and sucrose, respectively.

The free aldehyde groups of glucose formed after starch and sucrose digestion were allowed to react with 3, 5-dinitro salicylic acid reagent. The reduced dinitro salicylic acid is measured spectrophotometrically at 550 nm.

2.8 Determination of Transaminase Enzymes (AST and ALT): AST (aspartate aminotransaminase) and ALT (alanine aminotransaminase) were determined calorimetrically according to the method of Adel *et al.* [25] and Reitman and Frankle [26]. AST transfers the amino group from L-aspartate to Ketoglutaric acid, producing a new amino acid (L-glutamic) and a new keto acid (Oxaloacetate). ALT transfers the amino group from D, L-alanine to (-ketoglutaric acid), resulting in L-glutamic acid and pyruvic acid. The oxaloacetate and pyruvate formed from both reactions react with 2, 4-dinitrophenylhydrazine, forming oxaloacetate or pyruvate hydrazone which in alkaline medium form a brown colour, which can be measured spectrophotometrically at 546 nm.

2.9 Phosphatase Enzymes: Acid phosphatase (ACP) and alkaline phosphatase (ALP) were determined according to the method described by Powell and Smith [27]. In this method, the phenol released by enzymatic hydrolysis of disodium phenyl phosphate was measured spectrophotometrically.

2.10 Data Analysis

Data collected on the toxicity experiment were analyzed using percentages, while that of biochemical assay were subjected to univariate analysis of variance (ANOVA). Significant means were separated using New Duncan Multiple Range Test. T- test was used to compare the differences between the enzymes at the same treatment group. Significant means were accepted at $p < 0.05$ and presented as means \pm standard error of mean. All analysis were done using Statistical Packages for Social Sciences (SPSS) version, 23.0 (IBM Corporation, Armonk USA).

3. Results

3.1 Phytochemical Composition of the plants extracts

The results of the qualitative phytochemicals revealed the presence of different metabolites as shown in Table 1. Steroids and terpenes were strongly present in *H. africana* but *U. chamae* and *L. africana* had it in trace. Anthraquinones were in trace in *U. chamae* but absent in both *L. africana* and *H. africana*. Cardiac glycosides were strongly present in *U. chamae* but moderately present in both *L. africana* and *H. africana*. Saponins was detected to be strongly present in *L.*

africana but moderately present in both *H. africana* and *U. chamae*. Tannins and Phenols were strongly present in *L. africana* but moderately present in both *U. chamae* and *H. africana*. Flavonoids were strongly present in both *U. chamae* and *H. africana* but moderately present in *L. africana*. Alkaloids were strongly present in *U. chamae* but moderately present in both *H. africana* and *L. africana*. Also, phlobatannins was moderately present in both *U. chamae* and *H. africana* but absent in *L. africana*.

Table 1: Qualitative phytochemical analysis of the different extracts

	<i>U. chamae</i>	<i>L. africana</i>	<i>H. africana</i>	Test
Anthraquinones	+	-	-	Bortrager
Steroids/terpenes	++	+	+++	Liebermann-Burchard
Cardiac glycoside	+++	++	++	Keller-kiliani, Salkowski
Saponin	++	+++	++	Frothing, Fehling solution, Na ₂ CO ₃
Tannins and Phenols	++	+++	++	Ferric Chloride, Pb acetate
Flavonoids	+++	++	+++	NaOH, Mayer, Wagner
Alkaloids	+++	++	++	NaOH, Shinda
Phlobatannins	++	-	++	Dragendoff, Mayer, Wagner

+++ = Strongly present; ++ = moderately present; += trace; - = absent

Results on the toxicity assessment of both *Callosobruchus maculatus* for 96 hours after the application of the unitary and binary formulations of different plant extracts at 5 and 10 % concentrations are shown on Tables 2, and figure 1. Significant difference ($p < 0.5$) in mortality was observed among different treatments depending on the type of extracts combinations, concentration and time after treatment. Significantly higher mortality was recorded under binary formulations as compared to unitary formulations at both level of treatment. The highest mean mortality of both *C. maculatus* was recorded as the result of the synergy of *U. chamae* + *L. africana*, *U. chamae* + *H. africana* and *L. africana* + *H. africana* treatment at both concentrations. The result also showed that binary formulation where *U. chamae* was added has a highest percent mortality of 90%. The toxicity effect of the binary formulations was comparable with the synthetic insecticide (Aluminium Phosphide). Increased mortality of both weevils was observed after the treatment with higher concentration. Among the unitary formulations, the highest mortality was recorded at higher concentration (10%) after treatment with *U. chamae* followed by *H. africana* and *L. africana*. There was no significant difference in mortality in both unitary and binary formulations.

Table 2: Adult mortality of *Callosobruchus maculatus* (% mean \pm SD) on maize seeds admixed with unitary and binary formulations (5% and 10%) of different botanical extracts

Conc. (%)	Groups	Duration (Day)			
		7	14	21	28
10	<i>U. chamae</i>	0.75 ± 0.50 ^{ab1}	1.75 ± 0.50 ^{b2}	2.75 ± 0.50 ^{b2}	4.00 ± 0.00 ^{b3}
	<i>L. africana</i>	0.25 ± 0.50 ^{b1}	0.75 ± 0.50 ^{bc1}	1.25 ± 0.50 ^{b12}	2.00 ± 0.00 ^{b2}
	<i>H. africana</i>	0.50 ± 0.58 ^{ab1}	1.25 ± 0.96 ^{b12}	2.00 ± 0.82 ^{b2}	3.00 ± 0.82 ^{b2}
	<i>U. chamae</i> + <i>L. africana</i>	1.00 ± 0.00 ^{ab1}	2.25 ± 0.50 ^{b2}	3.25 ± 0.50 ^{b2}	4.50 ± 1.00 ^{b3}
	<i>U. chamae</i> + <i>H. africana</i>	1.00 ± 0.00 ^{ab1}	2.25 ± 0.50 ^{b2}	3.50 ± 0.58 ^{b3}	5.00 ± 0.82 ^{b4}
	<i>L. africana</i> + <i>H. africana</i>	1.00 ± 0.00 ^{ab1}	1.75 ± 0.50 ^{b2}	2.75 ± 0.50 ^{b2}	4.00 ± 0.00 ^{b3}
5	<i>U. chamae</i>	0.50 ± 0.58 ^{b1}	1.25 ± 0.50 ^{b1}	1.75 ± 0.96 ^{b1}	2.50 ± 1.29 ^{b1}
	<i>L. africana</i>	0.50 ± 0.58 ^{b1}	0.75 ± 0.96 ^{bc1}	1.00 ± 0.82 ^{bc1}	1.50 ± 1.29 ^{b1}
	<i>H. africana</i>	0.75 ± 0.50 ^{b1}	1.00 ± 0.82 ^{bc1}	1.50 ± 0.58 ^{b1}	2.50 ± 0.58 ^{b1}
	<i>U. chamae</i> + <i>L. africana</i>	0.50 ± 0.58 ^{b2}	1.00 ± 0.82 ^{bc2}	1.75 ± 0.96 ^{b12}	2.75 ± 0.96 ^{b2}
	<i>U. chamae</i> + <i>H. africana</i>	0.25 ± 0.50 ^{b1}	1.00 ± 0.82 ^{bc1}	1.50 ± 1.29 ^{b1}	2.50 ± 1.29 ^{b1}
	<i>L. africana</i> + <i>H. africana</i>	0.75 ± 0.50 ^{b1}	1.25 ± 0.50 ^{b1}	1.75 ± 0.96 ^{b1}	2.50 ± 1.00 ^{b1}
	Standard control	2.00 ± 0.00 ^{a1}	4.00 ± 0.00 ^{a2}	6.00 ± 0.00 ^{a3}	8.00 ± 0.00 ^{a4}
	Experimental control	0.00 ± 0.00 ^{b1}	0.00 ± 0.00 ^{c1}	0.00 ± 0.00 ^{c1}	0.00 ± 0.00 ^{c1}

Values as mean ± standard deviation. Values with different alphabet superscripts along a column were significantly different; while values with different numeric superscripts across a row were significantly different ($p < 0.05$).




Figure 1: Cumulative mortality effects of both unitary and binary formulations of different extract on adult of *Callosobruchus maculatus* applied at dosage rates of 5% (light blue) and 10% (brown) on maize seeds

Digestive Enzymes: The results of amylase and invertase enzymes activity revealed a significant decrease ($p < 0.05$) in the larvae treated with the *Lasianthera africana*, *Hippocratea africana* and *Uvaria chamae* thus: 47.70 ± 5.3 ; 43.23 ± 3.5 and 40.35 ± 2.5 $\mu\text{m}/\text{min}/\text{g}$ tissue for *C. maculatus* respectively when compared to controls (78.97 ± 1.3 $\mu\text{m}/\text{min}/\text{g}$) tissue and invertases with 46.27 ± 3.9 ; 42.85 ± 3.8 and 38.75 ± 3.9 $\mu\text{m}/\text{min}/\text{g}$ tissue for *C. maculatus* respectively when compared with controls. These values represent 51.3 % (amylase) inhibition and 54.7% (invertase) respectively as shown in the table. These enzymes are secreted to play role in digestion and utilization of starch and sucrose, so the impairment of these substrates availability might have inhibited the digestive enzymes activity in the tested weevil. It therefore shows that *U. chamae*, *L. africana* and *H. africana* delays feeding initiation and the actual movement of food bolus through the digestive tract.

Transaminase enzymes activities (AST and ALT): It is observed from Table 2 that treatment with different plant extracts reduces the activity of aspartate aminotransaminase (AST) and alanine aminotransaminase (ALT) as the percentages inhibition were 33.2% and 42.6% for *C. maculatus* respectively when compared to the controls. The mean average of ALT concentrations of different extract were 4.35 ± 0.25 , 4.23 ± 0.22 and 4.15 ± 0.11 in the case of treatment with *Lasianthera africana*, *Hippocratea Africana* and *Uvaria chamae* respectively compared to 6.15 ± 0.05 of the control. There was a significant difference between the treated and untreated larvae.

The activity of AST are found to be decreasing when treated with extracts of *L. africana*, *H. africana* and *U. chamae* in this order as compared to control. The average values of AST activity in *Sitophilus zeamais* were 3.0 ± 0.25 , 3.1 ± 0.14 and 2.9 ± 0.34 when compared to the control of 5.4 ± 0.39 . There was a significant difference between the treated and untreated larvae. Therefore, it is clear that the observed reduction in the enzymatic activities of AST was higher than that of ALT.

Phosphatase enzymes (ACP and ALP): The data obtained as shown in Tables 2 and 3 showed the effect of the tested plant extracts on the enzymatic activity of acid and alkaline phosphatases (ACP and ALP). The result showed a potent inhibitory effect of both enzymes which was 31.8% and 75.8% for ACP and ALP respectively on treated weevil when compared to the control. The mean average of ACP activities of different extract were 3.04 ± 0.16 , 3.04 ± 0.13 and 4.01 ± 0.25 U/100mg in the case of treatment with *Lasianthera africana*, *Hippocratea africana* and *Uvaria chamae* respectively compared to 4.2 ± 0.02 U/100mg of the control. There was a significant difference between the treated and untreated larvae.

The mean average of ALP activities of different extract were 72.6 ± 9.8 , 70.5 ± 8.7 and 68.4 ± 6.7 U/100mg in the case of treatment with *L. africana*, *H. africana* and *U. chamae* respectively compared to 245.8 ± 8.5 U/100mg of the controls in the weevil. There was a significant increase in ALP than ACP in the treated and untreated larvae.

Table 3: Enzymatic activity of larvae of *C. maculatus* treated with different extracts.

Enzymes Sample	Enzymes activities				
	<i>L. africana</i>	<i>H. africana</i>	<i>U. chamae</i>	Control	% inhibition
Digestive enzymes ($\mu\text{m}/\text{min}/\text{g}$ tissues)					
Amylase	47.7 ± 5.3	43.23 ± 3.5	40.35 ± 2.5	78.77 ± 1.3	51.3
Invertase	$46.27 \pm 3.9^{**}$	42.85 ± 3.8	38.75 ± 3.9	72.32 ± 4.4	54.7
Transaminase enzymes($\mu\text{m}/\text{min}/\text{g}$ tissues)					

AST	3.0 ± 0.25**	3.1 ± 0.14	2.9 ± 0.38	5.4 ± 0.39	33.2
ALT	4.35 ± 0.25**	4.32 ± 0.22	4.15 ± 0.11	6.15 ± 0.5	42.6
Phosphate enzymes(µm/min/g tissues)					
Acid Phosphate	3.04 ± 0.16	3.04 ± 0.13	4.01 ± 0.25	4.02 ± 0.2	31.8
Alkaline Phosphate	72.9 ± 9.8	70.5 ± 8.7	68.4 ± 6.7	245.8 ± 8.5	75.8

Discussions

The qualitative phytochemical screening of all the plants extract (*Uvaria chamae*, *Hippocratea africana* and *Lasianthera africana*) revealed the presence of alkaloids, saponins, tannins, flavonoids, phenols and cardiac glycosides. This result was in agreement with the report of [28, 29, 30] who carried out a phytochemical screening of some methanolic plants extracts and found tannins, flavonoids, alkaloids and saponins to be the most abundant phytochemical present. Anthraquinone was not present in this study, while Phlobotannins was moderately present. The report of this study was in agreement with the findings of [31, 32, 33] who carried out the phytochemical screening of extract of *L. africana* and *H. africana* and reported no trace of anthraquinone and moderately presence of Phlobatannins. The strong presence of cardiac glycosides observed in this study was in agreement with the findings of [34, 32, 30] who observed heavy presence of the same metabolites when carrying out the phytochemical screening of *L. africana* and *H. africana* but disagreed with the findings of Rajeswari *et al.* [35] who observed no trace of cardiac glycosides when screening the extract of *H. africana*.

The findings showed that ALT and AST, reared on the tested extracts in the 4th instar larvae, were less than in untreated control larvae. From the present findings it is very clear that there was inhibition of AST on the treated seed grains in the 4th instar larvae. These results agreed with [36] who reported that as far as the activity of transaminase is concerned, pyrethroid and Neem formulation compounds were found to have inhibitory effects and the decrease of AST was more than in ALT. Hassen [37] denoted that the activity of tissue-specific enzymes was used to diagnose the harm caused by chemical toxicity to particular organs and tissues. The two most diagnostic-potential enzymes are aspartate aminotransferase (AST) and alanine aminotransferase (ALT). Since transaminase enzymes help to produce energy, ALT and AST assist in the transition of amino groups and play an important role in the process of the Krebs or the process of high energy output, in other words, the amino acid, lipid, and carbohydrate metabolism [38, 39]. Since AST and ALT are essential anaplerotic enzymes that provide oxaloacetate and pyruvate as critical precursors of Krebs's cycle respectively, inhibition of these enzymes caused impairment of this process that could affect the normal reproduction and growth rate of the treated insects. It is in accordance with [40] who concluded that plant extracts induced inhibition of AST and ALT enzymes, thus decreasing *Aphis craccivora's* reproductive capacity. Therefore, the use of plant extracts, which have been found to suppress these enzymes, can help combat pests of insects. The results showed that alkaline phosphatase activity in treated larvae was lower than in control when using *L. africana*, while its activities increases when using *H. africana* and *U. chamae*

extract. The resulted increase in this study was in agreement with El-Gindi [41] who found that topical treatment of *Parasarcophaga argyrostoma* with juvenile hormone, pyriproxyfen at 1 % leads to highly significant increases in ALP activity. Acid phosphatase activity in the treated larvae was found to be lower than that in control. This result was in agreement with the findings of Mostafa [42, 43] who recorded significant reduction in the acid phosphatases activity at all times intervals when treatment having formulation of plant extracts (*Margason*) were administered to 4th and 6th instar larvae of *Spodoptera littoralis*. Since both enzymes are closely related to insect development, nutrition, egg maturation and metamorphosis, their inhibition could affect the transport of nutrients which in turn may impair the normal development of the insect. This was in agreement with the findings of [44, 45, 46] who reported that botanical extracts caused a reduction in ACP and ALP activities. The extracts was found to have more effects on the weevils as juvenile hormone on larvae. The findings here are in accordance with [47] who found that s-alp (Soluble alkaline phosphatase) activity was increased in all tissue whereas m-alp (membrane alkaline phosphatase) was increased in the midgut and hindgut by juvenile hormone analog (JHA) treatment, and also the larval duration was increased. This is also in agreement with [48] who observed that the plant oils may have a regulatory impact on juvenile or insect development. The rise in alkaline phosphatase may also be due to the *U. chamae*'s juvenile hormone effect because juvenile hormone contributes to increased alkaline phosphatase. The study has therefore, identified *C. maculatus* economic roles as serious threat to beans production in Nigeria.

CONCLUSION

The plants used in controlling the weevil, *Callosobruchus maculatus* are harmless to mammals at least at the dosage being reported and so efforts should be intensified towards their cultivation, packaging and application on a large scale as botanical insecticides.

References

1. Packiam SM, Baskar K, Ignacimuthu S. Insecticidal and histopathological effects of botanical formulations against *Helicoverpa armigera* (Hub.) (Lepidoptera: Noctuidae). *Journal of Agricultural Technology*, 2013;9(3): 553-563.
2. Thacker D, *Global Neem Update 2: 1*. University Press, 2002 London.
3. Muthu C, Baskar K, Kingsley S, Ignacimuthu S. Bioefficacy of *Clerodendrum phlomidis* Linn. F. and *Fleuggea leucopyrus* (Koen.) Willd. against *Earias vitelli* Fab. *Journal of Entomology*, 2013;9:332-342
4. Bongers F, Parren PME, Traore D. *Forest Climbing Plants of West Africa: Diversity, Ecology and Management*. CAB International, 2005 New Delhi.
5. Bassey ME, Etuk UI, Ekpo JU. Morphological diversity in the macrophyte genus *Lasianthera* (Icacinaceae) and the taxonomic implications. *Living System for Sustainable Development*, 2014;2: 1-5.
6. Dalziel JM, *Useful Plants of West Tropical Africa*. Crown Agents for Overseas Government, 1956 London.

7. Ntoukwam G, Murdock LL, Shode RE. Managing insect pests of cowpea in storage. Midcourse Research Meeting, 2006; Senegal.
<http://www.conr.msu.edu/oerseas/cowpea/proceal> Accessed November, 2013.
8. Profit M, Bruchid Research at Royal Holloway. University of London, 1997 London.
9. Houseman RM, Insect Pest of Stored Product. Division of Plant Science Entomology, University of Missouri Extension Service, University of Missouri, 2006 USA.
10. Dress BM, John J. Field Guide to Texas Insects. Gulf Publishing Company, 1999 Texas, USA.
11. Paranagama PA, Adhikari AACK, Abeywickrama KP, Bandara KANP. Toxicity and repellent activity of *Cymbopogon citratus* (DC) Stapf and *Murraya koenigii* Sprang against *Callosobruchus maculatus* (F.) (Coleoptera, Bruchidae). Tropical Agricultural Research and Extension, 2002;5(1-2): 22-28.
12. Kapila R, Agarwal HC. Biology of an egg parasite of *Callosobruchus maculatus* (Fab.) (Coleoptera: Bruchidae). *Journal of Stored Products Research*, 1995;31(4): 335-341.
13. Santana, PM, Miranda M, Payrol JA, Silva M, Hernandez V, Peralta E. Gas Chromatography- Mass Spectrometry Study from the Leaves Fractions Obtained from *Vernonanthura patens* (Kunth) H. Rob. *International Journal of Organic Chemistry*, 2013 3(2):32968. <http://dx.doi.org/10.4236/ijoc.2013.32011>
14. Mukhtar, M. D. and Huda, M. (2005). Prevalence of *Tinea capitis* in primary school and sensitivity of etiological agents of *Pisti astratiotes* extracts. *Nigerian Journal of Microbiology*, 19(1-2): 412-419.
15. Fatope MO, Ibrahim H, Takeda Y. Screening of higher plants reputed as pesticides using the brine shrimp lethality assay. *International Journal of Pharmacognosy*, 1999;31(4): 250-254, <https://doi.org/10.3109/13880209309.082949>
16. Prashant, T., Bimlesh, K., Mandeep, K., Gurpreet, K. and Harleen, K. Phytochemical screening and extraction: a review. *Internationale Pharmaceutica Scientia*, 2011; 1(1): 98-106.
17. Kokate CK, Purohit AP, Gokhale SB. *Pharmacognosy*. 40th Edition, Nirali Prakashan, 2008; India.
18. Evans WC. *Trease and Evans' Pharmacognosy*. 15th Edition, Elsevier, A Division of Reed Elsevier India Pvt. Limited, 2002; India.
19. Harborne JB. *Phytochemical Methods*. 2nd Edition, Chapman and Hall, 1984; New York.
20. Ousman A, Ngassoum MB, Essia-Ngang JJ, Ngamo LST, Ndjouvenkwu R. Insecticidal activity of spicy plant oils against *Sitophilus zeamais* in stored maize in Cameroun. *Agriculture Journal*, 2007;2(2): 192-196
21. Abbott WS. A method of computing the effectiveness of an insecticide. *Journal of Economic Entomology* 1925; 18: 65 – 66
22. Obeng-Ofori D, Reichmuth CH, Bekele J, Hassanali A. Biological activity of 1, 8 cineole, a major component of essential oil of *Ocimum kenyense* (Ayobangira) against stored product beetles. *Journal of Applied Entomology*, 1997; 121(1-5):237-243, <https://doi.org/10.1111/J-1439-0418.1997.tt01399>
23. Oboho DE, Eyo JE, Ekeh FN, Okweche S. Efficacy of *Cymbopogon citratus* Stapf leaf extract as seeds protectant against *Sitophilus zeamais* Motschulsky (Coleoptera: Curculionidae) on stored maize grains. *Journal of Biological Control*, 2016; 30(4): 220-225, <https://doi.org/10.18311/jbc/2016/15540>

24. Ishayaa I, Swirski E. Trehalase, invertase and amylase activities in the black scale Saissetiaoleae and their relation to host adaptability. *Journal of Insect Physiology*, 1976; 22: 1025- 1029, [https://doi.org/10.1016/0022-1910\(76\)90087-1](https://doi.org/10.1016/0022-1910(76)90087-1)
25. Adel MM, El-Hawary FM, Abdel-Aziz NF, Sammour EA. Some physiological, biochemical and histopathological effects of *Artemisia monosperma* against the cotton leafworm, *Spodoptera littoralis*. *Archives of Phytopathology and Plant Protection*, 2010;43(11): 1098-1110, <https://doi.org/10.1080/03235400802285562>
26. Reitman S, Frankel S. A calorimetric method for the determination of serum glutamic oxaloacetic and glutamic pyruvic transaminases. *American Journal of Clinical Pathology* 1957; 28: 56- 63
27. Powell MEA, Smith MJH. The determination of serum acid and alkaline phosphatases activity with 4-amino antipyrine. *Journal of Clinical Pathology*, 1954;7: 245-248, <https://dx.doi.org/10.1136%2Fjcp.7.3.245>
28. Basse ME, Etuk UI, Ekpo JU. Morphological diversity in the macrophyte genus *Lasianthera* (Icacinaceae) and the taxonomic implications. *Living System for Sustainable Development*, 2014;2: 1-5.
29. Folawewo AD, Madu AN, Agbaje-Daniels FV, Faboyede AO, Coker AR. Phytochemical screening and antibacterial activities of the root bark extracts of *Hippocratea africana* (Willd.) Loes. ex Engl. *European Journal of Medicinal Plants*, 2017;19(1): 1-8, <http://doi.org/10.9734/EJMP/2017/32765>
30. Oboho DE, Oyebadejo S, Edagha I, Ubulom PME, Ita BN, Nelson AU, Akpan AU, Eyo JE. Gas Chromatography-Mass Spectroscopy and Histopathological Effects of Methanol Leaf extract of *Uvaria chamae* on the Midgut of *Sitophilus zeamais*. *International Journal of Agriculture and Biology* 2021, 26(6): 695- 701
31. Ekanem NG, Mbagwu HOC, Harry GI. Phytochemical screening and hypoglycaemic activity of *Lasianthera africana* Beauv. (Aquifoliales: Stemonuraceae) leaf extract on diabetic rats. *Brazilian Journal of Biological Sciences*, 2016; 3(6): 293-298, <http://dx.doi.org/10.21472/bjbs.030606>
32. Ebana RU, Asamudo NU, Etok CA, Edet UO, Onyebuisi CS. Phytochemical Screening, Nutrient Analysis and Antimicrobial Activity of the leaves of *Lasianthera africana* and *Dennettia tripetala* on Clinical Isolates. *Journal of Advances in Biology and Biotechnology*, 2016;8(4): 1-9, <http://dx.doi.org/10.973-/JABB/2016/28291>
33. Oboho DE, Akwaowo UN, Edeke A, Okwor JI, Imakwu, CA, Akpan AU, Oyebadejo S, Eyo JE. Phytochemical screening, GC-MS and Histological Effects of Methanolic leaf extract of *Hippocratea africana* (Willd) on the midgut of *Sitophilus zeamais* (Motsch.). *London Journal of Science Research*, 2022 22(4):11-21
34. Rajeswari K, Ravi Kumar A, Subbu Rathinam KM. Phytochemical and antidiarrhoeal activity of *Hippocratea africana* roots. *Indian Journal of Research in Pharmacy and Biotechnology*, 2014; 2(4): 1357- 1359
35. Tabassum R, Naqvi SNH, Jahan M, Nurulain SM, Khan MF, Azmi, MA. Determination of the toxicities of fenpropathrin (Pyrethroid) and neem formulation (RB- a+ PBO+ Tx-100) against *Alphitobius diaperinus* adults and their effects on transaminases. *Turkish Journal of Zoology*, 1998; 22(4): 319-322.

36. Hassan HA. Biological and Biochemical Studies on the Effects of Some Botanical Extracts on Cotton Leafworm *Spodoptera littoralis* (Boisd.) (Lepidoptera: Noctuidae). M. Sc. Thesis, Faculty of Science, Ain Shams University, 2002 Egypt.
37. Sharma RK, Singh K (1977). Studies on glutamic-oxalacetic (GOT) and glutamic-pyruvic (GPT) transaminases of swine kidney worm *Stephanurus dentatus* (Diesing, 1839). Zeitschrift für Parasitenkunde, 1977;54 (3): 251-256.
38. Azmi MA, Naqvi SNH, Khan MF, Akhtar K, Khan FY. Comparative toxicological studies of RB-a (Neem extract) and Coopex (Permethrin + Bioallethrin) against *Sitophilus oryzae* with reference to their effects on oxygen consumption and GOT, GPT activity. Turkish Journal of Zoology, 1998; 22(4): 307-310.
39. El-Hawary FMA, Sammour EA. The bioactivity and mechanism of action of certain wild plant extracts on *Aphis craccivora*. Bulletin of the National Research Centre, Egypt, 2006;31: 547-556.
40. El-Gindi AM. Effect of pyriproxyfen on phosphatases and transaminases enzymes of last instar larvae of *Parasarcophaga argyrostoma* (Robieau-Desvoidy) (Diptera-Sarcophagidae). Journal of Egyptian and German Society of Zoology, 2000; 33(E): 327-338.
41. Mostafa SA. Biochemical Effect of Some Chemical Compounds on *Spodoptera littoralis* (Boisd). PhD Thesis, Faculty of Agriculture, Al-Azhar University, 1993; Egypt, <https://doi.org/10.1093/ajcp/28.1.56>
42. Younes MW, Othman SE, Elkersh MA, Youssef NS, Omar GA. Effect of seven plant oils on some biochemical parameters in Khapra beetle *Trogoderma granarium* Everts (Coleoptera: Dermestidae). Egyptian Journal of Experimental Biology (Zoology), 2011; 7(1): 53-61.
43. Bakr FR, El-Bermawy S, Emara S, Abolyazid I, Abdel-Wahab H. Biochemical studies on *Spodoptera littoralis* developmental stages after larval treatment with different botanical extracts. The Proceedings of the 2nd International Conference of Plant Protection Research Institute, Cairo, Egypt, 2002;1: 886- 897
44. Hussein NM, Gadallah AI, Tawfik SM, Hewady MA. The changes in the enzymatic activities in the larvae of bullworm induced by vertimec and /or neem azal in their artificial diet. Paper Presented at the First Conference of the Central Agricultural Pesticide Laboratory, September 3-5, 2002, Cairo Egypt.
45. Sammour EA, El-Hawary FM, Abdel-Aziz NF. Comparative study on the efficacy of neemix and basil oil formulations on the cowpea aphid *Aphis craccivora* Koch. Archives of Phytopathology and Plant Protection, 2011; 44(7): 655-670, <https://doi.org/10.1080/03235400903266495>
46. Ferraro RB, Sousa JL, Cunha RD, Meyer Fernandes JR. Characterization of an ecto-phosphatase activity in malpighian tubules of hematophagous bug *Rhodnius prolixus*. Archives of Insect Biochemistry and Physiology, 2004;57(1): 40-49.
47. Dua VK, Pandey AC, Dash AP. Adulticidal activity of essential oil of *Lantana camara* leaves against mosquitoes. Indian Journal of Medical Research, 2010 131: 434-439.
48. Werdin Gonzalez JO, Gutiérrez MM, Murray AP, Ferrero AA. Biological activity of essential oils from *Aloysia polystachya* and *Aloysia citriodora* (Verbenaceae) against the soybean

pest *Nezara viridula* (Hemiptera: Pentatomidae). Natural Product Communications, 2010; 5(2): 301-306.

UNDER PEER REVIEW