
On the Properties of Functions Defining Distinguished Varieties

Abstract

An inner toral polynomial is a polynomial in $\mathbb{C}[z, w]$ such that its zero set is contained in $\mathbb{D}^2 \cup \mathbb{T}^2 \cup \mathbb{E}^2$, where \mathbb{D} is the open unit disc, \mathbb{T} is the unit circle and \mathbb{E} is the exterior of the closed unit disc in \mathbb{C} . Given such a polynomial p , its zero set that lies inside \mathbb{D}^2 , i.e. $V = Z(p) \cap \mathbb{D}^2$ is called a distinguished variety, and p is called a polynomial defining the distinguished variety V . An inner toral polynomial always gives a distinguished variety and vice versa. Finite Blaschke products generate inner toral polynomials such a way that, given a finite Blaschke product $B(z)$, the numerator of $w^m - B(z)$ is an inner toral polynomial. In this paper, we investigate the conditions that define the sum and the composition of inner toral polynomials generated by finite Blaschke products on the inner toral.

Keywords: Automorphisms, Blaschke products, Distinguished varieties, Inner toral polynomials

2010 Mathematics Subject Classification: 30C15; 30J10; 32B99

1 Introduction

An algebraic variety is a subset of \mathbb{C}^n that can be written as a set of common zeroes to a collection of polynomials in $\mathbb{C}[z_1, z_2, z_3, \dots, z_n]$ ([7]). A distinguished variety is a special variety generated from a single two complex variables polynomial such that it lies in \mathbb{D}^2 and extends to \mathbb{C}^2 through the distinguished boundary \mathbb{T}^2 , where \mathbb{D} is the unit disc and \mathbb{T} is the unit circle in the complex plane. Jim Agler and John E. McCarthy recently initiated the work on distinguished varieties ([5]), though the history goes back to Rudin's time. Study on distinguished varieties and function spaces on distinguished varieties as well as operators on such spaces were later became interested topics among researchers ([1], [3], [10], [4], [2]). In 2010, Greg Knese ([4]) gave a new proof of a representation for distinguished variety by using sum of squares formula for two variables polynomials with no zeros on the bidisk. He also discussed symmetricity properties of polynomials defining distinguished varieties: Polynomials defining distinguished varieties are symmetric with respect to the distinguished boundary \mathbb{T}^2 . In this work we focus on exploring algebraic properties of polynomials defining distinguished varieties. It is straightforward the set of all polynomials defining distinguished

varieties is closed under scalar multiplication and product. However, as it turned out, the set of all polynomials defining distinguished varieties is not closed under addition and under composition. In ([4]), it was presented that given a finite Blaschke product $B(z)$, the zero set of $w^m - B(z)$ is a distinguished variety. In other words, the numerator of $w^m - B(z)$ is a polynomial defining distinguished variety. In this paper, we worked on polynomials generated by finite Blaschke products above manner to investigate the summation and the composition of polynomials defining distinguished varieties. In section 3, we presents results we obtained for the summation and the composition on such polynomials respectively.

2 Preliminaries

2.1 Polynomials Defining Distinguished Varieties

In this section we present the necessary preliminary materials so that the reader can smoothly move to the results section. Through out this article, we consider \mathbb{D} to be the open unit disc, \mathbb{T} to be the unit circle and \mathbb{E} to be the exterior of the closed unit disc in the complex plane. Among the couple of versions of the definition of distinguished varieties available in the literature ([4],[5],[10]), we will stay with the following version:

Definition 2.1. [5] A non-empty set V in \mathbb{D}^2 is called a distinguished variety if there is a polynomial $p \in \mathbb{C}[z, w]$ such that

$$V = \{(z, w) \in \mathbb{D}^2 : p(z, w) = 0\}$$

and the extension of V to \mathbb{C}^2 is a subset of $\mathbb{D}^2 \cup \mathbb{T}^2 \cup \mathbb{E}^2$.

In this case, we say p defines the distinguished variety V and we call p an *inner toral polynomial*. For example, $z^n - w^m$ for $n, m \in \mathbb{N}$ is an inner toral polynomial.

A polynomial in two complex variables z and w , $p(z, w) \in \mathbb{C}[z, w]$, is said to have *bidegree* (n, m) if p has degree n in z and degree m in w ([1]). As it proved in ([4]), polynomials defining distinguished varieties are symmetric with respect to the distinguished boundary \mathbb{T}^2 in the following way:

Definition 2.2. [4] Let $p \in \mathbb{C}[z, w]$ be a polynomial of bidegree (n, m) . The reflection polynomial \tilde{p} of p at the degree (n, m) given by,

$$\tilde{p}(z, w) = z^n w^m \overline{p\left(\frac{1}{z}, \frac{1}{w}\right)}$$

Definition 2.3. [4] A polynomial $p \in \mathbb{C}[z, w]$ is said to be essentially \mathbb{T}^2 -symmetric if there exists a unimodular constant c such that

$$\tilde{p}(z, w) = c p(z, w).$$

We call this coefficient c the *reflective coefficient* of p .

Proposition 2.1. [4] A Polynomial defining a distinguished variety is essentially \mathbb{T}^2 -symmetric.

For an example, the polynomial $p_1(z, w) = 4w^2 + izw^2 - 4z + i$ is an inner toral polynomial and defines a distinguished variety. Observe that $\tilde{p}_1(z, w) = 4zw^2\left(\frac{4}{w^2} + \frac{i}{zw^2} - \frac{4}{z} + i\right) = -1(4w^2 + izw^2 - 4z + i) = -p_1(z, w)$. Therefore p_1 is essentially \mathbb{T}^2 -symmetric with the reflective coefficient -1 .

The rest of this chapter is devoted to discuss a special type of polynomials defining distinguished varieties: polynomials generated by Blaschke products.

Definition 2.4. [9] The rational function $B(z) = \zeta \prod_{i=1}^k \left(\frac{z - a_i}{1 - \bar{a}_i z} \right)^{m_i}$ is called a finite Blaschke product of degree n , where $\zeta \in \mathbb{T}$, $a_i \in \mathbb{D}$, $n = \sum_{i=1}^k m_i$ and m_i is the multiplicity of the zero a_i for $i = 1, 2, 3, \dots, k$.

Proposition 2.2. [4] Let $B(z)$ be a finite Blaschke product of degree n . The set $V = \{(z, w) \in \mathbb{D}^2 \mid w^m - B(z) = 0\}$ is a distinguished variety.

In other words, the numerator of $w^m - B(z)$ is a polynomial defining a distinguished variety (or an inner toral polynomial). Proposition 2.2 was given as just a statement in page 10 in ([4]). We present a quick proof for the claim for the sake of the completeness.

Proof. Note that

$$|B(z)| \text{ is } \begin{cases} = 1 & \text{iff } |z| = 1 \\ < 1 & \text{iff } |z| < 1 \\ > 1 & \text{iff } |z| > 1 \end{cases}$$

If (z_0, w_0) is a zero of $w^m - B(z)$, then $|w_0|^m = |B(z_0)|$. Observe that $|w_0| = 1$ iff $|B(z_0)| = 1$ iff $|z_0| = 1$. Likewise, $|w_0| < 1$ iff $|z_0| < 1$ and $|w_0| > 1$ iff $|z_0| > 1$. Therefore, $(z_0, w_0) \in \mathbb{D}^2 \cup \mathbb{T}^2 \cup \mathbb{E}^2$. Therefore $\{(z, w) \in \mathbb{C}^2 : w^m - B(z) = 0\}$ is a distinguished variety. Therefore, the numerator of $w^m - B(z)$ is inner toral. \square

By Proposition 2.2, The numerator of $w^m - B(z)$, is an inner toral polynomial with the bidegree (n, m) . For the rest of the article, we focus on such inner toral polynomials.

3 Results

3.1 On the Summation of Inner Toral Polynomials Generated by Finite Blaschke Products

It is obvious that the scalar multiplication and products of inner toral polynomials are also inner toral. However, sum of two inner toral polynomials is not always inner toral. For example, $z - w$ and $z^2 - w$ are two inner toral polynomials, however the sum, $z^2 + z - 2w$, is not inner toral, because $(-1, 0)$ is a zero of $z^2 + z - 2w$, but it is not in $\mathbb{D}^2 \cup \mathbb{T}^2 \cup \mathbb{E}^2$. **In this chapter** we investigate the **factors that makes** a sum of two inner toral polynomials generated by Blaschke products is inner toral, and we present following results.

Proposition 3.1. *The reflective coefficient of an inner toral polynomial generated by the finite Blaschke product $B(z) = \zeta \prod_{i=1}^k \left(\frac{z - m_i}{1 - \bar{a}_i z} \right)^{m_i}$ is $-\bar{\zeta}$.*

Proof. Let $p(z, w)$ be the numerator $w^m - B(z)$ and let $n = m_1 + m_2 + \dots + m_k$. Note that $p(z, w) = \prod_{i=1}^k (1 - \bar{a}_i z)^{m_i} w^m - \zeta \prod_{i=1}^k (z - a_i)^{m_i}$ and by Proposition 2.2, p is an inner toral polynomial of bidegree (n, m) . Therefore, by Proposition 2.1, $p(z, w)$ is essentially \mathbb{T}^2 -symmetric. To find the

reflective coefficient, consider $\tilde{p}(z, w)$.

$$\begin{aligned} \tilde{p}(z, w) &= w^m z^n \overline{\left\{ \prod_{i=1}^k \left(1 - \frac{\bar{a}_i}{z}\right)^{m_i} \frac{1}{w^m} - \zeta \prod_{i=1}^k \left(\frac{1}{z} - a_i\right)^{m_i} \right\}} \\ &= w^m z^n \left\{ \prod_{i=1}^k \left(1 - \frac{a_i}{z}\right)^{m_i} \frac{1}{w^m} - \bar{\zeta} \prod_{i=1}^k \left(\frac{1}{z} - \bar{a}_i\right)^{m_i} \right\} \\ &= \prod_{i=1}^k \left(\frac{z - a_i}{z}\right)^{m_i} z^n - \bar{\zeta} w^m z^n \prod_{i=1}^k \left(\frac{1 - \bar{a}_i z}{z}\right)^{m_i} \\ &= \bar{\zeta} \left\{ \prod_{i=1}^k (1 - \bar{a}_i z)^{m_i} w^m - \zeta \prod_{i=1}^k (z - a_i)^{m_i} \right\} \\ &= -\bar{\zeta} p(z, w). \end{aligned}$$

Therefore the reflective coefficient of p is $-\bar{\zeta}$. □

Proposition 3.2. Let $B_t(z)$ is the Blaschke product of degree n given by $B_t(z) = \zeta_t \prod_{i=1}^{k_t} \left(\frac{z - a_{ti}}{1 - \bar{a}_{ti}z}\right)^{m_{ti}}$

where $t = 1, 2, i = 1, 2, \dots, k_t$ and $n = \sum_{i=1}^{n_t} m_{ti}$. Let $p_t(z, w)$ be the inner toral polynomial generated by $B_t(z)$. If $p = p_1 + p_2$ is inner toral, then $\zeta_1 = \zeta_2$.

Proof. Note that $p_t(z, w) = \prod_{i=1}^{k_t} (1 - \bar{a}_{ti}z)^{m_{ti}} w^m - \zeta_t \prod_{i=1}^{n_t} (z - a_{ti})^{m_{ti}}$ for $t = 1, 2$. By Proposition 2.2, both p_1 and p_2 are inner toral polynomials with bidegree (n, m) . Note that $p(z, w) = p_1(z, w) + p_2(z, w)$ has bidegree (n, m) if $\zeta_1 \neq -\zeta_2$. Now, assuming $\zeta_1 \neq -\zeta_2$, we have

$$\tilde{p} = \widetilde{p_1 + p_2} = w^m z^n \overline{(p_1 + p_1) \left(\frac{1}{z}, \frac{1}{w}\right)} = w^m z^n \overline{p_1 \left(\frac{1}{z}, \frac{1}{w}\right)} + w^m z^n \overline{p_2 \left(\frac{1}{z}, \frac{1}{w}\right)} = \tilde{p}_1 + \tilde{p}_2.$$

By Proposition 3.1, p_1 and p_2 are essentially \mathbb{T}^2 -symmetric with reflective coefficients $-\bar{\zeta}_1$ and $-\bar{\zeta}_2$ respectively. Therefore $\tilde{p} = -\bar{\zeta}_1 p_1(z, w) - \bar{\zeta}_2 p_2(z, w)$. If p is inner toral, then by Proposition 2.1, p is essentially \mathbb{T}^2 -symmetric with reflective coefficient $-\bar{\zeta}$ (say). That is $\tilde{p} = -\bar{\zeta} p$. Therefore, $\zeta_1 = \zeta_2 = \zeta$. □

A sort of a converse to the above result can be given as follows:

Proposition 3.3. Let $B(z) = \prod_{i=1}^{k-1} \left(\frac{z - a_i}{1 - \bar{a}_i z}\right)^{m_i}$ where $m_i \in \mathbb{N}$ and $i = 1, 2, \dots, (k-1)$. For $t = 1, 2$,

let $B_t(z) = \zeta_t B(z) \left(\frac{z - a_{k_t}}{1 - \bar{a}_{k_t} z}\right)$ and $p_t(z, w)$ be the inner toral polynomial generated by $B_t(z)$. If $\zeta_1 = \zeta_2$, then $p = p_1 + p_2$ is an inner toral polynomial.

Proof. Note that for $t = 1, 2$, $p_t(z, w) = \left[\prod_{i=1}^{k-1} (1 - \bar{a}_i z)^{m_i} \right] (1 - \bar{a}_{k_t} z) w^m - \zeta_t \left[\prod_{i=1}^{k-1} (z - a_i)^{m_i} \right] (z - a_{k_t})$.

Suppose that $\zeta_1 = \zeta_2 = \zeta$ (say). A simple calculation shows that,

$$p(z, w) = \left[\prod_{i=1}^{k-1} (1 - \bar{a}_i z)^{m_i} \right] \left[2 - (\bar{a}_{k_1} + \bar{a}_{k_2})z \right] w^m - \zeta \left[\prod_{i=1}^{k-1} (z - a_i)^{m_i} \right] [2z - (a_{k_1} + a_{k_2})].$$

Let $a = \left(\frac{a_{k_1} + a_{k_2}}{2}\right)$. Since $a_{k_1}, a_{k_2} \in \mathbb{D}$, a is also in \mathbb{D} . If $(z_0, w_0) \in Z(P)$, then $w_0^m = \zeta B(z_0) \left(\frac{z_0 - a}{1 - \bar{a}z_0}\right)$. It easily follows that $(z_0, w_0) \in \mathbb{D}^2 \cup \mathbb{T}^2 \cup \mathbb{E}^2$. Therefore, p is an inner toral polynomial. □

3.2 On the Composition of Inner Toral Polynomials Generated by Finite Blaschke Products

An automorphism is simply an isomorphism from one set to itself. Every automorphism on the unit disc \mathbb{D} , is of the form $\frac{az + b}{bz + \bar{a}}$ where $a, b \in \mathbb{C}$ and $|a|^2 - |b|^2 = 1$. Alternatively, an automorphism on \mathbb{D} has the form $e^{i\theta} \frac{z - c}{1 - \bar{c}z}$ where $c \in \mathbb{D}$ and $\theta \in (0, 2\pi]$. Every automorphism on the bidisc \mathbb{D}^2 , is of the form $f = (f_1, f_2)$, where f_1 and f_2 are automorphisms on the unit disc ([8],[6]). Every automorphism, f , on the complex plane is of the form $f(z) = az + b$, where $a, b \in \mathbb{C}$ and $a \neq 0$. An automorphism on the two dimensional complex plane is of the form $f = (f_1, f_2)$ where f_k for $k = 1, 2$ are automorphism on the complex plane; that is an automorphism $f : \mathbb{C}^2 \rightarrow \mathbb{C}^2$ can be given by $f(z_1, z_2) = (f_1(z_1), f_2(z_2))$ for all $(z_1, z_2) \in \mathbb{C}^2$ ([6]).

Definition 3.1. Let $f = f(x_1, x_2, \dots, x_n)$ be a function of n variables and let $g = (g_1, g_2, \dots, g_n)$ be an n -tuple of functions. The composition $f \circ g$ is defined as $f \circ g := f(g_1, g_2, \dots, g_n)$.

Proposition 3.4. Let $B(z)$ be a finite Blaschke product of degree n and let $p(z, w)$ be the inner toral polynomial generated by $B(z)$. If f is an automorphism on the bidisc, then the composition of p with f , $p \circ f$ also defines an inner toral polynomial.

Proof. Let $B(z) = \zeta \prod_{i=1}^k \left(\frac{z - a_i}{1 - \bar{a}_i z} \right)^{m_i}$ where $\zeta \in \mathbb{T}$, $a_i \in \mathbb{D}$, $n = \sum_{i=1}^k m_i$ and m_i is the multiplicity of the zero a_i for $i = 1, 2, 3, \dots, k$. Recall that polynomial p is of the form $p(z, w) = w^m \prod_{i=1}^k (1 - \bar{a}_i z)^{m_i} - \zeta \prod_{i=1}^k (z - a_i)^{m_i}$. Let f be an automorphism on the bidisc of the form $f = (f_1, f_2) = \left(e^{i\theta_1} \frac{z - a_1}{1 - \bar{a}_1 z}, e^{i\theta_2} \frac{w - a_2}{1 - \bar{a}_2 w} \right)$ where $a_1, a_2 \in \mathbb{D}$ and $\theta_1, \theta_2 \in (0, 2\pi]$. Observe that

$$p \circ f(z, w) = \left(e^{i\theta_2} \frac{w - a_2}{1 - \bar{a}_2 w} \right)^m \prod_{i=1}^k \left\{ 1 - \bar{a}_i \left(e^{i\theta_1} \frac{z - a_1}{1 - \bar{a}_1 z} \right) \right\}^{m_i} - \zeta \prod_{i=1}^k \left\{ \left(e^{i\theta_1} \frac{z - a_1}{1 - \bar{a}_1 z} \right) - a_i \right\}^{m_i}.$$

Let (z_0, w_0) be a zero of $p \circ f$, and let $Z_0 = e^{i\theta_1} \frac{z_0 - a_1}{1 - \bar{a}_1 z_0}$ and $W_0 = e^{i\theta_2} \frac{w_0 - a_2}{1 - \bar{a}_2 w_0}$. Observe that $p \circ f(Z_0, W_0) = W_0^m \prod_{i=1}^k (1 - \bar{a}_i Z_0)^{m_i} - \zeta \prod_{i=1}^k (Z_0 - a_i)^{m_i} = 0$. That is, (Z_0, W_0) is a zero of p .

Consequently, we have $W_0^m \prod_{i=1}^k (1 - \bar{a}_i Z_0)^{m_i} = \zeta \prod_{i=1}^k (Z_0 - a_i)^{m_i}$. Taking the modulus of both sides,

$$\text{we have } |W_0|^m = \prod_{i=1}^k \left| \frac{Z_0 - a_i}{1 - \bar{a}_i Z_0} \right|^{m_i}.$$

If $W_0 \in \mathbb{D}$, then $\left| \frac{Z_0 - a_i}{1 - \bar{a}_i Z_0} \right| < 1$ for $a_i \in \mathbb{D}$ where $i = 1, 2, 3, \dots, k$, and hence $Z_0 \in \mathbb{D}$. Likewise, if $Z_0 \in \mathbb{D}$, then $W_0 \in \mathbb{D}$. Similarly, if $Z_0 \in \mathbb{T}$, then $W_0 \in \mathbb{T}$ and vice versa, and if $Z_0 \in \mathbb{E}$, then $W_0 \in \mathbb{E}$ and vice versa. Therefore $(Z_0, W_0) \in \mathbb{D}^2 \cup \mathbb{T}^2 \cup \mathbb{E}^2$. Solving for z_0 and w_0 , we have $Z_0 = e^{-i\theta_1} \frac{z_0 - e^{i\theta_1} a_1}{1 - e^{i\theta_1} \bar{a}_1 z_0}$ and $W_0 = e^{-i\theta_2} \frac{w_0 - e^{i\theta_2} a_2}{1 - e^{i\theta_2} \bar{a}_2 w_0}$. Since $(Z_0, W_0) \in \mathbb{D}^2 \cup \mathbb{T}^2 \cup \mathbb{E}^2$, so is (z_0, w_0) . Therefore, $p \circ f$ defines an inner toral polynomial. \square

Proposition 3.5. Let $B(z) = \zeta z^n$, where $\zeta \in \mathbb{T}$ and let p be the inner toral polynomial generated by $B(z)$. Let f be an automorphism on the two dimensional complex plane \mathbb{C}^2 . The composition $p \circ f$ defines an inner toral polynomial iff f has the form $f = (az, cw)$ with $|a|^n = |c|^m$.

Proof. Suppose that f has the form $f = (az, cw)$ with $|a|^n = |c|^m$. Let (z_0, w_0) be a zero of $p \circ f$. So we have $p \circ f(z_0, w_0) = (cw_0)^m - \zeta(az_0)^n = 0$. That is $(cw_0)^m = \zeta(az_0)^n$. Taking the modulus of both sides, since $|a|^n = |c|^m$, we have $|z_0|^n = |w_0|^m$. Therefore $(z_0, w_0) \in \mathbb{D}^2 \cup \mathbb{T}^2 \cup \mathbb{E}^2$ and $p \circ f$ defines an inner toral polynomial.

Conversely, suppose that polynomial $p \circ f$ defines an inner toral polynomial. By Proposition 2.1, $p \circ f$ is essentially \mathbb{T}^2 symmetric. That is, $(cw + d)^m - \delta(az + b)^n = \gamma\{z^n(\bar{c} + \bar{d}w)^m - \bar{\delta}(\bar{a} + \bar{b}z)^n\}$ with $|\gamma| = 1$. By comparing coefficient, $|a|^n = |c|^m$ and $b = d = 0$. Therefore, f has the form $f = (az, cw)$ and $|a|^n = |c|^m$. \square

We further generalize this result in the following way. Proposition 3.6 provides the necessary and sufficient condition for a composition of an inner toral polynomial generated by a finite Blaschke product with an automorphism on \mathbb{C}^2 to be inner toral.

Proposition 3.6. Let $B(z)$ be a finite Blaschke product of degree n and p be the inner toral polynomial generated by $B(z)$. If f is an automorphism on \mathbb{C}^2 , then $p \circ f$ defines an inner toral polynomial iff $f = (az, cw)$ with $|a| = |c| = 1$.

Proof. Let $B(z) = \zeta \prod_{i=1}^k \left(\frac{z - a_i}{1 - \bar{a}_i z} \right)^{m_i}$, where $\zeta \in \mathbb{T}$, $a_i \in \mathbb{D}$, $n = \sum_{i=1}^k m_i$ and m_i is the zero of a_i

for $i = 1, 2, 3, \dots, k$. The polynomial p has the form $p(z, w) = w^m \prod_{i=1}^k (1 - \bar{a}_i z)^{m_i} - \zeta \prod_{i=1}^k (z - a_i)^{m_i}$.

Suppose that $f \in \text{Aut}(\mathbb{C}^2)$ has the form $f = (az, cw)$ with $|a| = |c| = 1$. For a zero (z_0, w_0) of $p \circ f$, we have $p \circ f(z_0, w_0) = (cw_0)^m \prod_{i=1}^k (1 - \bar{a}_i az_0)^{m_i} - \zeta \prod_{i=1}^k (az_0 - a_i)^{m_i} = 0$. That is,

$(cw_0)^m \prod_{i=1}^k (1 - \bar{a}_i az_0)^{m_i} = \zeta \prod_{i=1}^k (az_0 - a_i)^{m_i}$. Taking the modulus of both sides, since $|a| = |c| = 1$,

we have $|w_0|^m = \prod_{i=1}^k \left| \frac{z_0 - \frac{a_i}{a}}{1 - \frac{\bar{a}_i}{a} z_0} \right|^{m_i}$. Therefore $(z_0, w_0) \in \mathbb{D}^2 \cup \mathbb{T}^2 \cup \mathbb{E}^2$ and $p \circ f$ defines an inner toral polynomial.

Conversely, let f be an automorphism on \mathbb{C}^2 and suppose that $f \neq (az, cw)$, or $|a| \neq 1$ or $|c| \neq 1$. If $f \neq (az, cw)$, then f has the form $f = (az + b, cw)$ or $f = (az, cw + d)$ or $f = (az + b, cw + d)$ where $b, d \neq 0$. Now we consider the two cases separately.

Case 1: $f \neq (az, cw)$

Suppose $f = (az + b, cw)$ for $b \neq 0$. Let $B_1(z) = \frac{z - (1/2)}{1 - (1/2)z}$ and $p_1(z, w) = w\{1 - (1/2)z\} - \{z - (1/2)\}$ be the inner toral polynomial generated by $B_1(z)$. Also let $f_1 = (z + 1, w)$. Note that, $p_1 \circ f_1(z, w) = w\{1 - (1/2)(z + 1)\} - \{(z + 1) - (1/2)\}$ and $(1/2, 4)$ is a zero of $p \circ f$. However, $(1/2, 4) \notin \mathbb{D}^2 \cup \mathbb{T}^2 \cup \mathbb{E}^2$. Therefore $p_1 \circ f_1$ does not define an inner toral polynomial. Similarly, if $f = (az, cw + d)$ with $d \neq 0$, we can make choices for f and p such that $p \circ f$ does not define an inner toral polynomial. Now suppose $f = (az + b, cw + d)$ for $b, d \neq 0$. Let $f_2 = (z + 2, w + 1)$, $B_2(z) = \frac{z - (1/2)}{1 - (1/2)z}$, and $p_2(z, w) = w\{1 - (1/2)z\} - \{z - (1/2)\}$ be the inner toral polynomial generated by $B_2(z)$. Observe that $(0, -6)$ is a zero of $p_2 \circ f_2$. However, $(0, -6) \notin \mathbb{D}^2 \cup \mathbb{T}^2 \cup \mathbb{E}^2$. Therefore $p_2 \circ f_2$ does not define an inner toral polynomial.

Case 2: $|a| \neq 1$ or $|c| \neq 1$

Without loss of generality, assume $|a| \neq 1$. Let $f_3 = (2z, w)$, $B_3(z) = \frac{z - (1/2)}{1 - (1/2)z}$, and $p_3(z, w) = w\{1 - (1/2)z\} - \{z - (1/2)\}$ be the inner toral polynomial generated by $B_3(z)$. Observe that $(1/2, 1)$ is a zero of $p_3 \circ f_3$ that does not lie in $\mathbb{D}^2 \cup \mathbb{T}^2 \cup \mathbb{E}^2$, and hence $p_3 \circ f_3$ does not define an inner toral polynomial.

Therefore, if $f \neq (az, cw)$, or $|a| \neq 1$ or $|c| \neq 1$, then $p \circ f$ does not define an inner toral polynomial. By taking the contrapositive, if $p \circ f$ defines an inner toral polynomial, then $f = (az, cw)$ with $|a| = |c| = 1$. \square

4 CONCLUSIONS

In this study we have proved that the reflective coefficient of a polynomial that defining a distinguished variety, generated by a finite Blaschke product with Blaschke coefficient is $-\bar{\zeta}$ and if the sum of two polynomials that defining distinguished variety generated by Blaschke products gives distinguished variety, then the Blaschke coefficients are the same. In addition, we proved that the sum of two polynomials that defining distinguished variety generated by Blaschke products which are differ only by one factor with multiplicity one, is defining distinguished variety iff their Blaschke coefficients are the same. Moreover, we prove that when we taking the composition of a polynomial that defines a distinguished variety and that generated by Blaschke product, with an automorphism on the bidisc we get back a polynomial defining a distinguished variety. Further, we have generalized this result by considering composition such polynomials with automorphisms on the complex plane and obtained similar results.

References

- [1] J. Agler, G. Knese, and J. E. McCarthy. Algebraic pairs of isometries. *Journal of Operator Theory*, pages 215–236, 2012.
- [2] T. Bhattacharyya, P. Kumar, and H. Sau. Distinguished varieties through the berger-coburn-lebow theorem. *Analysis and PDE*, 15(2):477–506, 2022.
- [3] M. T. Jury, G. Knese, and S. McCullough. Nevanlinna–pick interpolation on distinguished varieties in the bidisk. *Journal of Functional Analysis*, 262(9):3812–3838, 2012.
- [4] G. Knese. Polynomials defining distinguished varieties. *Transactions of the American Mathematical Society*, 362(11):5635–5655, 2010.

-
- [5] J. E. McCarthy and J. Agler. Distinguished varieties. *Acta Mathematica*, 194(2):133–154, 2005.
 - [6] A. Persson. The automorphism groups on the complex plane, 2017.
 - [7] A. Sheffer. Basic real algebraic geometry. 2015.
 - [8] J. A. Strong. *Boundary Characterization of Iterated Automorphism Orbits on Bounded Domains*. PhD thesis, UC Riverside, 2016.
 - [9] S. Uçar and N. Y. Özgür. Finite blaschke products and decomposition. *arXiv preprint arXiv:1712.07850*, 2017.
 - [10] U. D. Wijesooriya. Algebraic pairs of pure commuting isometries with finite multiplicity. *Journal of Operator Theory*, 79(2):507–527, 2018.