

Original Research Article

Heterosis and Inbreeding Depression for Quantitative Characters in Bread Wheat (*Triticum aestivum* L. em. Thell.)

Abstract

Aim-: Determine the magnitude of heterosis, heterobeltiosis and inbreeding depression in bread wheat and identified superior cross combinations for commercial production of hybrid wheat.

Study design: The experiment was laid out in Randomized block design with three replications.

Place and Duration of Study: The experimental material was evaluated during Rabi 2021-22 at RARI, Durgapura.

Methodology: The ten genotypes of wheat viz. Raj 3077, Raj 4079, Raj 3765, PBW 343, PBW 590, DBW 17, HD 2967, HD 3059, WH 1021 and UP 2425 were crossed in all possible combinations excluding reciprocals during Rabi 2018-19 to generate F₁'s. In *kharif* 2019, half of F₁'s seed was raised at IARI Regional Station Wellington (Tamil Nadu) to get F₂'s seed for experimentation. Final experimental material comprising 10 parents along with 45 F₁'s and 45 F₂'s were evaluated during Rabi 2021-22 at RARI, Durgapura.

Results: The cross combinations Raj 4079 x Raj 3765, Raj 3765 x PBW 343, Raj 3765 x WH 1021, Raj 4079 x HD 2967, Raj 3765 x HD 2967 and Raj 4079 x PBW 343 exhibited significant and desired heterobeltiosis over better parent for grain yield per plant and also showed significant and desirable heterobeltiosis for other yield attributing traits. For grain yield Raj 3077 x HD 3059, PBW 590 x HD 2967 and HD 2967 x WH 1021 showed significant and negative inbreeding depression.

Conclusion: Nature and magnitude of heterosis helps in identifying superior cross combinations to obtain better transgressive segregants. Six superior hybrids may be exploited commercially for getting benefits of heterosis for grain yield and its component traits in wheat.

Introduction

Bread wheat (*Triticum aestivum* L. em. Thell) is staple food for all over the world which belongs to Poaceae (Gramineae) family. It is annual self-pollinated crop with allohexaploid chromosome number $2n = 6x = 42$. It is popularly known as 'Stuff of life or King of the cereals' because of its

Comment [Rk1]: Line No. 15, In place of were, use was

Formatted: Underline

30 high cultivation area and production (Sharma *et al.*, 2019) and the prominent position it holds in
31 the international food grain trade. Wheat grain contains starch (60-68%), protein (6-21%), fat
32 (1.5-2.0%), cellulose (2.0-2.5%), minerals (1.8%) and vitamins. The uniqueness of wheat in
33 contrast to other cereals is that wheat contains gluten protein which enables leavened dough to
34 rise by forming minute gas cells and this property enables bakers to produce light breads. In
35 India, it is grown in about 30.54 million hectares with the production of about 106.41 million
36 tonnes with an average productivity of 3484 kg/ha [1].

37 The exploitation of heterosis is a milestone innovation in modern agriculture and it is considered
38 to be one of the pillars of global food security. In a self-pollinated crop like wheat, the utilization
39 of heterosis depends mainly upon the direction and magnitude of heterosis. The study of
40 heterosis and inbreeding depression has a direct bearing on the breeding methodology to be
41 employed for varietal improvement. In case of self-fertilizing crops where identification of
42 superior cross combinations is the breeding objective heterobeltiosis is more useful than
43 heterosis from the breeder's point of view. However, heterosis in wheat crop will not be
44 worthwhile unless, it is practicable for development of hybrid varieties. F₁ hybrid carrying
45 heterotic effects, which are featured in all crop species, the yield gains are limited to the F₁
46 generation. The F₂ and succeeding generations obtained through selfing are discarded since
47 reduced yields and developmental characters (Wang *et al.*, [2]). Heterosis is considered as the
48 superiority of the hybrids in comparisons to either of its parents. It is the allelic or non-allelic
49 interaction of genes under the influence of specific environment. Heterosis has been estimated in
50 a range of cultivated crops and has been the purpose of considerable importance to study as mean
51 of increasing productivity of crop plant. The present study has been carried out to estimate the
52 heterosis (%) over mid parent (MP), and better parent (BP) and inbreeding depression (ID) for
53 quantitative and qualitative traits in a 10x10 diallel set in bread wheat to identify parental lines
54 that could be used for commercial production of hybrid wheat as well as isolation of pure lines
55 among the progenies of heterotic F₁ for further amelioration of grain yield in bread wheat.

56 **Materials and Methods**

57 The experimental material consisted of ten genetically diverse bread wheat varieties, viz. Raj
58 3077, Raj 4079, Raj 3765, PBW 343, PBW 590, DBW 17, HD 2967, HD 3059, WH 1021 and

Comment [Rk2]: Line no. 47, use uniform pattern for citation.

59 UP 2425. These genotypes were crossed in all possible combinations excluding reciprocals
60 during Rabi 2018-19 to generate F₁s. In *khariif* 2019, half of F₁'s seed was raised at IARI
61 Regional Station Wellington (Tamil Nadu) to get F₂'s seed for experimentation. Final
62 experimental material comprising 10 parents along with 45 F₁s and 45 F₂s were evaluated during
63 Rabi 2021-22 at RARI, Durgapura. The genotypes under study were planted in a Randomized
64 Block Design with three replications per entry and one row (3m) per replication. Plant to plant
65 spacing of 10 cm and row-to-row spacing of 30 cm were maintained. Standard agronomic
66 practices were used for raising and maintenance of the plants.

67 Data were recorded on plot basis for days to heading and days to maturity, while rest of the
68 characters *viz.* plant height (cm), productive tillers per plant, flag leaf area (cm²), spike length
69 (cm), number of grains per spike, 1000-grain weight (g), biomass per plant (g), grain yield per
70 spike (g), grain yield per plant (g) and harvest index (%) were recorded on 10 randomly selected
71 plants for parents, F₁'s and while 40 plants for F₂'s. Analysis of variance, heterosis,
72 heterobeltiosis and inbreeding depression were calculated as per standard procedures given by
73 Fonseca and Patterson [3] and Panse and Sukhatme [4].

74 **Results and Discussion**

75 The analysis of variance (Table 1) depicted significant differences among the genotypes,
76 parents, and generations for all studied characters, established the circumstances that characters
77 displayed the presence of ample genetic diversity among the genotypes, parents and generations.
78 Likewise, F₁ and F₂ generations also exhibited significant differences for all studied characters.
79 The mean squares due to F₁ *vs* F₂ were found significant for all studied characters.
80 Correspondingly, the differences among the parents *vs* generations were significant for all
81 studied characters revealed the presence of heterosis. The results are in agreement with those of
82 others obtained for different characters Zare-Kohan and Heidari [5], Singh *et al.* [6] and
83 Nageshwar *et al.* [7].

84 The magnitude of heterosis provides information on the extent of genetic diversity in
85 parents of a cross and helps in choosing the parents for superior F₁'s, so as to exploit hybrid
86 vigour. The extent of heterosis in a crop depends on its exploitation uses and achievability of

Comment [Rk3]: Citation pattern not uniform

87 hybrid seed production. The appropriate technique of hybrid seed production at commercial scale
88 is not yet available in wheat due to its self-pollinating nature. Therefore, currently in this crop
89 heterosis *per se* may not be of any economic importance. However, it indicates genetic potential
90 of parental combination and if the heterosis is due to epistatic gene effects, particularly of
91 additive x additive type or due to repulsion phase linked loci, exhibiting partial or complete
92 dominance, it is possible to fix the alleles at interacting state to preserve the heterotic effects in
93 the pure lines (Arunachalam *et al.* [8]). Means transgressive segregants are possible. The
94 allopolyploid nature of wheat will also favour preservation of such hybrid vigour for a
95 considerable number of generations. Nevertheless, understanding of degree and extent of
96 heterosis is imperative for determining the direction of imminent breeding programme and the
97 selection of promising crosses to achieve superior segregants in advance generations for future
98 improvement of grain yield in bread wheat.

99 Heterosis over better parent or heterobeltiosis is more imperative and useful in
100 determining the feasibility of commercial exploitation of heterosis and also specifying the
101 parental combination capable of producing the maximum level of transgressive segregants.
102 Hence, the heterosis measured in terms of superiority over the better parent is more treasured,
103 which decides whether an experimental hybrid is worth exploiting or not.

104 The magnitude of heterosis and number of cross combinations showing heterosis over
105 better parent and inbreeding depression for grain yield per plant and its related characters are
106 present in Table 2. Grain yield per plant in wheat is the character of economic importance for
107 which 28 hybrids over mid parent and 14 hybrids over better parent exhibited significant and
108 positive heterosis while 5 crosses exhibited significant inbreeding depression. Several hybrids
109 exhibited significant and desirable direction of heterobeltiosis and inbreeding depression for
110 various characters such as days to 50% heading (15 hybrids); days to maturity (15 and 1
111 hybrids); plant height (20 and 1 hybrids); number of productive tillers per plant (20 hybrids); flag
112 leaf area (18 and 5 hybrids); spike length (15 hybrids); number of grains per spike (13 and 1
113 hybrids); 1000 grain weight (15 and 5 hybrids); biomass per plant (17 and 2 hybrids); grain yield
114 per spike (14 and 5 hybrids) and harvest index (9 and 20 hybrids), respectively. No inbreeding
115 depression was observed in any cross for the days to 50% heading, productive tillers per plant,
116 spike length and grain yield per spike which can be used for maintain the specific gene pool for

117 further utilization of improvement of wheat. The hybrid, PBW 343 x WH 1021 for days to 50%
118 heading, days to maturity, flag leaf area and biomass per plant, HD 2967 x WH 1021 for days to
119 maturity, Raj 3765 x HD 2967 for flag leaf area, Raj 3765 x WH 1021 for spike length, Raj 3765
120 x UP 2425 for number of grains per spike, PBW 343 x HD 2967 for 1000-grain weight, WH
121 1021 x UP 2425 for biomass per plant, Raj 4079 x Raj 3765 for grain yield per plant and PBW
122 590 x HD 2967 for harvest index showed significant and maximum heterosis over better parent.
123 Similar finding was reported by Gaur *et al.* [9].

124 The utility of hybrid breeding technique lies in identification of the most heterotic and
125 useful combinations in order to make commercial cultivation of hybrid beneficial. The Table 3
126 reveals a significant relation between heterobeltiosis for grain yield per plant and its associated
127 traits *i.e.* crosses which have desirable heterobeltiosis for grain yield, also exhibited desirable
128 heterobeltiosis for most of the other yield attributes. The high yielding hybrids were in general
129 the most heterotic crosses which indicated close association between hybrid mean performance
130 and manifestation of heterosis. The relative ranking of hybrids based on the mean performance.
131 This suggested that selection of hybrids should be based on *per se* performance. Similar findings
132 have been reported by Kumar *et al.* [10].

133 The cross combinations Raj 4079 x Raj 3765, Raj 3765 x PBW 343, Raj 3765 x WH 1021, Raj
134 4079 x HD 2967, Raj 3765 x HD 2967 and Raj 4079 x PBW 343 exhibited significant and
135 desired heterobeltiosis over better parent for grain yield per plant and also showed significant
136 and desirable heterobeltiosis for Days to 50 % heading, days to maturity, plant height, productive
137 tillers per plant, flag leaf area, Spike length, number of grains per spike, 1000-grain weight,
138 biomass per plant, grain yield per spike and harvest index. These three superior hybrids may be
139 exploited commercially for getting benefits of heterosis for grain yield and its component traits
140 in wheat. The result of the present investigation clearly showed that considerable heterosis did
141 occur for all the characters studied. Hence, it is obvious that increase in yield of F₁ hybrids is the
142 result of increase in values of other yield contributing characters. Similar results were also
143 reported by Thomas *et al.* [11], Kumar *et al.* [12] and Choudhary *et al.* [13].

144 Generally, the heterotic expression decreases in F₂ generation as the dominance or
145 dominance interaction effects dissipate in this generation due to decreased heterozygosity,

Comment [Rk4]: Remove the before table3

146 resulting into inbreeding depression. For grain yield the crosses Raj 3077 x HD 3059, PBW 590
147 x HD 2967 and HD 2967 x WH 1021 revealed significant and negative inbreeding depression,
148 indicated F₂ plants accomplished higher grain yield per plant as compared to F₁ hybrids. The
149 significant and negative inbreeding depression for grain yield and other component characters
150 reported by several earlier researchers such Kumar *et al.* [12] and Nagar *et al.* [14].

151 Inbreeding depression based on genetic variability indicated the positive and negative
152 expression of genes in the population which could not be fixed for heterosis breeding. The
153 heterosis and inbreeding depression jointly implement the criteria for positive selection. As
154 indicated in the present study the succeeding generations showed the significant amount of
155 dominance in the population with accumulation of additive and epistatic genes. Bailey *et al.* [15]
156 observed that F₂ progenies performance for inbreeding depression might be a good indicator of
157 predicting heterosis performance in F₁ hybrid of wheat.

Table 1: Analysis of variance (mean squares) for parents, F₁'s and F₂'s for yield and its contributing traits

Characters	Source of variation								
	Replication	Genotypes	Parents	Generation	F ₁ 's	F ₂ 's	F ₁ vs F ₂	Parents vs generation	Error
Df	(2)	(99)	(9)	(89)	(44)	(44)	(1)	(1)	(198)
Days to 50% heading	0.57	101.46**	41.49**	107.83**	86.49**	130.69**	40.83**	73.67**	3.5
Days to maturity	0.37	131.06**	30.81**	139.88**	119.41**	160.65**	126.76**	248.43**	7.62
Plant height	39.2	242.74**	97.27**	258.57**	214.92**	302.45**	248.39**	143.69**	20.54
Productive tillers/plant	0.43	5.68**	2.07**	5.78**	3.81**	7.79**	3.95**	29.46**	0.37
Flag leaf area	10.04	218.75**	59.01**	232.29**	216.42**	252.86**	25.35*	452.01**	5.5
Spike length	0.99	6.98**	1.62**	7.48**	6.51**	8.54**	3.73**	10.81**	0.43
Number of grains/spikes	0.06	77.59**	23.98**	81.12**	66.46**	96.8**	36.36*	245.84**	5.77
1000-grain weight	2.58	21.96**	9.4**	21.72**	20.39**	23.36**	7.9*	156.3**	1.57
Biomass/plant	5.23	93.31**	37.11**	92.11**	67.81**	117.19**	57.57*	706.03**	10.79
Grain yield/plant	1.01	63.51**	13.75**	65.99**	60.85**	69.49**	138.07**	290.83**	4.2
Grain yield/spike	0.01	0.64**	0.41**	0.64**	0.56**	0.73**	0.18**	2.4**	0.02
Harvest index	0.25	109.41**	16.12*	119.25**	100.84**	120.89**	857.07**	73.11**	8.33

*, ** significant at 5 and 1% per cent levels, respectively

Table 2: Magnitude of heterosis (H), heterobeltiosis (HB) and inbreeding depression (ID) for different traits in bread wheat

Traits	Range of heterosis							Number of cross combinations showing heterosis		Number of cross combinations showing Inbreeding depression
	Heterosis (%)		Heterobeltiosis (%)			Inbreeding depression (%)		Heterosis (%)	Heterobeltiosis (%)	
	Min.	Max.	Min.	Max.	Best cross with heterotic effect	Min.	Max.			
Days to 50% heading	-12.79	11.35	-8.02	13.04	PBW 343 x WH 1021	-16.67	3.51	27	15	-
Days to maturity	-10.11	8.47	-8.33	10.22	PBW 343 x WH 1021	-10.54	5.15	26	15	1
Plant height	-15.11	14.87	-14.04	22.38	HD 2967 x WH 1021	-28.89	10.79	24	20	1
Productive tillers/plant	-10.20	31.01	-16.98	21.71	Raj 3765 x HD 2967	-11.07	29.07	30	20	-
Flag leaf area	-36.24	41.39	-46.15	31.86	PBW 343 x WH 1021	-16.48	17.25	27	18	5
Spike length	-21.61	23.22	-27.19	15.95	Raj 3765 x WH 1021	-5.57	16.43	25	15	-
Number of grains/spikes	-17.16	14.59	-18.80	13.10	Raj 3765 x UP 2425	-10.55	6.38	24	13	1
1000-grain weight	-7.31	13.48	-8.05	10.00	PBW 343 x HD 2967	-6.18	5.58	28	15	5
Biomass/plant	-9.33	23.42	-11.96	19.92	WH 1021 x UP 2425	-12.36	10.26	30	17	2
Grain yield/spike	-12.01	47.96	-21.65	29.86	PBW 343 x WH 1021	-4.70	17.13	30	15	-
Grain yield/plant	-26.22	38.73	-34.21	31.26	Raj 4079 x Raj 3765	-16.19	10.79	28	14	5
Harvest index	-29.26	15.10	-31.56	13.98	PBW 590 x HD 2967	-27.02	9.98	16	9	20

Table 3: Promising hybrids identified on the basis of per se performance and heterobeltiosis for grain yield per plant

S. No.	Hybrids	<i>Per se performance of grain yield per plant (g)</i>	Heterobeltiosis (%)	Significant heterobeltiosis for other traits in desired direction
1.	Raj 4079 x Raj 3765	32.68	31.26	DH, DM, PH, PTP, FLA, SL, GS, GW, BY, GYS, HI
2.	Raj 3765 x PBW 343	31.13	30.24	DH, DM, PH, PTP, FLA, SL, GS, GW, BY, GYS, HI
3.	Raj 3765 x WH 1021	28.85	29.86	DH, DM, PH, PTP, FLA, SL, GS, GW, BY, GYS, HI
4.	PBW 343 x HD 2967	31.04	29.33	DH, DM, PH, PTP, FLA, SL, GW, BY, GYS, HI
5.	Raj 4079 x HD 2967	32.06	28.77	DH, DM, PH, PTP, FLA, SL, GS, GW, BY, GYS, HI
6.	Raj 3765 x HD 2967	30.78	28.24	DH, DM, PH, PTP, FLA, SL, GS, GW, BY, GYS, HI
7.	Raj 4079 x PBW 343	31.59	26.88	DH, DM, PH, PTP, FLA, SL, GS, GW, BY, GYS, HI

DH: Days to 50 % heading, DM : days to maturity, PH : plant height, PTP : productive tillers per plant, FLA : flag leaf area, SL : Spike length, GS : number of grains per spike, GW : 1000-grain weight, BY : biomass per plant, GYS : grain yield per spike and HI : harvest index

References

1. Anonymous. Progress report of all India coordinated wheat and barley improvement project. Indian Institute of Wheat & Barley Research, Karnal, India; 2021-22a.
2. Wang L, Greaves IK, Groszmann M, Wua LM, Dennisa ES, Peacock WJ. Proceeding of the National Academy of Sciences, 2015 : E4959–E4967.
3. Fonseca S, Patterson FL.. Hybrid vigour in a seven parental diallel cross in common winter wheat (*Triticum aestivum* L.). Crop Sci. 1968 ; 51(9): 623-626.
4. Panse VC, Sukhatme PV. Statistical methods for agricultural workers. Published by ICAR, New Delhi; 1985.
5. Zare-kohan M, Heidari B. 2012. Estimation of genetic parameters for maturity and grain yield in diallel crosses of five wheat cultivars using two different models. *J. Agric. Sci.* 2012 ; 4(8): 74-85.
6. Singh K, Sharma SN, Sharma Y, Tyagi, BS.. Combining ability for high temperature tolerance and yield contributing traits in bread wheat. *J. Wheat Res.* 2012; 4(1): 29-37.
7. Nageshwar, Singh SV, Singh M, Singh L, Kumar S, Nan Kumar N, Singh AK. 2021. Selection of good combiner for further crop improvement by diallel analysis for central plan zone in winter wheat (*Triticum aestivum* L.). *The Pharma Innov. J.* ;10(12): 910-921.
8. Arunachalam V, Bandhyopadhyay A, Nigam SN. and Gibbons RW. Heterosis in relation to genetic divergence and combining ability in groundnut (*Arachis hypogaea* L.). *Euphytica.* 1984 ; 33(1): 33-39.
9. Gaur SC, Singh SN, Tiwari LP, Gaur LB. Heterosis and inbreeding depression in the inheritance of grain yield and its components in wheat (*Triticum aestivum*). *Current Advances in Agricultural Sciences*, 2014; 6(2): 186-189
10. Kumar A, Kumar V, Kerkhi SA, Kumar S, Chand P, Kumar N, Kumar D, Kumar M. Evaluation of heterosis for yield and yield related traits in bread Wheat (*Triticum aestivum* L.). *Progressive Agriculture*, 2014; 14 (1): 151-159.
11. Thomas N, Marker S, Lal GM, Dayal A. Study of heterosis for grain yield and its components in wheat (*Triticum aestivum*) over normal and heat stress condition. *Journal of Pharmacognosy and Phytochemistry*, 2017; 6(4): 824-830.
12. Kumar A, Razdan AK, Sharma V, Kumar N, Kumar D.. Study of heterosis and inbreeding depression for economic and biochemical traits in bread wheat (*Triticum aestivum* L.). *Journal of Pharmacognosy and Phytochemistry*, 2018; 7(4): 558-564.

13. Choudhary M, Singh, H., Punia, S. S., Gupta, D., Yadav, M., Get, S. and Bijarania, S. 2022. Estimation of heterosis for grain yield and some yield components in bread wheat (*Triticum aestivum* L. Em. Thell.). *The Pharma Innovation Journal*, **11**(2): 611-614.
14. Nagar SS, Kumar P, Singh C, Gupta V, Singh G, Tyagi BS. Assessment of heterosis and inbreeding depression for grain yield and contributing traits in bread wheat. *Journal of Cereal Research* 2019 ; 11(2): 125-130.
15. Bailey TB, Qualset CO, Cox DF. Predicting heterosis in wheat. *Crop Science*. 1980; 20:339-342.

UNDER PEER REVIEW

UNDER PEER REVIEW

