

Agronomic Zn Biofortification of Cereal Crops a Sustainable Way to Ensuring Nutritional Security: A Review

Abstract

Nutritional deficits in humans and animals constitute a hidden epidemic in many impoverished areas across the world. The staple foods of developing South Asian and African nations, such as rice, wheat, and maize, are poor in micronutrients. In recent past, a lack of food diversification i.e., cereal-based crops low in minerals, is another danger to nutritional quality and security. Because of the inherently low-level accumulation of nutrients in cereal crops, they are the primary target for bio-fortification among all crops. Among different micronutrients, zinc (Zn) is an important micronutrient that plays a vital role in a variety of physiological functions, and its scarcity will result in lower crop yields and productivity. Agronomic practices like application of fertilizers in soil, nutri-priming, foliar spray etc enhance the availability and uptake of Zn in crops. As a result, the growth and development, quality parameters and yield attributes of crop enhanced significantly. Therefore, agronomic biofortification of Zn in cereal crop is utmost important to achieve nutritional quality and food security. Furthermore, biofortification boosted the crop productivity to alleviate hidden hunger, in addition to quality aspects, proving to be a sustainable and cost-effective strategy. With soil and foliar fertiliser applications, including amendments, the agronomic interventions boost the Zn concentration in cereal crops. In this review the importance of agronomic Zn biofortification as a procedure to improve cereal yield and as an agricultural solution to solve nutritional quality and food security challenges is discussed.

Keywords: Agronomic interventions, Food security, Nutritional quality, Zn Biofortification.

1. Introduction

The ever-increasing population poses a global threat to the nutritional security. Nutrition security is defined as “one when all the people at all the time consume food of sufficient quantity and quality in terms of variety, diversity, nutrient content and safety to meet their dietary needs and food preferences for an active and healthy life, coupled with a sanitary environment, adequate health, education and care” (Anonymous, 2012). This issue is significantly more serious in low-income countries since farmers there use extensive agricultural approaches to boost productivity and profitability. Imbalance diets, a lack of variety in food sources, eating foods with lesser nutritional value, and food insecurity all have a negative impact on human health (Gundersen and Ziliak, 2015, Praharaj et al, 2021). Thus, nutritional security is utmost important for human being as well as animal. In the recent past years, the micronutrient deficiency become widespread across the world (Ernawati et al., 2021). Micronutrient deficiency is not only limited to developing countries but it also becoming prevalent where the staple food is cereals (Malik and Khan, 2021). The micronutrient insufficiency in soil restricting nutrient uptake in plants and, eventually, humans. Inadequate intake of these micronutrients has significant biological

consequences because they play such an important role in the human body's functioning (Allai et al., 2022). For plants, 17 mineral elements are deemed important, whereas for humans, the number exceeds 22 (White and Broadley, 2005). Out of those nutrients, deficiency of essential micronutrients such as Fe and Zn affects more than 2 billion individuals around the world (Huang et al., 2020, Mandi et al., 2021, Prasad and Shivay, 2020). Globally, Zn deficient soils are much more widespread than that of other micronutrient deficiencies. Zn constitute about 3000 different proteins and various classes of enzymes (Malik and Khan, 2021). Low Zn uptake may pose a threat to 60–70% of the population in Asia and Sub-Saharan Africa (Praharaj et al., 2021). Zn deficiency causes lead in DNA damage, a weaker immune system, cancer, heart disease, issues during female germ cell pregnancy, and other issues in humans (Gibson, 2012, Hotz and Brown, 2004). Besides this, mostly in developing low-income Asian and African countries cereals are staple food, accounting 55% of the dietary energy (Anonymous, 2008). Due to their maximum consumption and availability per person (464.6 g person⁻¹ day⁻¹) as well as their reactivity to micronutrient fertilisation, cereal grains are thought to have the most potential to serve as the finest carrier of micronutrients for relieving malnutrition (Shivay et al., 2008a, Mandi et al., 2021). Furthermore, wheat, rice, and maize are the three cereals that currently provide up to 60% of the daily energy intake of human populations (Tilman et al., 2002,). Low dietary intake, which is linked to a high consumption of cereal-based meals, is a common cause of zinc deficiency (Yaseen and Hussain, 2021). Cereal-based foods have very low Zn contents when compared to animal-based foods (Cakmak and Kutman, 2018). An adult's man daily Zn requirement is 9–18 mg day⁻¹, which cannot be met by cereal-based diet low in Zn content (Malik and Khan, 2021). When staple foods like wheat, rice, and maize are bio-fortified with zinc, it could significantly reduce nutritional insecurity because these foods are consumed by a huge number of people worldwide (Bouis and Welch, 2010, Shahane and Shivay, 2022, Wessells and Brown, 2012). Therefore, it is the need of the hour to increase the uptake and bioavailability of Zn in cereal crops to eradicate the nutritional security and improve the food quality. There are many options to improve nutritional security, including dietary diversification, food fortification, and medical supplementation. In fact, bio-fortification is a useful strategy for reducing malnutrition (De Valença et al., 2017, Saeid et al., 2019). Thus, to bridge the gap between nutrient supply and lower availability of Zn in the cereal grain, agronomic bio-fortification of Zn could be a novel approach to enhance the nutritional security.

2. Global status of Zn deficiency

In order to maintain better health, one must consume enough calories per day and a variety of foods to meet supplemental nutritional demands. Undernourishment is a major risk factor for death and other negative health effects, especially in children and mothers. In 2018, 9.2% of the world population were defined as severely food insecure. As a share of the population, food insecurity is highest in Sub-Saharan Africa where nearly one-third are defined as severely insecure (Anonymous, 2019). The highest prevalence of undernourishment was in Africa followed by Asia (Figure 1). Globally the status of Zn deficiency varies and it is shown in Figure 2. Country with highest population of Zn deficiency was found in Tazikistan (Figure 2). In high-income regions such as North America, Europe, Oceania, and Central Asia, Zn insufficiency affects less than 5–10% of the population, whereas it affects 15–50% of the population in Sub-Saharan Africa and South Asia, with the greatest prevalence of 54% in the Democratic Republic of Congo (Dhaliwal et al., 2022). It is projected that 60–70% of the population in Asia and Sub-Saharan Africa may be deficient in zinc (Gibson, 2006). This numbers to 2 billion

people in Asia and 400 million people in Sub-Saharan Africa in absolute terms (Anonymous, 2006). Zn deficiencies is estimated to be associated for around 4% of the global burden of morbidity and mortality in children under the age of five (Walker et al., 2009, Prasad et al., 2014). Pregnancy difficulties, low birth weight, reduced immunological competence, mother and infant mortality and morbidity, and development failure in infancy and childhood may all be linked to zinc deficiency. Zn deficiency is also widespread in different types of soil. Its deficiency is mostly prevalent in sodic soil, saline soil, calcareous soil, paddy soil, sandy soil, and highly weathered soil (Singh et al., 2005). The Zn deficiency first detected in rice crop at Pantnagar, Uttarakhand in 1965 by Nene (1966). Currently, 48% of Indian soils are deficient in zinc, with that number anticipated to rise to 63% by 2025 (Singh, 2009). In India, about 10 mha area are affected by zinc deficiency, and about 85% of rice-wheat system cropping occurs in the Indo-Gangetic plain, which has calcareous soils with high pH and consequently limited Zn availability. Improving the production from this cereal belt is consequently critical for the country's grain production to maintain sustainability (Singh et al., 2005). Based on the foregoing, it has been theorized that human Zn deficiency is linked to soil Zn content and relative cereal intake in the diet.




Figure 1: Prevalence of undernourishment in the world (Anonymous, 2020)




Figure 2: Countries with highest population of Zn deficiency (Anonymous, 2015)

3. Functions vis-à-vis deficiency symptoms of Zn in plant system

Zn plays a vital role in plant nutrition. Zn influence hydrogenase and carbonic anhydrase activity, ribosomal fraction stabilisation, and cytochrome synthesis in plants (Tisdale et al., 1984). It plays a role in carbohydrate metabolism. The enzyme involved is carbonic anhydrase. Zn helps in detoxification of superoxide radicle, the enzyme involved is copper zinc superoxide dismutase. It has a role in anaerobic root respiration in rice. The enzyme involved is ADH (alcohol dehydrogenase). Zn imparts resistance to disease in plants (Cakmak, 2000). Zn is required for the synthesis of tryptophan, which is a precursor to IAA, and it also plays a role in the creation of auxin, an important growth hormone (Alloway et al., 2004, Brennan et al., 1993). The Zn is also necessary for maintaining the integrity of the cellular membrane in order to keep macromolecules and ion transport systems in their structural orientation. Its interaction with phospholipids and membrane proteins' sulphhydryl groups aids in membrane preservation (Kabata-Pendias et al., 2004). Zn also aids nitrogen metabolism, chlorophyll production, protein synthesis, and photosynthesis, as well as providing resilience to stress. Zn also increases nitrogen metabolism, chlorophyll production, protein synthesis, and photosynthesis, as well as providing tolerance to stress (Hassan et al., 2021, Hassan et al., 2020). As Zn is relatively immobile in plant system, its deficiency symptoms generally appear on the growing young tissue. In rice, the characteristic symptom of Zn deficiency is bronzing. Zinc deficiency appear 2–3 weeks after rice seedlings are transplanted, with leaves developing brown blotches and streaks that may fuse to completely cover older leaves, and plants remaining stunted, whereas in severe cases, the plants may die, and those that recover will show significant delay in maturity and yield reduction (Sajwan and Lindsay, 1988, Yoshida and Tanaka, 1969, Prathap et al., 2022). Because Zn is immobile in deficit, the deficiency symptoms show first on young leaves. The upper leaf surfaces acquire interveinal chlorosis and necrotic spots, which eventually unite to form brown necrotic and brittle patches (Brennan et al., 1993).

4. Functions vis-à-vis deficiency symptoms of Zn in human physiology and health

Zn plays a pivotal role in biological system. Zn has several functions in human physiology i.e., reproduction system, nervous system, endocrine system, heart disease and hyper tension (Table 1). It is essential for the expression of genetic potential; nucleic acid synthesis, repair, and structural integrity all require Zn. As a result, it's no surprise that a lack of Zn stunts growth in practically all biological systems by reducing cell multiplication (Fairweather-Tait, 1988). The human body has 1.5–2.5 g of zinc, with 60% of it found in muscle and roughly 30% in bones. Zn is advised at 11 mg day⁻¹ for adult men and 8 mg day⁻¹ for adult women (Brown et al., 2001). Zn is also engaged in membrane stabilisation and cellular protection by preventing lipid peroxidation and lowering free radical production. The significance of zinc in reproduction has piqued people's interest (Apgar, 1985, Black et al., 2001). Poor Zn nutrition during pregnancy may cause low birth weight. Zn is such an essential element for human health that even a small shortage can be fatal. Loss of appetite, anorexia, loss of smell and taste, and other symptoms are caused by a lack of Zn in humans, and it can also impair the immune system, producing arteriosclerosis and anaemia (Gondal et al., 2021). Because it is a cofactor in the creation of numerous enzymes, DNA, and RNA, zinc is a vital element that serves a variety of activities in the body (Anonymous, 1996). Skin issues, recurrent infections, and dwarfism are all common in toddlers and school-aged children (Prasad et al., 2014, Roohani et al., 2013). Over 300 enzymes rely on zinc for their operation (Zastrow and Pecoraro, 2014). The important sources of Zn in human diet are marine foods such as crab, oysters, lobster, and other shellfish, red meat and poultry, dairy products such as cheese, legumes and nuts, whole grain cereals, and other sources (Prasad et al., 2016, Shahane et al., 2022). The amount of Zn absorbed from the food varies depending on a variety of dietary and physiological factors. The dietary factors which having affecting Zn bioavailability are phytate (plus calcium), high-fibre foods especially bran, histidine, hemicellulose, animal protein, oxalate (Kelsay, 1983), orange juice, ascorbic acid, iron etc, which are basically act as bioavailability inhibitors (Fairweather-Tait, 1988). In humans, a lack of Zn causes a variety of issues, including immune system dysfunction, skin difficulties, lower IQ levels, joint discomfort, memory loss, night blindness, and sexual maturation issues (Maret and Sandstead, 2008).

Table 1: Function of Zn in human physiology

Human physiology	Role of Zn	References
1. Reproduction system	Control the growth, fertility, and pregnancy of female germ cell, ovarian function, antral follicular development.	Garner et al. (2021), Foresta et al. (2014), Vickram et al. (2021)
2. Nervous system	Required for normal brain development and function, Increased blood-brain barrier permeability, hibits antagonist binding to muscarinic acetylcholine receptors,	Vallee and Falchuk (1993), Takeda (2001), Frederickson et al. (2001)
3. Endocrine system	thyroid hormones, androgens, and notably growth hormones metabolism, Zinc enhances circulation levels of IGF-I and potentiates IGF-I membrane signalling and intracellular second messengers that drive cell proliferation by influencing growth hormone	Favier (1992), MacDonald (2000), Salgueiro et al. (2002), Baltaci et al. (2019)

	synthesis and secretion as well as hepatic IGF-I synthesis.	
4. Diabetes mellitus	Involved in insulin secretion, storage, metabolism, and signaling.	Taylor (2005), Dunn (2005)
5. Heart disease and hypertension	Helps to defend against the oxidative stress that causes atherosclerosis in endothelial cells and reduce arterial hypertension.	Tubek (2007), Hennig et al. (1999)

5. Factors affecting Zn deficiency in plant system

Zinc deficiency can be seen in almost every country, and almost all crops react well to Zn application (Welch, 2002). Because over half of India's soils are deficient in accessible Zn, Zn shortage has gotten a lot of attention. There has been great emphasis on Zn fertilization in cereal crops in north-western India. Zn deficiency is occurred in almost all type of soil: submerged soil, sandy soil, loamy soil, soils formed from basalt, granite, leached soil etc. Zn deficiency is more prominent in calcareous and alkaline soil (Norouzi et al., 2014). The factors mostly affecting the availability of Zn are pH, calcium carbonate content, parent material, organic matter, soil moisture, soil temperature, plant species and genotype. The parent material one of the most important factors that influences Zn availability and determines Zn concentration in soils. The amount of Zn in soil is determined by the weathering of the rocks from which it is made (Rashid and Ryan, 2008, Wang and Busbey, 2005). The solubility of Zn is highly pH dependent and decrease 100 folds with each unit increase in soil pH. Zn deficiency is mostly occurred in alkaline soil rather than acid soil. Because soil Zn solubility is lower in alkaline soils, the availability of Zn is lowered. The soluble Zn in bulk soil generally varies from 4×10^{-10} M to 4×10^{-6} M (Barber, 1995) whereas in calcareous soil of rice-wheat cropping system of India soluble Zn concentration may be as low as 10×10^{-9} M (Hacisalihoglu and Kochian, 2003). Adequate soil moisture is utmost important in Zn uptake by plant. Lack of soil moisture results in Zn deficiency in soil. The most prominently Zn deficiency was observed in aerobic rice system (Prasad, 2011). Flooded soils are more commonly related with zinc deficiency than dry soils. Rice plants, for example, suffer from Zn shortage in calcareous soil when submerged. The interaction of zinc with free sulphide caused zinc shortage owing to floods. Because finer texture soils, such as clay, have higher CEC values, they have more reactive sites and can hold more Zn than lighter textured soils. Zn interacts positively with primary nutrient N and K as well as negatively with P (Prasad et al., 2016a). Zn reacts antagonistically with all the secondary nutrients. High concentration of Ca and Mg reduces the absorption and translocation of Zn in plant system (Prasad, 2011).

6. Factors affecting Zn deficiency in human

Zn deficiency in the human body is caused due to reduce bio-availability of Zn. Most chemicals reduce Zn availability by forming an unabsorbable combination with soluble Zn in the intestinal lumen. Phytic acid (myo-inositol hexa-phosphate), found in most cereal grains and seed legumes, is perhaps the most effective inhibitor of Zn absorption (Fairweather-Tait, 1988). Zn absorption is reduced primarily due to its high phytate content (Farah et al., 1984, Foster et al., 2012).

7. Response of cereal crops to Zn fertilization

Zinc (Zn) deficiency in cereal crops has become a major issue, resulting in lower yield and nutritional quality of the grain, harming human health. Agronomic management of cereal crops with Zn fertiliser has been reported as a promising method for improving seedling germination and production, as well as grain Zn concentration in cereal crops (NemeñoA, 2010, Phattarakul et al., 2016). Several parameters, including fertiliser supply, application timing and method, and soil chemical characteristics, have been reported to influence the efficacy of Zn fertiliser application (Wang et al., 2014). The adequate Zn fertilization application can help in increasing cereal production. Different response of Zn fertilization in cereal crops are presented in the Table 2.

Crops	Treatment	Increase in Zn concentration	References
1. Rice	Zn at 60 kg ha ⁻¹	Increased grain Zn content was 0.5 mg plant ⁻¹ (29.1%) when Zn fertilizer was applied	Tuiwong et al. (2021)
	4.0% ZnCU+0.2% ZnFS (ZnSO ₄ ·7H ₂ O)	enhanced the grain Zn concentration by 54.5% over control and 2.3–2.5 times higher Zn concentration was observed in grain as compared to straw.	Bana et al. (2021)
	Zn at 10 kg ha ⁻¹	When Zn was applied to wheat grain, it resulted in the highest level of Zn uptake (115.4 g ha ⁻¹).	Lakshmi et al. (2021)
	3.5% Zn (ZnO- enriched urea)	Zn concentration in aromatic rice increased by 16.4 mg kg ⁻¹ grain.	Shivay et al. (2008b)
	1% urea+0.5% ZnSO ₄ (N+Zn+)	Grain Zn concentration was the highest under foliar application of N+Zn+, with a 37.9% increase compared with control.	Tuiwong et al. (2022)
2. Wheat	8 mg Zn kg ⁻¹ soil	Increased grain Zn concentration of Zincol-2016 and Faisalabad-2008 by respectively 32 and 18% in industrial-zone soil, and by 15 and 2% in peri-urban soil.	Qaswar et al. (2017)
	Foliar application of 4 kg ZnSO ₄ ·7H ₂ O ha ⁻¹ , and soil application of 50 kg ZnSO ₄ ·7H ₂ O ha ⁻¹	Improved the grain Zn concentration of wheat by 28% and 89% during the first and second growing seasons, respectively.	Wang et al. (2014)

3. Maize	ZnO nanoparticles	Maximum increase in grain Zn concentration (82%)	Umar et al. (2021)
	Zinc oxide nanoparticles (ZnONPs 20 mg L ⁻¹)	Increase the Zn concentration 4.167 mg kg ⁻¹ in shoot.	Tondey et al. (2021)

8. Critical level of Zn in plant tissues

One of the main constraints to global food production is zinc deficiency. Identification of Zn-deficient areas and the various causes of shortage is therefore critical. According to Dobermann and Fairhurst (2000), the critical limits of Zn deficiency in plants are: 10 mg kg⁻¹ definite deficiency, 10–15 mg kg⁻¹ very likely, 15–20 mg kg⁻¹ likely, and >20 mg kg⁻¹ unlikely (sufficient). Zn sufficiency ranges from 15 to 50 ppm in the dry matter of mature plants in most crop species, with 15 ppm Zn regarded crucial in most cases.

9. Major approaches to achieve nutritional security

Three important strategies to achieving nutritional security are medical supplementation, dietary diversification and biofortification are presented in figure 3. (Dhaliwal et al., 2022). The first strategy, dietary diversity is a qualitative measure of food consumption that represents household access to a diverse range of foods and serves as a proxy for an individual's nutrient sufficiency. This strategy is involved in shift from monotonous staple starchy food crop to fruits, vegetable, pulses and animal diet which are rich in nutrition (Dhaliwal et al., 2022, White and Broadley, 2005). Furthermore, this strategy is suitable in regions where people have a wide variety of foods; nevertheless, it is not relevant in all areas like developing countries where the majority of people are depends on starchy staple food (Bouis and Welch, 2010). Nutrient supplementation meaning supply of nutrients which are not available in sufficient quantity in daily foods. Nutrient supplementation is done by vitamins, amino acid, minerals etc. When there is Zn deficiency, Zn supplementation can be done. Furthermore, this approach is commonly recommended during acute Zn shortage, particularly during pregnancy and early childhood (Gibson et al., 1998, Stein et al., 2007, Allen, 1998). However, Zn supplementation is quite expensive and is only recommended when a quick response is required (Stein et al., 2007). Biofortification is the process of increasing the micronutrient content in edible sections of a target crop without losing agronomic features such as pest resistance, yield, or drought resilience (Klikocka and Marks, 2018, Dhaliwal et al., 2022). Biofortification has the potential to solve the problem of micronutrient insufficiency (Bouis and Welch, 2010). Furthermore, it significantly increased crop yield in Zn deficient situations, and this method is thought to be a long-term solution to Zn insufficiency (Bouis and Welch, 2010). Different approaches of bio-fortification i.e., agronomic bio-fortification, genetic bio-fortification and genome engineering are presented in the figure 4.




Figure 3: Approaches to achieve nutritional security

10. Agronomic interventions for Zn biofortification

Agronomic biofortification is the deliberate application of mineral fertilisers to raise the proportion of a specific mineral in consumable crop parts in order to enhance dietary intake of the mineral (Prasad and Shivay, 2020, Shahane and Shivay, 2022). Agronomic biofortification is not a competitor to genetic biofortification. Different types of agronomic interventions involved in biofortification are soil application of fertilizers, foliar spraying of fertilizer, nanoparticle fertilizers, phyto-siderophores and EDTA chelated Zn, nutrient priming etc (Figure 4). The majority of management approaches are aimed at increasing soil quality, which improves micronutrient availability for plant uptake and consequently food concentration. The plant's nutrient usage efficacy is determined by a variety of elements including soil physical, chemical, and biological features. Agronomic biofortification is a quickest and cost-effective tool to improve the aforesaid characteristics. The application of fertilisers significantly enhanced grain Zn content, and it could be a quick solution to the problem of Zn insufficiency. Additionally, this method also boosts Zn availability and crop yield grain yield and Zn content in the grain (Yadav et al., 2011).




Figure 4: Different approaches of biofortification

10.1. Agronomic Zn biofortification via soil application of Zn

Micronutrient delivery through soil is the most adaptable and successful method due to its numerous advantages (Dhaliwal and Manchanda, 2009, Dhaliwal et al., 2012). Despite the fact that Zn deficit reduces crop productivity and grain Zn content, Zn deficiency can be addressed with Zn application. When substantial amounts of nutrients are required, the best technique for supplying them is by soil application. The treatment of high Zn is usually suggested for crops that are particularly vulnerable to Zn deficiency (Hassan et al., 2021). However, because of the reduced FUE, this approach necessitates a larger fertiliser treatment (Singh et al., 2005). Soil application of $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$ (21% Zn) at 62.5 kg ha^{-1} or $\text{ZnSO}_4 \cdot \text{H}_2\text{O}$ (33% Zn) at 40 kg ha^{-1} alleviate Zn deficiency in several crops (Ram et al., 2015).

10.2. Agronomic Zn biofortification via foliar feeding of Zn

Foliar micronutrient feeding increases grain Zn by reducing nutrient loss and allowing nutrients to be directly absorbed by plants (Hassan et al., 2021, Johnson et al., 2005). Zn sprayed foliarly increases grain Zn and bioavailability. Foliar-applied Zn is phloem-mobile and can easily be translocated into cereal grains during development (Haslett et al., 2001, Erenoglu et al., 2011). The time of foliar Zn fertiliser application determines its biofortification efficiency. Foliar Zn applications, in both wheat and rice, are more successful at enriching the grain with Zn if applied later rather than early in the growth process, preferably during grain filling (Boonchuay et al., 2013, Prasad et al., 2013). The fertiliser method for agronomic biofortification of wheat with zinc increases Zn concentrations in the endosperm, which is important for target groups who consume high amounts of white flour (Kutman et al., 2011). Even though a little amount of Zn is used in foliar feeding compared to soil applied Zn, foliar feeding is more effective in increasing the Zn content in grain and flour than soil application. The most essential and effective strategy for addressing Zn deficiency in cereals is foliar feeding. Furthermore,

because of the increase in direct Zn absorption in arid and semi-arid locations, foliar application is considered a unique application strategy (Chapagain and Wiesman, 2004).

10.3. Agronomic Zn biofortification via Seed priming

Seed priming is the regulated hydration of seeds to allow them to complete their pre-germination metabolic activities without causing radical emergence (Lutts et al., 2016). This promotes many physiological processes in plants to combat the damaging effects of abiotic stressors while also increasing the nutritional value and production of crops (Wojtyla et al., 2016, Kranner and Colville, 2011, Sundaria et al., 2018). Seed priming not only reduces the time it takes for seedlings to emerge, but it also allows for more uniform germination and a higher germination rate. Primed seeds have a greater potential for uniform stand establishment and yield than dry seeds (Farooq et al., 2006). The seed priming with Zn fertilisers improved wheat productivity and Zn uptake by seeds substantially (Farooq et al., 2012). Wheat seeds can be efficiently treated with ZnO NPs to improve Zn nutrition (Elhaj Baddar and Unrine., 2018). Seed priming with Zn is cost-effective, environmentally benign, and produces a large boost in yield; nonetheless, SP has occasionally proven to be a non-beneficial approach.

11. Agronomic Zn biofortification via nanotechnology

Nano-fertilizers are typically utilized for the controlled release of nutrients into the soil, which can improve the availability of nutrients to various plant organs, resulting in higher production and quality (Khan et al., 2021, Sekhon, 2014). Nano-fertilizers are much more beneficial of plant development and environmental safety than a comparable amount of traditional fertiliser due to their capacity to cover a larger surface area and their efficient absorption by plants. These are used in smaller amounts, resulting in less leaching and less gas emissions into the atmosphere (Manjunatha et al., 2016, Adisa et al., 2019). The effects of nano ZnO differ depending on the plant species, growth stage, application type and duration, and doses used. Zinc oxide NPs had a potential advantage over ZnSO₄ since they were less harmful to wheat crops at high concentrations, while also improving grain yield and Zn concentration (Du et al., 2019). Zinc hydroxide nitrate (Zn₅(OH)₈(NO₃)₂H₂O) has recently been investigated for use as a foliar spray. During the first growth stage, the novel source influenced Zn accumulation in maize stems and leaves, followed by its remobilization from the stems to other plant organs during the second development stage (Dhaliwal et al., 2022, Ivanov et al., 2019). The influence of weathered nano ZnO on grain Zn content was detected, weathered nano Zn showed 186 % increase in grain Zn content, whereas fresh nano ZnO showed a 229% increase (Dimkpa et al., 2020). Different bio-fortified cereal crops and their characteristics are discussed in the Table 3.

Sl. no.	Crop	Biofortified varieties	Characteristics
1.	Rice	CR Dhan 310 (Protein rich)	Contains 10.3% protein in polished grain as compared to 7.0–8.0% in popular varieties

		DRR Dhan 45 (Zn rich)	High in zinc content (22.6 ppm) in polished grains in comparison to 12.0–16.0 ppm in popular varieties.
2.	Wheat	WB 02 (Fe and Zn rich)	Rich in zinc (42.0 ppm) and iron (40.0 ppm) in comparison to 32.0 ppm zinc and 28.0–32.0 ppm iron in popular varieties.
		HPBW 01 (Fe and Zn)	high iron (40.0 ppm) and zinc (40.6 ppm).
3.	Maize	Pusa Vivek QPM9 Improved	High provitamin-A (8.15 ppm), lysine (2.67%) and tryptophan (0.74%).
		Pusa HM4 Improved	0.91% tryptophan and 3.62% lysine rich.
		Pusa HM8 Improved	tryptophan (1.06%) and lysine (4.18%) rich
		Pusa HM9 Improved	0.68% tryptophan and 2.97% lysine rich.
4.	Pearl millet	HHB 299	High iron (73.0 ppm) and zinc (41.0 ppm).
		AHB 1200	Rich in iron (73.0 ppm).

12. Agronomic biofortification in the light of cost effectiveness

Agronomic bio-fortification is widely used in various parts of the world, and this method has a major impact on grain Zn content. The agronomic fertilisation technique is a quick and cost-effective way to address the Zn deficiency (Bouis and Welch, 2010). Foliar feeding is a cost-effective strategy, however, due to the need to spray multiple times during the growth season, it can become uneconomical. Agronomic Zn biofortification is the most efficient and cost-effective approach to boost mineral and molecule content in food crops (Prasad and Shivay, 2020). Biofortification reduces the burden of micronutrient deficiencies, according to several research (De Steur et al., 2011). It also has to figure out how much the biofortification process costs in order to accomplish these burden reductions. Initial expenses for basic breeding and research to develop micronutrient-enriched biofortified lines are followed by marginal costs for testing, adaptive breeding, maintenance breeding, dissemination, and extension efforts in crop biofortification. According to a World Bank report from 1997, public health interventions that cost less than USD 150 DALY⁻¹ (Disability-Adjusted Life Years) saved are extremely cost-effective. Zn-enriched beans, rice, and wheat found that the bulk of the costs per DALY saved for biofortification fell into the extremely cost-effective category (Meenakshi et al., 2010).

13. Agronomic biofortification in the light of nutritional security and alleviating malnutrition

In comparison to non-biofortified crops, biofortified crops are nutritionally rich, assuming equivalent micronutrient bioavailability (La Frano et al., 2014). Because biofortified staple crops improve overall micronutrient intake and retention during cooking, processing, and storage. Nano-Zn particles improve the grain nutrient content in plants, making them a primary source of human health and easing malnutrition, their availability and level of accumulation should be closely regulated, as excessive intake can pose health hazards (El-Ramady et al., 2021, Rizwan et al., 2021). Many studies have shown that eating zinc-biofortified wheat starch boosts overall zinc absorption by 30–70% (Signorell et al., 2019, Lowe et al., 2020). As “Ending hunger, achieving food security and improving nutrition, and promoting sustainable agriculture” are the UN’s second sustainable development goals, which can be achieved through Zn biofortification of cereal crop. The use of biofortified cultivars has a lot of potential for improving human health and well-being. Several studies have shown that these biofortified crops have beneficial impacts on people (Yadava et al., 2018). The development and promotion of biofortified varieties thus would be helpful in addressing malnutrition and achieving the SDGs (Paroda et al., 2019). In keeping with the Sustainable Development Goals set, it is paramount that agronomically Zn biofortified cereal crops be developed and provided to the rural poor to alleviate malnutrition.

14. Conclusion

In developing and undeveloped countries, the great reliance of the human population on cereal-based diets is the primary source of micronutrient insufficiency. To overcome micronutrient insufficiency in the human population, agronomic biofortification by fertilisation appears to be a viable and cost-effective solution. Breeding techniques for bio-fortification might be a lengthy, costly, and resource-intensive process. Conversely, agronomic bio-fortification of cereal crops is a promising strategy that has significantly increased grain yield and grain Zn levels to fulfil human demands.

15. References

- Adisa, I.O., Pullagurala, V.L.R., Peralta-Videa, J.R., Dimkpa, C.O., Elmer, W.H., Gardea-Torresdey, J.L., White, J.C., 2019. Recent advances in nano-enabled fertilizers and pesticides: A critical review of mechanisms of action. *Environmental Science: Nano* 6(7), 2002–2030.
- Allai, F.M., Gul, K., Zahoor, I., Ganaie, T.A., Nasir, G., Azad, Z.R., 2022. Malnutrition: Impact of Zinc on Child Development. In *Microbial Biofertilizers and Micronutrient Availability*. Springer, Cham. ISBN 978-3-030-76608-5. ISBN 978-3-030-76609-2 (eBook). DOI 10.1007/978-3-030-76609-2-4, 87-100.
- Allen, L.H., 1998. Zinc and micronutrient supplements for children. *The American Journal of Clinical Nutrition* 68(2), 495S–498S.
- Alloway, T.P., Gathercole, S.E., Willis, C., Adams, A.M., 2004. A structural analysis of working memory and related cognitive skills in young children. *Journal of Experimental Child Psychology* 87(2), 85–106.
- Anonymous, 1996. Trace elements in human nutrition and health. World Health Organization (WHO). Available at <https://apps.who.int/iris/handle/10665/37931>. Accessed on 18th May, 2022.

- Anonymous, 1997. World Development Report 1997: The State in a Changing World. The World Bank. Available at <http://hdl.handle.net/10986/5980>. Accessed on 18th May, 2022.
- Anonymous, 2006. Bringing Hope, Improving Lives: Strategic Plan 2007–2015. International Rice Research Institute, Manila, Philippines, 61.
- Anonymous, 2008. Dietary, Diversity and Nutrition: The State of Food Insecurity in the World: Economic Growth is Necessary but not Sufficient to Accelerate Reduction of Hunger and Malnutrition. The Food and Agriculture Organization (FAO) of the United Nations, Rome. Available at <https://www.fao.org/3/i0291e/i0291e>. Accessed on 24th May, 2022.
- Anonymous, 2012. The State of Food Insecurity in the World 2012. FAO, Rome. Available at <https://reliefweb.int/report/world/state-food-insecurity-world-2012-economic-growth-necessary-not-sufficient-accelerate>. Accessed on 24th May, 2022.
- Anonymous, 2019. Hunger and Undernourishment. Available at OurWorldInData.org. <https://ourworldindata.org/hunger-and-undernourishment>. Accessed on 21st September, 2022.
- Anonymous, 2020. The State of Security and Nutrition in the World. FAO, Rome. Available at <http://www.fao.org/3/ca9692en/online/ca9692en.html#tab1>. Accessed on 14th May, 2022.
- Apgar, J., 1985. Zinc and reproduction. *Annual Review of Nutrition* 5(1), 43–68.
- Bana, R.C., Gupta, A.K., Bana, R.S., Shivay, Y.S., Bamboriya, S.D., Thakur, N.P., Puniya, R., Gupta, M., Jakhar, S.R., Choudhary, R.S., Bochalya, R.S., 2021. Zinc-coated urea for enhanced zinc biofortification, nitrogen use efficiency and yield of basmati rice under typical fluvents. *Sustainability* 14(1), 104.
- Barber, S.A., 1995. *Soil Nutrient Bioavailability: A Mechanistic Approach*. John Wiley & Sons, Inc., 605 Third Avenue, New York. ISBN 0-471-58747-8.
- Black, R.E., 2001. Micronutrients in pregnancy. *British Journal of Nutrition* 85(S2), S193–S197.
- Boonchuay, P., Cakmak, I., Rerkasem, B., Prom-U-Thai, C., 2013. Effect of different foliar zinc application at different growth stages on seed zinc concentration and its impact on seedling vigor in rice. *Soil Science and Plant Nutrition* 59(2), 180–188.
- Bouis, H.E., Welch, R.M., 2010. Biofortification- a sustainable agricultural strategy for reducing micronutrient malnutrition in the global south. *Crop science* 50(S1), 20S–30S.
- Brennan, R.F., Armour, J.D., Reuter, D.J., 1993. Diagnosis of zinc deficiency. In: Robson., A.D. (Ed.). *Zinc in Soils and Plants* (2nd Edn.). Springer, Dordrecht, 167–181.
- Brown, K.H., Wuehler, S.E., Peerson, J.M., 2001. The importance of zinc in human nutrition and estimation of the global prevalence of zinc deficiency. *Food and Nutrition Bulletin* 22(2), 113–125.
- Cakmak, I., 2000. Possible roles of zinc in protecting plant cells from damage by reactive oxygen species. *The New Phytologist* 146(2), 185–205.

- Cakmak, I., Kutman, U.A., 2018. Agronomic biofortification of cereals with zinc: A review. *European Journal of Soil Science* 69(1), 172–180.
- Chapagain, B.P., Wiesman, Z., 2004. Effect of Nutri-Vant-PeaK foliar spray on plant development, yield, and fruit quality in greenhouse tomatoes. *Scientia Horticulturae* 102(2), 177–188.
- De Steur, H., Gellynck, X., Blancquaert, D., Lambert, W., Van Der Straeten, D., Qaim, M., 2012. Potential impact and cost-effectiveness of multi-biofortified rice in China. *New Biotechnology* 29(3), 432–442.
- De Valença, A.W., Bake, A., Brouwer, I.D., Giller, K.E., 2017. Agronomic biofortification of crops to fight hidden hunger in sub-Saharan Africa. *Global Food Security* 12, 8–14.
- Dhaliwal, S.S., Manchanda, J.S., 2009. Critical level of boron in typic ustochrepts for predicting response of mungbean (*Phaseolus aureus* L.) to boron application. *Indian Journal of Ecology* 36(1), 22–27.
- Dhaliwal, S.S., Sadana, U.S., Khurana, M.P.S., Sidhu, S.S., 2012. Enrichment of wheat grains with zinc through Ferti-fortification. *Indian Journal of Fertilisers* 8(7), 34–45.
- Dhaliwal, S.S., Sharma, V., Shukla, A.K., Verma, V., Kaur, M., Shivay, Y.S., Nisar, S., Gaber, A., Brestic, M., Barek, V., Skalicky, M., 2022. Biofortification- A frontier novel approach to enrich micronutrients in field crops to encounter the nutritional security. *Molecules* 27(4), 1340.
- Dimkpa, C.O., Andrews, J., Sanabria, J., Bindraban, P.S., Singh, U., Elmer, W.H., Gardea-Torresdey, J.L., White, J.C., 2020. Interactive effects of drought, organic fertilizer, and zinc oxide nanoscale and bulk particles on wheat performance and grain nutrient accumulation. *Science of the Total Environment* 722, 137808.
- Dobermann, A., Fairhurst, T.H., 2000. *Rice: Nutrient Disorders & Nutrient Management* (2nd Edn.). International Rice Research Institute, Potash and Phosphate Institute. ISBN 981-04-2742-5.1-192.
- Du, W., Yang, J., Peng, Q., Liang, X., Mao, H., 2019. Comparison study of zinc nanoparticles and zinc sulphate on wheat growth: From toxicity and zinc biofortification. *Chemosphere* 227, 109–116.
- Dunn, M.F., 2005. Zinc-ligand interactions modulate assembly and stability of the insulin hexamer- A review. *Biometals* 18(4), 295–303.
- Elhaj Baddar, Z., Unrine, J.M., 2018. Functionalized-ZnO-nanoparticle seed treatments to enhance growth and Zn content of wheat (*Triticum aestivum*) seedlings. *Journal of Agricultural and Food Chemistry* 66(46), 12166–12178.
- El-Ramady, H., Abdalla, N., Elbasiouny, H., Elbehiry, F., Elsakhawy, T., Omara, A.E.D., Amer, M., Bayoumi, Y., Shalaby, T.A., Eid, Y., Zia-ur-Rehman, M., 2021. Nano-biofortification of different crops to immune against COVID-19: A review. *Ecotoxicology and Environmental Safety* 222, 112500.
- Erenoglu, E.B., Kutman, U.B., Ceylan, Y., Yildiz, B., Cakmak, I., 2011. Improved nitrogen nutrition enhances root uptake, root to shoot translocation and remobilization of zinc (⁶⁵Zn) in wheat. *New Phytologist* 189(2), 438–448.

- Ernawati, F., Syaury, A., Arifin, A.Y., Soekatri, M.Y., Sandjaja, S., 2021. Micronutrient deficiencies and stunting were associated with socioeconomic status in Indonesian children aged 6–59 months. *Nutrients* 13(6), 1802.
- Fairweather-Tait, S.J., 1988. Zinc in human nutrition. *Nutrition Research Reviews* 1(1), 23–37.
- Farah, D.A., Hall, M.J., Mills, P.R., Russell, R.I., 1984. Effect of wheat bran on zinc absorption. *Human nutrition: Clinical Nutrition* 38(6), 433–441.
- Farooq, M., Cheema, Z.A., Wahid, A., 2012. Seed priming with boron improves growth and yield of fine grain aromatic rice. *Plant Growth Regulation* 68(2), 189–201.
- Farooq, M., Tabassum, R., Afzal, I., 2006. Enhancing the performance of direct seeded fine rice by seed priming. *Plant Production Science* 9(4), 446–456.
- Favier, A.E., 1992. Hormonal effects of zinc on growth in children. *Biological Trace Element Research* 32(1), 383–398.
- Foresta, C., Garolla, A., Cosci, I., Menegazzo, M., Ferigo, M., Gandin, V., De Toni, L., 2014. Role of zinc trafficking in male fertility: From germ to sperm. *Human Reproduction* 29(6), 1134–1145.
- Foster, M., Karra, M., Picone, T., Chu, A., Hancock, D.P., Petocz, P., Samman, S., 2012. Dietary fiber intake increases the risk of zinc deficiency in healthy and diabetic women. *Biological Trace Element Research* 149(2), 135–142.
- Frederickson, C.J., Suh, S.W., Silva, D., Frederickson, C.J., Thompson, R.B., 2000. Importance of zinc in the central nervous system: the zinc-containing neuron. *The Journal of Nutrition* 130(5), 1471S–1483S.
- García-Gómez, C., Obrador, A., González, D., Babín, M., Fernández, M.D., 2017. Comparative effect of ZnO NPs, ZnO bulk and ZnSO₄ in the antioxidant defences of two plant species growing in two agricultural soils under greenhouse conditions. *Science of the Total Environment* 589, 11–24.
- Garner, T.B., Hester, J.M., Carothers, A., Diaz, F.J., 2021. Role of zinc in female reproduction. *Biology of Reproduction* 104(5), 976–994.
- Gibson, R.S., 2006. Zinc: the missing link in combating micronutrient malnutrition in developing countries. *Proceedings of the Nutrition Society* 65(1), 51–60.
- Gibson, R.S., 2012. Zinc deficiency and human health: Etiology, health consequences, and future solutions. *Plant and soil* 361(1), 291–299.
- Gibson, R.S., Yeudall, F., Drost, N., Mtitimuni, B., Cullinan, T., 1998. Dietary interventions to prevent zinc deficiency. *The American Journal of Clinical Nutrition* 68(2), 484S–487S.
- Gondal, A.H., Zafar, A., Zainab, D., Toor, M.D., Sohail, S., Ameen, S., Ijaz, A.B., Ch, B.I., Hussain, I., Haider, S., Ahmad, I.A., 2021. A detailed review study of zinc involvement in animal, plant and human nutrition. *Indian Journal of Pure & Applied Biosciences* 9(2), 262–271.

- Gundersen, C., Ziliak, J.P., 2015. Food insecurity and health outcomes. *Health affairs* 34(11), 1830–1839.
- Hacisalihoglu, G., Kochian, L. V., 2003. How do some plants tolerate low levels of soil zinc? Mechanisms of zinc efficiency in crop plants. *New Phytologist* 159(2), 341–350.
- Haslett, B.S., Reid, R.J., Rengel, Z., 2001. Zinc mobility in wheat: uptake and distribution of zinc applied to leaves or roots. *Annals of Botany* 87(3), 379–386.
- Hassan, M.U., Aamer, M., Nawaz, M., Rehman, A., Aslam, T., Afzal, U., Shahzad, B.A., Ayub, M.A., Ahmed, F., Qiaoying, M., Qitao, S., 2021. Agronomic bio-fortification of wheat to combat zinc deficiency in developing countries. *Pakistan Journal of Agricultural Research* 34(1), 201–217.
- Hassan, M.U., Aamer, M., Umer Chattha, M., Haiying, T., Shahzad, B., Barbanti, L., Nawaz, M., Rasheed, A., Afzal, A., Liu, Y., Guoqin, H., 2020. The critical role of zinc in plants facing the drought stress. *Agriculture* 10(9), 396.
- Hennig, B., Meerarani, P., Toborek, M., McClain, C.J., 1999. Antioxidant-like properties of zinc in activated endothelial cells. *Journal of the American College of Nutrition* 18(2), 152–158.
- Hotz, C., Brown, K.H., 2004. Assessment of the risk of zinc deficiency in populations and options for its control. *Food and Nutrition Bulletin* 25(1), S91–S204.
- Huang, S., Wang, P., Yamaji, N., Ma, J.F., 2020. Plant nutrition for human nutrition: hints from rice research and future perspectives. *Molecular Plant* 13(6), 825–835.
- Ivanov, K., Tonev, T., Nguyen, N., Peltekov, A., Mitkov, A., 2019. Impact of foliar fertilization with nanosized zinc hydroxy nitrate on maize yield and quality. *Emirates Journal of Food and Agriculture* 31(8), 597–604.
- Johnson, S.E., Lauren, J.G., Welch, R.M., Duxbury, J.M., 2005. A comparison of the effects of micronutrient seed priming and soil fertilization on the mineral nutrition of chickpea (*Cicer arietinum*), lentil (*Lens culinaris*), rice (*Oryza sativa*) and wheat (*Triticum aestivum*) in Nepal. *Experimental Agriculture* 41(4), 427–448.
- Kabata-Pendias, A., 2004. Soil–plant transfer of trace elements- An environmental issue. *Geoderma* 122(2–4), 143–149.
- Kang, B.T., Okoro, E.G., 1976. Response of flooded rice grown on a vertisol from northern Nigeria to zinc sources and methods of application. *Plant and Soil* 44(1), 15–25.
- Khan, W.A., Shabala, S., Cuin, T.A., Zhou, M., Penrose, B., 2021. Avenues for biofortification of zinc in barley for human and animal health: a meta-analysis. *Plant and Soil* 466(1), 101–119.
- Klikocka, H., Marks, M., 2018. Sulphur and nitrogen fertilization as a potential means of agronomic biofortification to improve the content and uptake of microelements in spring wheat grain DM. *Journal of Chemistry* 2018(spl.), 1–12.

- Kranner, I., Colville, L., 2011. Metals and seeds: Biochemical and molecular implications and their significance for seed germination. *Environmental and Experimental Botany* 72(1), 93–105.
- Kutman, U.B., Yildiz, B., Cakmak, I., 2011. Improved nitrogen status enhances zinc and iron concentrations both in the whole grain and the endosperm fraction of wheat. *Journal of Cereal Science* 53(1), 118–125.
- Lakshmi, P.V., Singh, S.K., Pramanick, B., Kumar, M., Laik, R., Kumari, A., Shukla, A.K., Abdel Latef, A.A.H., Ali, O.M., Hossain, A., 2021. Long-term zinc fertilization in calcareous soils improves wheat (*Triticum aestivum* L.) productivity and soil zinc status in the rice–wheat cropping system. *Agronomy* 11(7), 1306.
- Lowe, N.M., Zaman, M., Moran, V.H., Ohly, H., Sinclair, J., Fatima, S., Broadley, M.R., Joy, E.J., Mahboob, U., Lark, R.M., Zia, M.H., 2020. Biofortification of wheat with zinc for eliminating deficiency in Pakistan: Study protocol for a cluster-randomised, double-blind, controlled effectiveness study (BIZIFED2). *BMJ Open* 10(11), e039231.
- Lutts, S., Benincasa, P., Wojtyła, L., Kubala, S., Pace, R., Lechowska, K., Quinet, M., Garnczarska, M., 2016. Seed priming: New comprehensive approaches for an old empirical technique. In: Araujo, S., Balestrazzi, A. (Eds.). *New Challenges in Seed Biology-Basic and Translational Research Driving Seed Technology*. Intechopen, 1–46.
- MacDonald, R.S., 2000. The role of zinc in growth and cell proliferation. *The Journal of Nutrition* 130(5), 1500S–1508S.
- Malik, A., Khan, S.T., 2021. *Microbial Biofertilizers and Micronutrient Availability: The Role of Zinc in Agriculture and Human Health*. Springer Nature. ISBN 978-3-030-76608-5. ISBN 978-3-030-76609-2 (eBook). DOI 10.1007/978-3-030-76609-2.
- Mandi, S., Shivay, Y.S., Prasanna, R., Kumar, D., Purakayastha, T.J., Pooniya, V., Nayak, S., Raihan, O., Baral, K., Pal, M., Hussain, S., 2022. Improving micronutrient density in basmati rice and durum wheat through summer green manuring and elemental sulfur fertilisation. *Crop and Pasture Science* 73(8), 804–816.
- Manjunatha, S., Biradar, D., Aladakatti, Y.R., 2016. Nanotechnology and its applications in agriculture: A review. *Journal of Farm Sciences* 29, 1–13.
- Maret, W., Sandstead, H. H., 2006. Zinc requirements and the risks and benefits of zinc supplementation. *Journal of Trace Elements in Medicine and Biology* 20(1), 3–18.
- Meenakshi, J.V., Johnson, N.L., Manyong, V.M., DeGroote, H., Javelosa, J., Yanggen, D.R., Naher, F., Gonzalez, C., Garcia, J., Meng, E., 2010. How cost-effective is biofortification in combating micronutrient malnutrition? An ex ante assessment. *World Development* 38(1), 64–75.
- NemeñoA, G.A., 2010. Effect of water management on zinc concentration in rice grains. In: *Proceedings of the 19th World Congress of Soil Science*. Brisbane, Australia, 1–6th August.

- Norouzi, M., Khoshgoftarmanesh, A.H., Afyuni, M., 2014. Zinc fractions in soil and uptake by wheat as affected by different preceding crops. *Soil Science and Plant Nutrition* 60 (5), 670–678.
- Paroda, R.S., Joshi, P.K., 2019. Sustainable development goals: Role of agriculture. In: Chaturvedi, S., James, T.C., Saha, S., Shaw, P. (Eds.). *2030 Agenda and India: Moving from Quantity to Quality*. Springer, Singapore, 17–40. ISBN 978-981-32-9090-7. ISBN 978-981-32-9091-4 (eBook).
- Phattarakul, N., Rerkasem, B., Li, L.J., Wu, L.H., Zou, C.Q., Ram, H., Sohu, V.S., Kang, B.S., Surek, H., Kalayci, M., Yazici, A., 2012. Biofortification of rice grain with zinc through zinc fertilization in different countries. *Plant and Soil* 361(1), 131–141.
- Praharaj, S., Skalicky, M., Maitra, S., Bhadra, P., Shankar, T., Brestic, M., Hejnak, V., Vachova, P., Hossain, A., 2021. Zinc biofortification in food crops could alleviate the zinc malnutrition in human health. *Molecules* 26(12), 3509.
- Prasad, R., 2011. Aerobic rice systems. *Advances in Agronomy* 111, 207–247.
- Prasad, R., Shivay, Y.S., 2020. Agronomic biofortification of plant foods with minerals, vitamins and metabolites with chemical fertilizers and liming. *Journal of Plant Nutrition* 43(10), 1534–1554.
- Prasad, R., Shivay, Y.S., Kumar, D., 2013. Zinc fertilization of cereals for increased production and alleviation of zinc malnutrition in India. *Agricultural Research* 2(2), 111–118.
- Prasad, R., Shivay, Y.S., Kumar, D., 2014. Agronomic biofortification of cereal grains with iron and zinc. *Advances in Agronomy* 125, 55–91.
- Prasad, R., Shivay, Y.S., Kumar, D., 2016. Interactions of zinc with other nutrients in soils and plants- A review. *Indian Journal of Fertilisers* 12(5), 16–26.
- Prathap, S., Thiyareshwari, S., Krishnamoorthy, R., Prabhakaran, J., Vimalan, B., Gopal, N.O., Anandham, R., 2022. Role of zinc solubilizing bacteria in enhancing growth and nutrient accumulation in rice plants (*Oryza sativa*) grown on zinc (Zn) deficient submerged soil. *Journal of Soil Science and Plant Nutrition* 22(1), 971–984.
- Qaswar, M., Hussain, S., Rengel, Z., 2017. Zinc fertilisation increases grain zinc and reduces grain lead and cadmium concentrations more in zinc-biofortified than standard wheat cultivar. *Science of the Total Environment* 605, 454–460.
- Raj, A.B., Raj, S.K., 2019. Seed priming: An approach towards agricultural sustainability. *Journal of Applied and Natural Science* 11(1), 227–234.
- Ram, H., Sohu, V.S., Cakmak, I., Singh, K., Buttar, G.S., Sodhi, G.P.S., Gill, H.S., Bhagat, I., Singh, P., Dhaliwal, S.S., Mavi, G.S., 2015. Agronomic fortification of rice and wheat grains with zinc for nutritional security. *Current Science* 109(6), 1171–1176.
- Rashid, A., Ryan, J., 2004. Micronutrient constraints to crop production in soils with Mediterranean-type characteristics: A review. *Journal of Plant Nutrition* 27(6), 959–975.

- Rizwan, M., Ali, S., ur Rehman, M.Z., Riaz, M., Adrees, M., Hussain, A., Zahir, Z.A., Rinklebe, J., 2021. Effects of nanoparticles on trace element uptake and toxicity in plants: A review. *Ecotoxicology and Environmental Safety* 221, 112437.
- Roohani, N., Hurrell, R., Kelishadi, R., Schulin, R., 2013. Zinc and its importance for human health: An integrative review. *Journal of research in medical sciences. Isfahan University of Medical Sciences* 18(2), 144.
- Saeid, A., Patel, A., Jastrzębska, M., Korczyński, M., 2019. Food biofortification. *Journal of Chemistry* 2019, 1–2.
- Sajwan, K.S., Lindsay, W.L., 1988. Effect of redox, zinc fertilization and incubation time on DTPA-extractable zinc, iron and manganese. *Communications in Soil Science and Plant Analysis* 19(1), 1–11.
- Salgueiro, M.J., Zubillaga, M.B., Lysionek, A.E., Caro, R.A., Weill, R., Boccio, J.R., 2002. The role of zinc in the growth and development of children. *Nutrition* 18(6), 510–519.
- Sekhon, B.S., 2014. Nanotechnology in agri-food production: an overview. *Nanotechnology, Science and Applications* 7, 31.
- Shahane, A.A., Shivay, Y.S., 2022. Agronomic biofortification of crops: Current research status and future needs. *Indian Journal of Fertilisers* 18(2), 164–179.
- Shivay, Y.S., Kumar, D., Prasad, R., 2008a. Effect of zinc-enriched urea on productivity, zinc uptake and efficiency of an aromatic rice–wheat cropping system. *Nutrient Cycling in Agroecosystems* 81(3), 229–243.
- Shivay, Y.S., Prasad, R., Rahal, A., 2008b. Relative efficiency of zinc oxide and zinc sulphate-enriched urea for spring wheat. *Nutrient Cycling in Agroecosystems* 82(3), 259–264.
- Signorell, C., Zimmermann, M.B., Cakmak, I., Wegmüller, R., Zeder, C., Hurrell, R., Aciksoz, S.B., Boy, E., Tay, F., Frossard, E., Moretti, D., 2019. Zinc absorption from agronomically biofortified wheat is similar to post-harvest fortified wheat and is a substantial source of bioavailable zinc in humans. *The Journal of Nutrition* 149(5), 840–846.
- Singh, B., Natesan, S.K.A., Singh, B.K., Usha, K., 2005. Improving zinc efficiency of cereals under zinc deficiency. *Current Science* 88(1), 36–44.
- Singh, M.V., 2009. Micronutrient nutritional problems in soils of India and improvement for human and animal health. *Indian Journal of Fertilisers* 5(4), 11–16.
- Stein, A.J., Nestel, P., Meenakshi, J.V., Qaim, M., Sachdev, H.P.S., Bhutta, Z.A., 2007. Plant breeding to control zinc deficiency in India: how cost-effective is biofortification? *Public Health Nutrition* 10(5), 492–501.

- Sundaria, N., Singh, M., Upreti, P., Chauhan, R.P., Jaiswal, J.P., Kumar, A., 2019. Seed priming with iron oxide nanoparticles triggers iron acquisition and biofortification in wheat (*Triticum aestivum* L.) grains. *Journal of Plant Growth Regulation* 38(1), 122–131.
- Takeda, A., 2001. Zinc homeostasis and functions of zinc in the brain. *Biometals* 14(3), 343–351.
- Taylor, C.G., 2005. Zinc, the pancreas, and diabetes: Insights from rodent studies and future directions. *Biometals* 18(4), 305–312.
- Tilman, D., Cassman, K.G., Matson, P.A., Naylor, R., Polasky, S., 2002. Agricultural sustainability and intensive production practices. *Nature* 418(6898), 671–677.
- Tisdale, S.L., Nelson, W.L., Beaten, J.D., 1984. Zinc in soil fertility and fertilizers (4th Edn.). Macmillan Publishing Company, New York, 382–391.
- Tondey, M., Kalia, A., Singh, A., Dheri, G.S., Taggar, M.S., Nepovimova, E., Krejcar, O., Kuca, K., 2021. Seed priming and coating by nano-scale zinc oxide particles improved vegetative growth, yield and quality of fodder maize (*Zea mays*). *Agronomy* 11(4), 729.
- Tubek, S., 2007. Role of zinc in regulation of arterial blood pressure and in the etiopathogenesis of arterial hypertension. *Biological Trace Element Research* 117(1), 39–51.
- Tuiwong, P., Lordkaew, S., Prom-u-thai, C., 2021. Improving grain zinc concentration in wetland and upland rice varieties grown under waterlogged and well-drained soils by applying zinc fertilizer. *Agronomy* 11(3), 554.
- Tuiwong, P., Lordkaew, S., Veeradittakit, J., Jamjod, S., Prom-u-thai, C., 2022. Seed priming and foliar application with nitrogen and zinc improve seedling growth, yield, and zinc accumulation in rice. *Agriculture* 12(2), 144.
- Umar, W., Hameed, M.K., Aziz, T., Maqsood, M.A., Bilal, H.M., Rasheed, N., 2021. Synthesis, characterization and application of ZnO nanoparticles for improved growth and Zn biofortification in maize. *Archives of Agronomy and Soil Science* 67(9), 1164–1176.
- Vallee, B.L., Falchuk, K.H., 1993. The biochemical basis of zinc physiology. *Physiological Reviews* 73(1), 79–118.
- Vickram, S., Rohini, K., Srinivasan, S., Veenakumari, D.N., Archana, K., Anbarasu, K., Jeyanthi, P., Thanigaivel, S., Gulothungan, G., Rajendiran, N., Srikumar, P.S., 2021. Role of zinc (Zn) in human reproduction: a journey from initial spermatogenesis to childbirth. *International Journal of Molecular Sciences* 22(4), 2188.
- Walker, C.L.F., Ezzati, M., Black, R.E., 2009. Global and regional child mortality and burden of disease attributable to zinc deficiency. *European Journal of Clinical Nutrition* 63(5), 591–597.
- Wang, L.C., Busbey, S., 2005. Acquired acrodermatitis enteropathica. *New England Journal of Medicine* 352(11), 1121–1121.

- Wang, Y.Y., Wei, Y.Y., Dong, L.X., Lu, L.L., Feng, Y., Zhang, J., Pan, F.S., Yang, X.E., 2014. Improved yield and Zn accumulation for rice grain by Zn fertilization and optimized water management. *Journal of Zhejiang University Science B* 15(4), 365–374.
- Welch, R., 2002. The impact of mineral nutrients in food crops on global human health. *Plant and Soil* 247(1), 83–90.
- Wessells, K.R., Brown, K.H., 2012. Estimating the global prevalence of zinc deficiency: results based on zinc availability in national food supplies and the prevalence of stunting. *PLoS One* 7(11), e50568.
- White, P.J., Broadley, M.R., 2005. Biofortifying crops with essential mineral elements. *Trends in Plant Science* 10(12), 586–593.
- Wojtyła, Ł., Lechowska, K., Kubala, S., Garnczarska, M., 2016. Molecular processes induced in primed seeds—increasing the potential to stabilize crop yields under drought conditions. *Journal of Plant Physiology* 203, 116–126.
- Yadav, R.S., Patel, A.M., Dodia, I.N., Aglodiya, A.V., Patel, G.A., Augustine, N., 2011. Agronomic bio-fortification of wheat (*Triticum aestivum* L.) through iron and zinc enriched organics. *Journal of Wheat Research* 3(1), 46–51.
- Yadava, D.K., Hossain, F., Mohapatra, T., 2018. Nutritional security through crop biofortification in India: Status & future prospects. *The Indian Journal of Medical Research* 148(5), 621.
- Yaseen, M.K., Hussain, S., 2021. Zinc-biofortified wheat required only a medium rate of soil zinc application to attain the targets of zinc biofortification. *Archives of Agronomy and Soil Science* 67(4), 551–562.
- Yoshida, S., Tanaka, A., 1969. Zinc deficiency of the rice plant in calcareous soils. *Soil Science and Plant Nutrition* 15(2), 75–80.
- Zastrow, M.L., Pecoraro, V.L., 2014. Designing hydrolytic zinc metalloenzymes. *Biochemistry* 53(6), 957–978.