

Effect of Incorporation of Sewage Sludge and Fly ash on soil Physico-chemical Properties and Okra in Inceptisols of Prayagraj U.P

Abstract

The present research was conducted to determine whether it would be feasible to grow Okra in amended soil utilising sewage sludge and fly ash. Different types of nutrients and heavy metals were found in both soil amendments and for growth of the plant and crop yield they are used as ameliorate in acidic soils. In the research trial, soil properties like Bulk density, Particle density and pH are found to be positively non-significant and Pore space, Water holding capacity, EC, OC, Nitrogen, Phosphorus, Potassium, Iron, Manganese, Zinc and copper are found to be positively significantly low to medium range, which comprises yellowish brown sandy loam textured neutral to alkaline soil that is non-saline in nature among all the treatment combination applied.

Keyword : Sewage Sludge, Fly ash, soil , Okra , nutrients and heavy metals

Introduction

“Okra (*Abelmoschus esculentus* (L.) Moench] is an important annual fruit vegetable commonly grown in the tropics and warmer temperate regions of the world” (Patil *et al.*, 2015). “Okra is generally a self-pollinating crop belonging to the Malvaceae” (Oppong-Sekyere *et al.*, 2011). “It probably originated in Ethiopia and is widely spread all over tropical, subtropical and warm temperate regions of the world. Okra plays an important role in the human diet by supplying fats, proteins, carbohydrates, minerals and vitamins. Moreover, its mucilage is suitable for certain medical and industrial applications. therefore, young fruits of okra have reawakened beneficial interest in bringing this crop into commercial production. Okra requires warm temperatures. The optimum temperatures are in the range of 20-30°C, with minimum temperatures of 18°C and maximum of 35°C” (Liu *et al.* 2005, Kumar *et al.* 2009) “Okra needs rather high quantity of water despite having considerable drought resistance. The plant forms a deeply penetrating tap root with dense shallow feeder roots reaching out in all directions in the upper 45 cm of soil. The composition of okra pods per 100 g edible portion (81% of the product as purchased, ends trimmed) is: water 88.6 g, energy 144.00 kJ (36 kcal), protein 2.10 g, carbohydrate 8.20 g, fat 0.20 g, Fibre 1.70 g, Ca 84.00 mg, P 90.00 mg, Fe 1.20 mg, β -carotene 185.00 μ g, riboflavin 0.08 mg, thiamin 0.04

mg, niacin 0.60 mg, ascorbic acid 47.00 mg. Moreover, okra mucilage is suitable for industrial and medicinal applications” (Akinyele & Temikotan 2007). “The edible part of okra or its capsule (pod) measures approximately 15–20 cm in length and has a pyramidal-oblong, pentagonal, hispid appearance. Historically, okra pods were utilized for various purposes, such as in food, appetite boosters, astringents, and as an aphrodisiac. Furthermore, okra pods have also been recommended to cure dysentery, gonorrhoea, and urinary complications”(Islam *et al.*, 2019). “Extracts of young okra pods have also been reported to display moisturizing and diuretic properties, whereas the seeds of this plant have been reported to possess anticancer and fungicidal properties” (Durazzo *et al.*, 2018). In India characteristically, the sewage sludge contained total nitrogen (15400-1920 mg Kg⁻¹), available nitrogen (4600 -6300 mg Kg⁻¹), available phosphorus (44-60 mg Kg⁻¹) and available potassium (290-410 mg Kg⁻¹). Besides this higher contents of different heavy metals like arsenic (8-23 mg Kg⁻¹), Cd (2-9 mg Kg⁻¹), Cr (66-1098 mg Kg⁻¹), Hg (7-32 mg Kg⁻¹), Ni (12-596 mg Kg⁻¹) and Pb (26-154 mg Kg⁻¹) were reported by Dubey *et al.* (2006). Kumar and Chopra (2013) and Kumar *et al.* (2016) also reported higher contents of nutrients and heavy metals in the sewage sludge.

“Fly ash (FA), the by-product of coal combustion for energy production is a major environmental problem worldwide. FA management mainly includes its use for cement production, road-base construction, mine reclamation, mineral wool production, recovery of metals, and as aggregate substitute material. Recently FA has gained attention as a potential resource in terrestrial carbon sequestration. The incorporation of FA may also improve physical, chemical, and biological properties and thus benefit agricultural production” (Ukwattage *et al.* [2013](#)). “Sewage sludge (SS) from the other hand, a by-product of the sewage effluents treatment, constitutes also a serious environmental problem requiring a safe and economical disposal. Due to its composition and properties, SS may be beneficially used in agriculture as substitute of fertilizers or soil amendment. It was found that co-application of FA and SS may enhance the beneficial effects from the soil application of these two by-products. However, FA and SS as inputs may have many benefits for agriculture like facing of nutrient deficiencies, while these by-products contain a number of toxic substances like toxic heavy metals and organic pollutants. Therefore, a proper attention should be paid on the impacts of the use of these materials on soil health, crop quality and heavy metal toxicity and leaching”. (Sipkova *et al.*, 2014)

“Therefore, the management of both by-products, i.e., FA and SS, is of significant importance for India. Research work showed that both by-products FA and SS may be used in a beneficial and sustainable way facing thus serious environmental problems” (Samaras *et al.*, 2008; Tsadilas *et al.*, 2009). The purpose of the present study was to investigate the possibility to use FA and SS in Okra crop separately or together and its influence on yield and soil properties including essential nutrients content as well as toxic heavy metals accumulation.

Materials and Methods

The experiment was laid out at Research Farm of SSAC, SHUATS, Prayagraj, Uttar Pradesh which is analysed by randomised block design with 12 treatments and 3 replications. The treatments comprises of three levels of sewage sludge, i.e. 0, 13, 26, 39 and 52 tonns ha⁻¹ also levels of fly ash taken 0, 13, 26, 39 and 52 tonns ha⁻¹ incorporated as soil application with or without NPK according to the treatment combination. The soil samples were collected randomly from two depths after harvest of okra. The soil is characterized after incorporation of Sewage sludge and fly-ash and the properties are

Weight of oven dried soil (Mg)

a) Bulk density (Mgm⁻³) = Volume of soil (m⁻³)

Mass of soil solid (Mg)

b) Particle density (Mgm⁻³) = Volume of solids (m⁻³)

Bulk Density

c) % pore space = (1-Particle Density × 100)

Vol. of water absorbed by soil

d) Water holding capacity = Vol. of soil taken × 100

In chemical parameters through method by

e) Soil pH - by using Digital pH meter of globe instruments given by (Jackson, 1973)

f) Soil EC (dSm⁻¹)-Digital EC meter of globe instruments.

- g) Organic Carbon (%) - through titration given by Walkley and Black method (1934)
- h) Available Nitrogen (Kg ha^{-1})-Kjeldhal Method (Subbaih and Asija, 1956)
- i) Available Phosphorus (Kg ha^{-1})- Colorimetric method by using Jasper single beam U.V Spectrophotometer at 660 nm wavelength.
- j) Available Potassium (Kg ha^{-1})- Flame photometric method by using Metzer Flame Photometer.
- k) Available Fe, Mn, Zn and Cu (ppm)- Atomic Absorption Spectrophotometer by instrument perkinelmer given by (Lindsay and Norwell)

Results and Discussion-

Table 1- Effect of Incorporation of Sewage Sludge and Fly ash on soil Physical properties-

Treatment/ depth	D _b (Mgm ⁻³) (2020)		D _b (Mgm ⁻³) (2021)		D _p (Mgm ⁻³) (2020)		D _p (Mgm ⁻³) (2021)		Percent pore space (2020)		Percent pore space (2021)		Water Holding Capacity (%) (2020)		Water Holding Capacity (%) (2021)	
	0-15 cm	15-30 cm	0-15 cm	15-30 cm	0-15 cm	15-30 cm	0-15 cm	15-30 cm	0-15 cm	15-30 cm	0-15 cm	15-30 cm	0-15 cm	15-30 cm	0-15 cm	15-30 cm
T ₁	1.22	1.25	1.23	1.26	2.25	2.3	2.32	2.41	45.78	45.65	46.3	46.82	41.83	40.16	42.11	40.42
T ₂	1.08	1.13	1.10	1.15	2.12	2.15	2.18	2.27	48.06	47.44	49.08	50.04	45.43	43.61	45.38	43.57
T ₃	1.13	1.17	1.15	1.20	2.13	2.16	2.19	2.28	46.95	45.83	47.46	47.97	45.72	43.89	45.67	43.84
T ₄	1.14	1.18	1.16	1.22	2.16	2.21	2.22	2.31	47.22	46.61	47.73	48.24	43.71	41.96	43.66	41.92
T ₅	1.15	0.20	1.18	1.22	2.18	2.26	2.25	2.34	47.25	44.15	47.76	48.26	43.75	42.00	43.7	41.95
T ₆	1.10	1.12	1.13	1.15	2.13	2.18	2.19	2.28	48.36	47.62	48.86	49.35	45.21	43.40	45.17	43.36
T ₇	1.09	1.14	1.11	1.15	2.15	2.2	2.21	2.3	49.3	48.18	47.32	49.76	44.57	42.78	44.52	42.74
T ₈	1.20	1.22	1.22	1.23	2.21	2.27	2.28	2.37	45.7	46.26	46.23	46.75	42.16	40.47	42.82	41.10
T ₉	1.17	1.20	1.18	1.24	2.18	2.2	2.25	2.34	46.33	45.45	46.85	47.36	43.11	41.39	43.06	41.34
T ₁₀	1.16	1.19	1.18	1.21	2.17	2.23	2.24	2.32	46.54	46.64	47.06	47.57	43.08	41.35	43.03	41.31
T ₁₁	1.12	1.16	1.13	1.19	2.14	2.19	2.2	2.29	47.66	47.03	48.17	48.67	46.55	44.69	46.49	44.63
T ₁₂	1.07	1.11	1.09	1.14	2.1	2.13	2.16	2.25	49.31	48.32	49.54	50.3	47.05	45.17	47.00	45.12
F Test	NS	NS	NS	NS	NS	NS	NS	NS	S	S	S	S	S	S	S	S
SEm±	-	-	-	-	-	-	-	-	0.52	0.85	0.64	0.62	1.02	0.75	0.8	0.71
CD (P=0.05)	-	-	-	-	-	-	-	-	1.53	2.48	1.89	1.82	2.98	2.2	2.36	2.08

As depicted in the Table 1, shows that the effect of sewage sludge and fly ash on the physical properties of soil at both depth in both the years. The B.D (Mgm^{-3}) was non-significant at both the depths and in both the years which was maximum in 1.26 in 2021 at 15-30 cm and minimum was 1.07 in 2020 at 0-15 cm. The PD (Mgm^{-3}) was non-significant in both the depths and in both the years which was maximum in 2.41 in 2021 at 15-30 cm and minimum was 2.1 in 2020 at 0-15 cm. The Pore-space (%) was significant in both the depths and in both the years which was maximum in 50.3 in 2021 at 15-30 cm and minimum was 45.7 in 2020 at 15-30 cm. The WHC (%) was significant in both the depths and in both the years which was maximum in 47.0 in 2020 at 0-15 cm and minimum was 40.16 in 2021 at 15-30 cm.

Table 2- Effect of Incorporation of Sewage Sludge and Fly ash on soil Chemical properties-

Treatment /depth	pH (2020)		pH (2021)		EC (dS m ⁻¹) (2020)		EC (dS m ⁻¹) (2021)		OC (%) (2020)		OC (%) (2021)		Available Nitrogen (kg ha ⁻¹) (2020)		Available Nitrogen (kg ha ⁻¹) (2021)		Available Phosphorus (kg ha ⁻¹) (2020)		Available Phosphorus (kg ha ⁻¹) (2021)		Available Potassium (kg ha ⁻¹) (2020)		Available Potassium (kg ha ⁻¹) (2021)	
	0-15 cm	15- 30 cm	0-15 cm	15- 30 cm	0-15 cm	15- 30 cm	0-15 cm	15- 30 cm	0-15 cm	15- 30 cm	0-15 cm	15- 30 cm	0-15 cm	15-30 cm	0-15 cm	15-30 cm	0-15 cm	15-30 cm	0-15 cm	15-30 cm	0-15 cm	15-30 cm	0-15 cm	15-30 cm
T ₁	7.58	7.62	7.46	7.58	0.29	0.28	0.28	0.32	0.52	0.5	0.45	0.43	276.08	258.42	256.08	268.89	29.2	27.68	28.64	25.62	185.6	172.28	191.6	173.78
T ₂	7.33	7.45	7.35	7.41	0.28	0.27	0.32	0.36	0.5	0.48	0.52	0.48	257.16	240.68	267.46	252.95	20.42	20.16	19.42	18.75	179.16	167.24	195.36	175.64
T ₃	7.1	7.16	7.32	7.36	0.3	0.29	0.34	0.38	0.6	0.58	0.56	0.52	275.13	252.38	297.23	269.53	27.92	26.98	30.84	28.54	219.52	204.26	249.42	217.86
T ₄	7.18	7.24	7.42	7.47	0.25	0.24	0.3	0.34	0.62	0.6	0.55	0.46	279.8	265.98	302.68	277.65	30.7	29.02	29.35	27.78	241.42	232.17	258.62	246.97
T ₅	6.98	7.06	7.15	7.19	0.26	0.25	0.33	0.37	0.66	0.64	0.58	0.54	293.68	280.14	315.88	294.47	31.98	31.44	30.21	28.96	255.16	243.64	272.26	257.34
T ₆	6.88	7.01	7.22	7.26	0.28	0.27	0.31	0.36	0.73	0.71	0.53	0.51	309.63	295.16	330.43	309.53	34.46	36.52	34.68	32.88	263.82	256.87	281.62	264.57
T ₇	7.3	7.36	7.3	7.32	0.26	0.25	0.34	0.39	0.57	0.55	0.59	0.55	291.56	285.31	312.26	301.76	22.06	23.25	24.76	21.57	220.64	208.24	238.74	216.74
T ₈	7.31	7.41	7.42	7.5	0.27	0.26	0.32	0.37	0.66	0.64	0.62	0.57	329.54	320.14	350.34	336.68	28.55	27.28	28.54	24.55	251.41	238.16	269.31	244.96
T ₉	7.01	7.12	7.21	7.27	0.28	0.27	0.34	0.37	0.68	0.66	0.65	0.59	344.6	331.74	366.26	345.56	33.16	32.25	34.58	31.86	261.12	250.68	278.32	263.48
T ₁₀	6.92	7.05	7.19	7.23	0.31	0.3	0.36	0.4	0.73	0.71	0.67	0.61	368.2	350.54	381.82	368.37	36.23	36.26	37.35	33.21	290.4	277.61	309.14	285.41
T ₁₁	6.9	7.04	7.15	7.19	0.32	0.29	0.35	0.39	0.74	0.72	0.7	0.64	379.9	370.44	403.19	389.69	38.28	37.36	41.82	36.34	298.16	289.36	318.26	296.66
T ₁₂	7.29	7.41	7.36	7.51	0.34	0.32	0.37	0.41	0.78	0.75	0.76	0.66	407.92	403.67	430.02	417.25	46.64	46.18	44.89	39.78	317.31	305.42	335.21	317.52
F Test	NS	NS	NS	NS	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S
SEm±	-	-	-	-	0.00	0.00	0.01	0.00	0.01	0.01	0.01	0.01	4.65	3.86	4.69	4.02	0.55	0.46	0.44	0.40	3.52	4.36	4.29	4.42
CD (P=0.05)	-	-	-	-	0.01	0.01	0.02	0.01	0.03	0.02	0.02	0.02	13.64	11.32	13.76	11.80	1.61	1.35	1.28	1.18	10.32	12.80	12.58	12.96

As depicted in the Table 2, shows that the effect of sewage sludge and fly ash on soil chemical properties. The pH was non-significant in both the depths and in both the years which was maximum in 7.58 in 2020 at 15-30 cm and minimum was 6.88 in 2020 at 0-15 cm. The EC (dSm⁻¹) was significant in both the depths and in both the years which was maximum in 0.41 in 2021 at 15-30 cm and minimum was 0.25 in 2020 at 15-30 cm. The OC (%) was significant in both the depths and in both the years which was maximum in 0.78 in 2020 at 0-15 cm and minimum was 0.48 in 2020 at 15-30 cm. The Nitrogen (kg ha⁻¹) was significant in both the depths and in both the years which was maximum in 438.67 in 2021

at 0-15 cm and minimum was 252.95 in 2020 at 15-30 cm. The Phosphorus (kg ha^{-1}) was significant in both the depths and in both the years which was maximum in 46.64 in 2020 at 0- 15 cm and minimum was 20.42 in 2021 at 15- 30cm. The Potassium (kg ha^{-1}) was significant in both the depths and in both the years which was maximum in 335.21 in 2021 at 0-15 cm and minimum was 167.24 in 2021 at 15-30 cm.

Table 3- Effect of Incorporation of Sewage Sludge and Fly ash on Available Micro-nutrients in soil-

Treatment/ depth	Available Iron (mg kg^{-1}) (2020)		Available Iron (mg kg^{-1}) (2021)		Available Manganese (mg kg^{-1}) (2020)		Available Manganese (mg kg^{-1}) (2021)		Available Zinc (mg kg^{-1}) (2020)		Available Zinc (mg kg^{-1}) (2021)		Available Copper (mg kg^{-1}) (2020)		Available Copper (mg kg^{-1}) (2021)	
	0-15 cm	15-30 cm	0-15 cm	15-30 cm	0-15 cm	15-30 cm	0-15 cm	15-30 cm	0-15 cm	15-30 cm	0-15 cm	15-30 cm	0-15 cm	15-30 cm	0-15 cm	15-30 cm
T₁	6.62	4.4	6.951	4.62	6.22	5.42	6.53	5.69	1.04	1.01	1.29	1.06	2.04	1.12	2.142	1.8375
T₂	18.42	16.05	19.341	16.8525	13.24	11.04	13.90	11.59	1.23	1.84	1.09	0.88	2.24	1.87	2.352	1.9635
T₃	17.68	15.42	18.564	16.191	14.56	12.36	15.29	12.97	1.5	1.52	1.58	1.16	3.09	2.45	3.2445	2.5725
T₄	19.85	17.2	20.8425	18.06	15.42	13.34	16.19	14.01	2.45	1.9	2.57	1.99	3.85	2.86	4.0425	3.003
T₅	28.94	25.42	30.387	26.691	17.84	15.45	18.73	16.22	3.02	2.45	3.72	3.10	4.1	3.24	4.305	3.402
T₆	33.233	30.2	34.8946	31.71	19.46	17.24	20.43	18.10	3.12	2.32	4.03	3.19	3.9	3.01	4.095	3.1605
T₇	26.78	23.54	28.119	24.717	15.54	13.24	16.32	13.90	2.04	1.85	1.09	0.89	2.2	2.65	2.31	1.7325
T₈	21.2	19.85	22.26	20.8425	17.42	15.32	18.29	16.09	2.53	1.72	1.61	1.08	2.94	1.98	3.087	2.079
T₉	21.85	20.23	22.9425	21.2415	21.09	19.05	22.14	20.01	2.65	1.94	2.78	2.04	3.04	2.24	3.192	2.352
T₁₀	20.82	19.24	21.861	20.202	21.45	19.62	22.52	20.60	2.98	2.16	3.13	2.20	3.68	3.01	3.864	3.1605
T₁₁	30.54	27.34	32.067	28.707	22.34	20.45	23.46	21.47	3.18	2.48	3.25	2.52	3.94	3.36	4.137	3.36
T₁₂	36.54	33.08	38.367	34.734	25.84	23.64	27.13	24.82	3.54	2.85	3.71	2.99	4.34	3.78	4.557	3.969
F Test	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S
SEm±	0.40	0.25	0.53	0.40	0.16	0.24	0.31	0.23	0.05	0.04	0.04	0.04	0.05	0.04	0.06	0.05
CD (P=0.05)	1.16	0.75	1.55	1.17	0.46	0.72	0.91	0.67	0.13	0.10	0.13	0.11	0.16	0.13	0.19	0.13

As depicted in the Table 3, shows that the effect of sewage sludge and fly ash on soil micronutrients. The Iron was significant in both the depth and in both the years which was maximum in 38.36 in 2021 at 0-15 cm and minimum was 4.4 in 2020 at 15-30 cm. The Manganese was significant in both the depths and in both the years which was maximum in 27.13 in 2021 at 0-15 cm and minimum was 5.42 in 2020 at 15-30 cm. The Zinc was significant in both the depths and in both the years which was maximum in 4.03 in 2021 at 0-15 cm and minimum was 1.01 in 2020 at 15-30 cm. The Copper was significant in both the depths and in both the years which was maximum in 4.55 in 2021 at 0-15 cm and minimum was 1.12 in 2020 at 15-30 cm.

Conclusion-

The research study has revealed that the use of sewage sludge and fly ash has improved the soil physico-chemical properties, the combined use of FA and sewage sludge has been proposed to reduce the bioavailability of heavy metals these ameliorant has potential liming capabilities, decreasing pH and having a long-term residual effect. It can be seen as a slow-release supply of components needed for plant growth, as well as a good source of nutrients required for plant growth. At the same time some toxic metal at higher level reduce the productivity of yield. The high concentration of micro nutrient and macronutrient presents in sewage sludge and fly ash increases the yield of okra.

Reference

- Akinyele, B.O. and Temikotan, T. (2007): Effect of variation in soil texture on the vegetative and pod characteristics of okra (*Abelmoschus esculentus* (L.)). *International. Journal of Agriculture Research*. 2: 165-169.
- Dubey, S.K., Yadav, R.K., Chatuvedi, P.K., Goyel, B., Yadav, R. and Minhas, P.S. (2006). Agricultural uses of sewage sludge and water and their impact on soil water and environmental health in Haryana, India. Abstract of 18th *World Congress of Soil Science, Philadelphia*.
- Durazzo, A., Lucarini, M., Novellino, E., Souto, E.B., Daliu, P. and Santini, A. (2018) *Abelmoschus esculentus* (L.): Bioactive Components' Beneficial Properties- Focused on Antidiabetic Role-For Sustainable Health Applications. *Molecules*. 24-38.
- Islam, M.T. (2019) Phytochemical information and pharmacological activities of Okra (*Abelmoschus esculentus*): A literature-based review. *Phytother Research*. **33**:72–80.

- Jackson, M.L. (1956). Soil Chemical Analysis—Advance Course. University of Wisconsin. Madison, 991p.
- Kumar, R., Patil, M.B., Patil, S.R. and Paschapur, M.S. (2009): Evaluation of *Abelmoschus esculentus* mucilage as suspending agent in paracetamol suspension. *International Journal of Pharm Technology Research*. **1**: 658-66
- Kumar, V. and Chopra, A.K. (2013a). Accumulation and translocation of metals in soil and different parts of French bean (*Phaseolus vulgaris* L.) amended with sewage sludge. *Bulletin of Environmental Contamination and Toxicology*, **92 (1)**: 103-108.
- Kumar, V. and Chopra, A.K. (2016b). Effects of sugarcane pressmud on agronomical characteristics of hybrid cultivar of eggplant (*Solanum melongena* L.) under field conditions. *International Journal of Recycling of Organic Waste in Agriculture*.
- Lindsay, W.L. and Norvell, W.A. (1978) Development of a DTPA soil test for zinc, iron, manganese and copper. *Soil Science Society of America Journal*, **42**: 421-428.
- Liu, I.M., Liou, S.S., Lan, T.W., Hsu, F.L. and Cheng, J.T. (2005): Myricetin as the active principle of *Abelmoschus moschatus* to lower plasma glucose in streptozotocin-induced diabetic rats. *Planta Medica* **71**: 617-621
- Muthuve, P., Udayasoorian, C., Natesan, R. and Ramaswami, P.R. (1992) Introduction to Soil Analysis, Tamil Nadu Agricultural University, Coimbatore.
- Olsen, S.R., Cole, C.V., Watanabe, F.S. and Dean, L.A. (1954) Estimation of available Phosphorus in soils by extraction with sodium carbonate. *U.S.D.A. Cir.*, **933**:1-10.
- Oppong, S., D., Akromah, R., Nyamah, E. Y., Brenya, E., and Yeboah, S. (2011). Characterization of okra (*Abelmoschus* spp. L.) germplasm based on morphological characters in Ghana. *Journal of Plant Breeding & Crop Sciences*. **3**:368–379.
- Patil, P., Sutar, S., Joseph, J. K., Malik, S., Rao, S., and Yadav, S. (2015). A systematic review of the genus *Abelmoschus* (Malvaceae). *Rheedea* **1**: 14–30.
- Samaras, V., Tsadilas, C.D. and Stamatiadis, S. (2008) Effects of repeated application of municipal sewage sludge in a cotton field: effects on soil fertility, crop yield, and nitrate leaching. *Agronomy Journal* **100**:477–483
- Sipkova, A., Szakova, J. and Thustos, P. (2014) Affinity of selected elements to individual fractions of soil organic matter. *Water Air Soil Pollution*. **225**:1-11
- Subbiah, B. V. and Asija, G. L. (1956) A rapid procedure for the estimation of available nitrogen in soils. *Journal of Current Sciences*, **25(8)**: 259-260.

- Toth, S.J. and Prince, A.L. (1949) Estimation of Cation exchange capacity and exchangeable Ca, K and NA content of soil by flame photometer technique. *Soil Science*; **67**:439-445.
- Tsadilas, C., Shaheen, S., Samaras, V., Gizas, D. and Hu, Z. (2009) Influence of fly ash application on copper and zinc sorption by acidic soil amended with sewage sludge. *communication Soil Science & Plant Analysis***40(1-6)**:273-284
- Ukwattage, N.L., Ranjith, P.G. and Bouazza, M. (2013) The use of coal fly ash as a soil amendment in agricultural lands (with comments on its potential to improve food security and sequester carbon. *Fuel***109**:400-408
- Walkley, A. and Black, I. A. (1934). An examination of the Digestion method for determining soil organic matter, and a proposed modification of the chromic acid titration method. *Soil Science*. **37(1)**: 29-38.
- Wilcox, L.V. (1950) Electrical conductivity. *American water works Association Journal*; **42**:775-776.