

### **Impact of Biotic and Abiotic Stresses and their management prospects on Vegetative Growth, Fruit yield and quality of Grafted Solanaceous Vegetables for hi-valued horticultural production**

#### **ABSTRACT**

Grafted vegetable seedlings are an unparalleled horticultural technology practiced for a long time. This technique was introduced into Europe and other countries in the late 20th century along with improved grafting methods suitable for commercial production and productivity of grafted vegetable seedlings. Mostly related to disease incidences, and abiotic stress tolerance including the soil and water pollution stress. Considering various research works on grafting it can be assumed that it is one of the best methods to be included in organic farming practices. Grafting improves quality of the plant and is used to induce resistance against low and high temperatures. Growth, yield and fruit quality of the scion solanaceous crops is greatly influenced by the type of rootstock used. Commercial grafting of vegetables has started for several decades and the area under vegetable grafting is gradually increasing. This review discusses recent literature on vegetable grafting with a focus on the genetics and breeding of the rootstock, and delves into current issues affecting the grafting industry. Fruit quality of solanaceous crops plants grafted on *S. sisymbriifolium*, *S. torvum* and *S. toxicarium* (all resistant to soil-borne diseases) rootstocks was compared with that of plants on their own roots.

**Key words:** rootstock, solanaceous crops, tomato, brinjal, chilli and grafting

#### **INTRODUCTION**

In vegetable production, India comes as a second largest producer in the world after China and mostly areas covered under vegetables is 10.29 million ha and total production 176.17 million Tonnes (NHB, 2018). Grafting is a method of growing together two plant parts (a rootstock and a scion) by the process of tissue regeneration and finally grows as a one plant. Grafting of fruit trees has been practiced for many years, but in Olericulture, grafting of vegetable is a new technique. Vegetable grafting reduces the agrochemicals dependence on organic production (Rivardet *et al.*, 2008). The advantages of vegetable grafting have been noticed by many workers. Grafts were used to induce resistance against low and high temperatures, enhance nutrient uptake increase synthesis of endogenous hormones improve water use efficiency reduce uptake of persistent organic pollutants from agricultural soils improve alkalinity tolerance, raise salt tolerance and limit the negative effect of heavy metal toxicity Rivero *et al.*, (2003), Venema *et al.*, (2008), Pulgar *et al.*, (2000), Colla *et al.*, (2010a), Rouphael *et al.*, (2008), Otani and Seike (2007) Colla *et al.*, (2010), Martinez-Rodriguez *et al.*, (2008), Savvas *et al.*, (2010). The scion variety affects size, yield, and quality of fruit in grafted plants, but rootstock effects can drastically alter these quality characteristics Davis *et al.*, (2008). The quality characteristics might be affected by grafting as a result of the translocation of metabolites associated with fruit quality to the scion through the xylem and/or modification of physiological processes of the scion Rouphael *et al.*, (2008).A

report from India reveals that bacterial wilt can cause 27% losses eggplant Peddy P (1986). To causal organism resistant rootstock is an effective technology to prevent the bacterial wilt. A report from Bangladesh indicated that grafting of eggplant on resistant rootstocks to prevent bacterial wilt. Grafting tomato was introduced commercially in the 1960's (Lee and Oda, 2003). Along with the rapid development of intensive protected cultivation technologies which prevented farmers from continuing traditional crop rotation, vegetable grafting became a crucial tool to overcome soil borne diseases and other pests. In the Mediterranean, where land use is very intensive and continuous cropping is a common practice, vegetable grafting is considered an innovative technique and is in increasing demand by farmers (Khah *et al.* 2006). Vegetable grafting, primarily for tomatoes, recently increased (Atasayar 2006) and Khah *et al.* (2006) showed that tomato grafting on suitable rootstocks has positive effects on cultivation performance, especially in greenhouse conditions. Grafting of chili peppers (*C. annuum* L.) is a recent practice where *C. annuum* scions are grafted onto *C. annuum* rootstocks that have soilborne disease resistance to fungi and nematodes (Morra and Bilotto, (2006), in Rodriguez and Bosland, 2010). To promote grafttake, foliar applications of the phytohormone abscisic acid, to reduce defoliation, and the vitamin ascorbic acid, to accelerate callus formation at the cut stem surfaces, were suggested for grafting sweet peppers (*C. annuum* L.) (Johkan *et al.*, 2008, 2009). Pepper grafting may have positive (Colla *et al.*, 2008; Palada and Wu, 2008; López-Marín *et al.*, 2009; Gisbert *et al.*, 2010) or negative (M'hamdi *et al.*, 2010; GarcíaRodríguez *et al.*, 2010) influences on plant growth or yield traits.

**Challenges in grafting** Small amounts of alkaloids to the fruits from the rootstock may occur as observed when tomatoes were grafted onto jimson weed (Lowman and Kelly, 1946). The improved resistance and better yield with grafted plants was inconsistent in some cases. Proper knowledge rootstock-scion compatibility is required which may not be available and understood properly by the rural farmers.

#### **Grafting method**

Selection of grafting is done on crop with the help of the farmers experience, the number of grafts required, the motive of grafting and available facilities required for machinery and infrastructure (Lee *et al.*, 2010). Manual grafting is better and mostly practices as compared of using machines and grafting robots (Lee *et al.*, 2010).

#### **1. Tongue / approach grafting**

Equal size of rootstock and scion are used for grafting. This method requires more labor and space but percentage of seedling survival is high and most commonly practice by small nurseries farmers. This method is not used in rootstocks with hollow hypocotyls Thakur and Savita (2020).

#### **2. Cleft grafting**

It is also called apical or wedge grafting. In this method slant angle is made in lower stem and scion is pruned with 1-3 true leaves and scion is placed into split and clip is attach in between scion and rootstock (Johnson *et al.*, 2011). This is done in mainly Solanaceous crops.

#### **3. Slant grafting:**

This is also known as onecotyledon grafting. It has recently been adopted by commercial seedling nurseries. It is applicable to most vegetables. It has been developed for robotic grafting. Rootstock should be sown 7 –10 days before scion sowing to ensure uniform diameter of hypocotyls and to hold the scion on rootstock perfectly Lakshmi *et al.*, (2019).

#### **4. Hole insertion grafting:**

This is also called astop insertion grafting. This is most popular incucurbits. When scion and rootstock have hollow hypocotyls, this method is preferred. In this method grafting can be performed by making hole on the top of the root stock and by inserting the scion in that hole which should be prepared in such a way to similar diameter as with hole measurements of stock Lakshmi *et al.*, (2019).

### 5. Splice grafting

This method is mostly used by growers. It can be practice by hand or machines in most Solanaceous vegetable crops Thakur and Savita (2020). One cotyledon spliced grafting. This method is most widely used and preferred by producers and commercial graded transplant companies. It can be carried out by hand or machines in most vegetables. This method is popular for vegetable solanaceous crops Ahmed Hashim (2019).

### 6. Pin grafting

In this method designed pins are used to hold the grafted position instead of placing grafting clips. It is similar method as splice grafting Thakur and Savita (2020). The ceramic pin is nearly about 15 mm long and 0.5 mm in diagonal width of the hexagonal cross-section. The pins are made of natural ceramic so it can be left on the plant without any problem Lakshmi *et al.*, (2019). Experimental results revealed that bamboo pins, rectangular in cross-sectional shape, could successfully replace the expensive ceramic pins at much lower price.

**Table 1: Grafting methods and rootstocks used in Solanaceous crops**

Scion Plant	Rootstock	Method
Tomato	<i>Lycopersicum pimpinellifolium</i>	Cleft grafting (Ahmed Hashim (2019)
	<i>Solanum nigrum</i>	Tongue and cleft grafting (Ahmed Hashim (2019)
Brinjal	<i>Solanum torvum</i>	Tongue and cleft grafting (Ahmed Hashim (2019)
	<i>Solanum sissymbriifolium</i>	Cleft method (Ahmed Hashim (2019)
	<i>Solanum khasianum</i>	Tongue and cleft method (Ahmed Hashim (2019)

**Table.2: Potential rootstocks of solanaceous vegetable for biotic stress tolerance**

Crop	Biotic stress	Potential root stocks
Tomato	Fusarium wilts	Efialto (Bolandnazar <i>et al.</i> , 2014)
	Bacterial wilt	Dai Honmei, RST-04-105-T, VI046103 (EG195), VI045276 (EG203), Shin cheong gang F1, and Nordox (Rivard <i>et al.</i> , 2012)
	Fusarium crown and root rot	Natalia, Cuore di Bue F1 and He-Man F1 (Keatinge <i>et al.</i> , 2014)
	Southern blight	Beaufort, and Maxifort (Onduso, 2014)
	Verticillium wilt	Beaufort (Vitale <i>et al.</i> , 2014)
	Root knot nematode	Big Power, VI046103(EG195) and VI045276 (EG203) (Hibar <i>et al.</i> , 2006)

<b>Pepper</b>	Phytophthora blight	PR 920', 'PR 921', and 'PR 922 ( Kaskavalci <i>et al.</i> , 2009)
	Corky root	Eldorado, Beaufort, Snooker ( Jang <i>et al.</i> , 2012)
<b>Brinjal</b>	Bacterial Wilt	<i>Solanum toxicarium</i> , <i>S. torvum</i> and <i>S.integrifolium</i> (Rivard <i>et al.</i> , 2010)
	Nematode	Beaufort (Al- Chaabi <i>et al.</i> , 2009)

**Table 3: Potential rootstocks of solanaceous vegetable for abiotic stress tolerance**

<b>Crop</b>	<b>Abiotic stress</b>	<b>Potential root stocks</b>
<b>Egg plant</b>	High temperature tolerance	<i>Solanum integrifolium</i> × <i>Solanum melongena</i> (Lee <i>et al.</i> , (2010)
	Tolerant to drought	<i>Solanum macrocarpum</i> , <i>Solanum gilo</i> , PKM-1 (Lee <i>et al.</i> , (2010)
	Higher yield even at low temperature	<i>S. integrifolium</i> x <i>S. melongena</i> (Pandey and Rai (2003)
<b>Tomato</b>	Resistant to water-logging	<i>Solanum laciniatum</i> (Pandey and Rai (2003)
	Low and high temperature tolerance	<i>Solanum melongena</i> (Lee <i>et al.</i> , (2010)
	Resistant to drought	<i>Solanum chilense</i> (Pandey and Rai (2003)
	Tolerant to drought	<i>Solanum pennelli</i> (Penella <i>et al.</i> , 2014)
	Tolerance to cold and chilling	<i>Solanum habrochaites</i> (Penella <i>et al.</i> , 2014)
	Tolerant to humidity	<i>Solanum cerasiforme</i> (Penella <i>et al.</i> , 2014)
	Resistant to salt	<i>Solanum cheesmanii</i> (Penella <i>et al.</i> , 2014)
	Tolerance to salt	<i>Solanum galapagense</i> (Penella <i>et al.</i> , 2014)
	Flood- and drought-tolerances	<i>Solanum torvum</i> , 'EG195' o'EG203, 'PP0237-7502, PP0242-62(Penella <i>et al.</i> , 2014)
	<b>Chilli</b>	High temperature tolerant
<b>Pepper</b>	Flooding tolerance	Chili accessions 'PP0237-7502', 'PP0242-62' and 'Lee B' (Pandey and Rai (2003)
	Water stress tolerant	Atlante, C-40, Serrano, PI-152225, ECU-973, BOL-58 and NuMex Conquistador (Penella <i>et al.</i> , 2014)

### **EFFECT OF ROOTSTOCK ON GROWTH, YIELD AND QUALITY OF SOLANACEOUS VEGETABLE CROPS:**

According to Tuberosa, 2012 drought responsive trait affects the crop quality and yield when grafted under water stressed conditions. Altunlu and Gul, 2012 observed that when tomato hybrid "Beaufort" (*Solanum lycopersicum* L. × *Solanum habrochaites* S.) grafted onto rootstock ("Resistar") reduced the plant growth due to occurrence of water stress condition. Petran, 2013 reported that when tomato scions ("Celebrity")

and “3212”) onto turkey berry rootstock (*Solanum torvum* S.) under water-stressed conditions helps in delayed wilting of plant. According to Nilsen *et al.* (2014) when tomato cv. “BHN602” grafted onto the tomato rootstock “Jjak Kkung under well-water condition leads to reduction in the leaf area and growth of plant. Penella *et al.* (2014) concluded that when (*Capsicum annuum* L. cv. Verset) grafted onto the rootstocks “Atlante,” “PI-15225,” and “ECU-973 under water-stressed conditions helps in improving the marketable yield of crop by maintaining the photosynthetic rate. Ibrahim *et al.* (2014) and Al-Harbi *et al.* (2016) stated this when grown under full or deficit irrigation on tomato cv. Faridah grafted onto the tomato hybrid “Unifort” (*Solanum lycopersicum* L. × *Solanum pimpinellifolium* L.) results in the expanding of yield. López-Marín *et al.* (2017) stated when rootstock “Atlante” used as the scion acts differently when the grafting combination of Atlante/Verset and Atlante/Hermínio. Similarly, it is important to test it for drought tolerance because it influences yield and growth component. In grafted plants the movement of endogenous flowering substances across the graft union is easy. The early flowering in grafted plants may be due to better and improved root system of the rootstocks used, which has resulted in increased water and nutrient uptake. These results are in conformity with the findings of Ibrahim *et al.* (2014). The early harvest in grafted plants may be due to the compatibility of various physiological traits such as photosynthetic rate, nutrient use efficiency, proper water flow and hormonal response which also influenced plant growth and biomass production. The results are in line with the findings of Khah *et al.* (2006), Gisbert *et al.* (2010) and Ibrahim *et al.* (2014).

Higher root weight of tomato plants grafted on rootstocks are expected, since, according to Bletsos and Olympios (2008), most of the rootstocks used are breeding products with well-developed root system that result in higher biomass in scion. In pathogen-free fields, ‘He-man’ has been reported previously to have no effects on stem length in greenhouse grown tomato cv. ‘Big Red’, whereas it increased this characteristic in open field crops (Khah *et al.*, 2006). In addition, this rootstock, in another study, increased dry weight of the above-ground parts in tomato cv. ‘Rita’ (Romano and Paratore, 2001). Grafting of tomato cv. ‘Cherokee Purple’ on ‘Beaufort’ also increased growth in open field crops (Buller *et al.*, 2013). These results are in agreement with those reported by Pogonyi *et al.* (2005) who found increased productivity, expressed as a higher average fruit weight in tomato cv. ‘Lemance’ grafted on ‘Beaufort’. Also, increase of fruit size and weight as well as of total yield per plant of tomato cultivars ‘Yeni Talia’, ‘Swanson’, and ‘Beril’ grafted on ‘Beaufort’ rootstock were reported by Turhan *et al.* (2011). On the contrary Buller *et al.* (2013), reported no increase in yield of tomato cv. ‘Cherokee Purple’ grafted on ‘Beaufort’, whereas ‘He-man’ was found to increase productivity of tomato cv. ‘Big Red’ grown in greenhouse (Khah *et al.*, 2006). The strong effect of rootstock-scion combination could be the reason for these differences. The increased yield produced by plants infected with *V. dahliae* and grafted on ‘Nova’, ‘Beaufort’, and ‘Vigomax’ is in accordance with earlier reports, where the effect of rootstocks on tomato yield was independent of the disease pressure, in case of another soil-borne pathogen, such as *Pyrenochaeta lycopersici* R.W. Schneid & Gerlach (Michel and Lazzeri, 2010). In addition, yield is strongly affected by factors like pathogen identity and virulence, soil and weather conditions, and the crop cultivar used as scion (Michel and Lazzeri, 2010). Grafting on rootstocks variably affects fruit characteristics of tomato and other vegetables, which are of a major concern for consumers (Rouphael *et al.*, 2010). Among the quality characteristics affecting consumers' preferences are the total soluble solids (TSS) and pH that are connected to palatability of fruit, flesh firmness and size that define fruit appearance, and the pericarp thickness that is connected to postharvest injuries and storage. TSS is an important quality criterion reflecting concentration of sugars in fruit (Flores *et al.*,

2010). Lower TSS content recorded in tomatoes grafted on rootstocks could be attributed to cytokinins produced in rootstock, as reported in grafted watermelon by Lopez-Galarza *et al.* (2004). More specifically, these researchers provided evidence that cytokinins, after transporting to the scion, reduce sugar content. In addition, Bari and Jones (2009) reported that production of cytokinins can be increased as part of the plant's responses to the presence of pathogens (Bari and Jones, 2009). However, the slightly lower TSS content (0 Brix) and dry matter, found in fruit of tomatoes grafted on certain rootstocks, in the present study, fell within the normal range for tomato, and, therefore, are considered as not important (Flores *et al.*, 2010) and Buller *et al.* (2013) also reported that grafting on 'Beaufort' had no influence on fruit TSS of tomato cv. 'Cuore di Bue', cv. 'oxheart', and cv. 'Cherokee Purple'. Furthermore, differences were not recorded in TSS content between tomato un-grafted tomato cv. 'Big Red' and grafted on 'Heman' and 'Primavera' rootstocks grown under open-field and greenhouse conditions (Khah *et al.*, 2006). In contrast to these findings, Pogonyi *et al.* (2005) reported that soluble solid content in fruit of tomato cv. 'Lemance' was lower in plants grafted on 'Beaufort'. Turhan *et al.* (2011) also reported that grafting tomato on 'Beaufort' increased TSS content depending on cultivars used as a scion but it did not have any effect on pH.

In an experiment conducted by Khah (2005) it was reported that grafted plants have higher plant height, leaf area index, as well as the fresh and dry weights of stems and leaves and yield was comparatively higher than the nongrafted plants. Harvesting of fruits was also 7 days earlier for the grafted tomato plants. Similar observation was also recorded by Gisbert *et al.*, 2011 and Rahman *et al.*, 2002 when they grafted eggplant cultivars onto perennial and wild Solanaceous species. Augmented water and nutrient uptake improved stomatal conductance in tomato when grafted onto vigorous rootstock (Fernandez- Garcia *et al.*, 2004a). Improved photosynthesis rates of grafted plants increase the crop productivity (Matsuzoe *et al.*, 1993). According to Besri, (2003) 15-20% of yield increase was reported for the grafted tomatoes even when only half of the plants were planted. Grafting resulted in bigger fruit size in tomatoes, and thus increased the yield (Pogonyi *et al.*, 2005). Grafting is an effective approach to improve fruit quality under both optimum growth conditions and salinity. The fruit quality of shoot, at least partially, depends on the root system (Flores *et al.*, 2010). In soil less tomato cultivation, grafted plants had higher marketable yield, fruit quality (Gebologlu *et al.*, 2011). Grafting of eggplants onto *S. torvum* increased the fruit size without any effect on quality and yield. Sugar, flavor, colour, carotene content and texture can be affected by grafting and the type of rootstock used (Davis *et al.*, 2008). In contrast, grafting eggplant on *Solanum torvum* and *Solanum sisymbriifolium* negatively affected vitamin C content, firmness and some sensory attributes but overall impression was not influenced (Arvanitoyannis *et al.*, 2005). Di-Gioia *et al.*, 2010 recorded no significant differences in total soluble solids by tomato "Oxheart" grafted onto 2 inter-specific *S. lycopersicum* × *S. habrochaites* and also found that vitamin C content was decreased by 14-20 % if tomato plants grafted onto Beaufort F1 and Maxifort F1. So, there is need of further research regarding improvement of qualitative traits via grafting techniques. In domesticated tomato, variability for abiotic stresses especially salt tolerance is limited. But, wild species such as *S. pimpinellifolium*, *S. peruvianum*, *S. cheesmaniae*, *S. habrochaites*, *S. chmielewskii* and *S. pennellii* in the genus *Solanum* possessed sources of tolerance and the same have been exploited as rootstock to enhance productivity under wide environmental stress conditions, whereas, introgression with commercial tomato is difficult due to the problem of crossibility. To sum up, grafting in vegetable crops especially in solanaceous vegetables is highly useful to eliminate various challenging biotic and abiotic stressors globally. Proper selection of scion and

rootstock is the key factor for higher fruit yield and quality. Under changing climate, this technique could play significant role to improve vegetable productivity. Thus, there is a need to identify more resistant sources by exploiting wild and elite germplasm.

**Role of grafting on abiotic stress tolerance:** Environmental stress is the most important limiting factor in the current climatic situation for plant growth and horticultural productivity worldwide. Extreme temperature, draught and salinity are the major ones which have influenced crop productivity the most (Schwarz et al., 2010). Increasing global warming, expansion of the saline affected area and lack of availability quality water storage in arid lands are becoming a limiting factor for crop production and climatologists believe that the combined environmental stress in the tropics will only get worse over time. Though several stress tolerant cultivars have been produced through breeding but the complex nature of the genetic tolerance of environmental factors makes it difficult and time consuming (Ashraf and Foolad, 2007) and grafting can be a rapid alternative to develop environmental stress tolerant planting material.

## **IMPACT & SUSTENANCE ROLE OF BIOTIC STRESS TOLERANCE ON GRAFTED SOLANACEOUS VEGETABLES:**

### **Disease tolerance**

Soil borne diseases like Fusarium (*Fusarium oxysporum* f. sp. *lycopersici*, race 1 and race 2) and verticillium wilts (*Verticillium dahliae*, races 1 and races 2), bacterial speck (*Pseudomonas syringae* p.v. tomato), root knot nematodes (*Melodogyne* spp.) and corky root (*Pyrenochaeta lycopersici*) have been reported to be among the most destructive for protected tomato crops (Besri, 2002; Poffley, 2003). These used to be controlled by fumigants from which methyl bromide but after the Copenhagen Amendment to the Montreal Protocol in 1992, most of the countries around the globe phases out methyl bromide by 1 January 2005 as it was added to the list of substances that deplete the ozone layer (Anon, 2014). With rapid development in intensive protected cultivation grafting has become a crucial tool. Root stocks of certain species have excellent tolerance to various soil-borne diseases like Fusarium, Verticillium, Phytophthora, nematodes, and other pests, this kind of resistance developed in the grafted plants may be resulted from limited colonization of bacteria or pathogen in the lower stem and preventing them from invading xylem tissues (Grimault and Prior, 1994). One more reason could be the inherent resistance of the rootstocks and improved plant nutrient uptake which improves disease control in grafted vegetable plants. Few experiments have also reported to provide some protection from viruses when the susceptible scions were grafted on specific root socks. Mahmoud, (2014) found that grafting increased TYLCV tolerance in susceptible plants, delayed the appearance of TYLCV symptoms and increase of yield components compared to non-grafted plants in tomato. Iouannou (2001) observed grafted eggplants showed lower disease incidences with higher yield and fruit size. Similar type of observations were obtained when tomato was grafted onto Beaufort' rootstocks (*S. lycopersicum* and *S. habrochaites*) (Hasna et al., 2009). Tomatoes (*Solanum lycopersicum* L.) when grafted onto commercial rootstock and subjected to infection with *Verticillium dahliae*, the grafted plant exhibited resistance to the verticillium wilt disease (Paplomatas et al., 2002). Grafting of chilli peppers (*Capsicum annum* L.) is a recent practice where *C. annum* scions are grafted onto *C. annum* rootstocks that have soilborne disease and nematode resistance (Morra and Biloto 2006). Rivard and Louws (2008) found that 'German Johnson' heirloom tomatoes had 0% fusarium wilt incidence in

infested soils when grafted onto resistant CRA 66 or Hawaii 7996 tomato rootstock, compared to a 79% incidence on non-grafted controls.

### **CONCLUSION:**

Vegetable grafting is also developing around the globe as a means to overcome a-biotic and biotic stresses which are responsible for 70% and 30% of the yield gap, respectively. Among the major a-biotic stresses, salinity and drought are the ones forecasted to rise due to global climate change. Not with standing, the enormous significance of grafting as a means for securing yield stability and quality in vegetables crops, commercial its practice has heavily centered on the production of high-valued solanaceous and cucurbitaceous vegetable crops. Among future perspectives is the extension of grafting practice to other seasonal crops including combinations where both rootstock and scion deliver harvestable products. The various innovative pilot research studies nevertheless demonstrated the potential of creating harvestable rootstock–scion combinations as a means of saving growth space and minimizing waste. Such unique grafting model systems may assist in elucidating scion–rootstock synergy and sink competition in production of high-valued horticultural crops.

### **REFERENCES:**

- Ahmed Hashim., A.A.R., 2019. Grafting Techniques in Vegetables Crops: A Review. *Plant Archives* Vol. 19 No. 1, 2019 pp. 49-51
- Al- Chaabi, S., Koutifani, O., Safeih, M.H. Asmar, J., 2009. Management of Root-Knot Nematodes and Corky Root Disease of Pepper Plants by Grafting Technique onto Resistant Rootstocks under Plastic House. February 2009.
- Al-Harbi, A., Hejazi, A., Al-Orman, A., Responses of grafted tomato (*Solanum lycopersicum* L.) to abiotic stresses in Saudi Arabia. *Saudi Journal Biological Sciences*.26(6):1274-1280.
- Ali,M., Alam ,MZ., Akanda ,MAM., 1994. Graiing: A technique of control soil-borne diseases of tomato and eggplant. *Studies in Agriculture*.
- Altunlu ,H., Gul ,A., 2012 Increases drought tolerance of tomato plants by grafting. *Acta Horticulture* 960:183-190.
- Anonymous 2014. Introduction: Methyl Bromide and the montreal protocol. Phasing-out Methyl Bromide in Developing Countries. Pp: 11-14.
- Arvanitoyannis Ioannis, Khah Ebrahim M, Christakou Evangelia C. and Bletsos Fotios A. (2005). Effect of grafting and modified atmosphere packaging on eggplant quality parameters during storage. *International Journal of Food Science & Technology* 40(3):311 – 322.
- Atasayar ,A., 2006. The usage of grafted watermelon seedling in Turkey. *Hasad Horticulture Magazine*, 252: 87–91.
- AVRDC 2003. Guide. Grafting tomatoes for the production in the hot-wet season. Asian Vegetable Research and Development Center, Pub. No 03-551, Shanhua, Tainan, Taiwan, 6pp.
- AVRDC 2009. Guide. Grafting sweet peppers for the production in the hotwet season. Asian Vegetable Research and Development Center, Pub. No 09- 722, Shanhua, Tainan, Taiwan, 8pp.

- Bari, R., Jones, D.G., 2009. Role of plant hormones in plant defence responses. *Plant Molecular Biology*. 69, 473e488.
- Besri, M. 2002. Tomato grafting as an alternative to methyl bromide in Morocco. Institute Agronomique et Veterinaire Hasan II. Morocco.
- Besri, M., 2002. Tomato grafting as an alternative to methyl bromide in Morocco. Institute Agronomique et Veterinaire Hasan II. Morocco
- Bletsos Fotios A. and Olympos Christos (2008). The European journal of Plant Science and Biotechnology 2 (1) 62-73.
- Bletsos, F.A., 2005. Use of grafting and calcium cyanamide as alternatives to methyl bromide soil fumigation and their effects on growth, yield, quality and fusarium wilt control in melon. *J. Phytopathol.* 153, 155–161
- Bolandnazar, S., Moghbeli, E.M., Panahandeh, J. and Arzanlou, M. 2014. Biological control of fusarium wilt in greenhouse tomato by Mycorrhizal fungi and resistant rootstock. *Acta Horticulturae*. 1041:127-132.
- Buller, S., Inglis, D., Miles, C., 2013. Plant growth, fruit yield and quality, and tolerance to Verticillium wilt of grafted watermelon and tomato in field production in the Pacific Northwest. *Horticulture Science* 48.
- Buller, S., Inglis, D. & Miles, C. 2013 Plant growth, fruit yield and quality, and tolerance to verticillium wilt of grafted watermelon and tomato in field production in the Pacific Northwest *HortScience* 48 1003 1009
- Colla, G., Suarez, CMC., Cardarelli, M., Roupheal, Y., 2010 Improving nitrogen use efficiency in melon by grafting. *Horticulture Science* 45: 559-565.
- Colla, G., Roupheal Y, Cardarelli M, O, Temperini, E, Rea, Salern., A, Pierandrei F, 2008. Influence of grafting on yield and fruit quality of pepper (*Capsicum annuum* L.) grown under greenhouse conditions. *Acta Horticulturae* (ISHS) 782, 359–364.
- Colla, G., Roupheal, Y., Cardarelli, M., Salerno, A., Rea, E., 2010a The effectiveness of grafting to improve alkalinity tolerance in watermelon. *Environment Experiment Botany* 68: 283-291.
- Davis, AR., Perkins-Veazie, P., Hassell, R., Levi, A., King, SR., Zhang, X., 2008. Grafting Effects on Vegetable Quality. *Horticulture Science* 43(6): 1670-1672.
- Di-Gioia F, Serio F, Buttaro D, Ayala O, Santamaria P. Influence of rootstock on vegetative growth, fruit yield and quality in 'Cuore di Bue', an heirloom tomato *Journal of Horticultural Sciences and Biotechnology* 2010;85(6):477-482.
- Fernández-García, N., Carvajal M., and Olmos. E. 2004. graft union formation in tomato plants: peroxidase and catalase involvement. *Annals of Botany* 93(1):53-60.
- Fernandez-Garcia, N., Martinez, V., Cerda, A., and Carvajal, M. 2004b. Fruit quality of grafted tomato plants grown under saline conditions. *Journal of Horticultural Science and Biotechnology*. 79: 995–1001.
- Flores F.B., Sanchez-Bel P., Estan M.T., Martinez-Rodriguez M.M., Moyano E., Morales B., Campos J.F., Garcia-Abellan J.O., Egea M.I., Fernandez-Garcia N., Romojaro F., Bolarín M.C., 2010. The effectiveness of grafting to improve tomato fruit quality. *Scientia Horticulturae* 125: 211-217.

- García-Rodríguez M.R., Chiquito-Almanz., E, LoezaLara., P.D, Godoy-Hernández., H, Villordo-Pineda., E, PonsHernández., J.L, González-Chavira., M.M, AnayaLópez., J.L, 2010. Production of ancho chili graft on criollo de morelos 334 for the control of *Phytophthora capsici*. *Agrociencia* 44, 701–709.
- Gebologlu N, Yilmaz E, Cakmak P, Aydin M and Kasap Y 2011. Determining of the yield, quality and nutrient content of tomatoes grafted on different rootstocks in soilless culture. *Scientific Research and Essays* 6(10): 2147- 2153.
- Gisbert C., Sánchez-Torres, P. Raigón M.D. and F. Nuez 2010. *Phytophthora capsici* resistance evaluation in pepper hybrids: Agronomic performance and fruit quality of pepper grafted plants. *Journal of Food, Agriculture and Environment* 8, 116–121.
- Gisbert, C., Prohens, J., Raigón, M.D., Stommel, J.R. and Nuez, F. 2011. Eggplant relatives as sources of variation for developing new rootstocks: Effects of grafting on eggplant yield and fruit apparent quality and composition. *Scientific Horticulturae*.128(1): 14-22.
- Grimault, V. and Prior, P. 1994. Grafting tomato cultivars resistant or susceptible to bacterial wilt: Analysis of resistance mechanisms. *Journal of Phytopathology*. 141: 330–334.
- Hasna, M.K., Ogren, E., Persson, P., Ma´rtensson, A. and Ramert, B. 2009. Management of corky root disease of tomato in participation with organic tomato growers. *Crop Protection*. 28: 155– 161.
- Ibrahim, A., Wahb-Allah,M., Abdel-Razzak ,H., Alsadon, A., 2014. Growth, yield, quality and water use efficiency of grafted tomato plants grown in greenhouse under different irrigation levels. *Life Science Journal* 11:118-126
- Jang, Y., Yang, E., Cho, M., Um, Y., Ko, K., Chun, C. 2012. Effect of grafting on growth and incidence of phytophthora blight and bacterial wilt of pepper (*Capsicum annum* L.). *Horticulture, Environment, and Biotechnology*. 53(1): 9–19
- Johkan Masahumi, Mitukuri Kazuhiko, Yamasaki Satoshi and Mori Genjirou (2008a). Causes of defoliation and low survival rate of grafted sweet pepper plants. *Scientia Horticulturae* 119(2):103-107
- Johkan M., K. Mitukuri, S. Yamasaki, G. Mori and M. Oda, (2009b). Causes of defoliation and low survival rate of graft -ed sweet pepper plants. *Scientia Horticulturae* 119, 103–107.
- Johnson, S., Kreider, P., Miles C. 2011. Vegetable Grafting Eggplants and Tomatoes. Washington State University 4.
- Keatinge, J.D.H., Lin, L.-J., Ebert, A.W., Chen, W.Y., Hughes, J.d'A., Luther, G.C., Wang, J.-F. Ravishanka, M., 2014. Overcoming biotic and abiotic stresses in the Solanaceae through grafting: current status and future perspectives. *Biological Agriculture & Horticulture*. 30(4): 272-287.
- Khah E.M., Kakava E., Mavromatis A., Chachalis D., Goulas C., 2006. Effect of grafting on growth and yield of tomato (*Lycopersicon esculentum* Mill.) in greenhouse and open-field. *Journal of Applied Horticulture*, 8: 3–7.
- Khah, E. M. 2005. Effect of grafting on growth, performance and yield of aubergine (*Solanum melongena* L.) in the field and greenhouse. *Journal of Food, Agriculture & Environment*. 3(3&4) : 92 - 94.
- Lakshmi, T., Tirupathamma., Venkata,C.,Ramana, Naram ., L, Naidu ., Sasikala ,K., 2019. Vegetable Grafting: A Multiple Crop Improvement Methodology. *Current Journal of Applied Science and Technology*. 33(3): 1-10.

- Lee JM, Oda M (2003) Graing of herbaceous vegetable and ornamental crops. Hort Rev 28: 61-124.
- Lee, Jung-Myung., Kubota, C., Tsao, S.J., Bie, Z., Echevarria, H., Morra, P.L. and Oda, M.2010. Current status of vegetable grafting: Diffusion, grafting techniques, automation. Scientia Horticulturae. 127(2,8): 93-105.
- Lopez-Galarza S., San Bautista A., Perez D. M. and Miguel A. (2004). Effects of grafting and cytokinin – Induced fruit setting on colour and sugar – Content traits in glasshouse – Grown triploid watermelon. Journal of Horticultural Science and Biotechnology 79(6):971-976
- López-Marín J., A. Gálvez, A. González and J.A. Fernández, 2009. Agronomic behaviour of grafted sweet pepper grown in a greenhouse in Mediterranean area. Acta Horticulturae (ISHS) 807, 655–660.
- López-Marín, J., Gálvez, A., Del Amol, FM., Albacete, A., Fernandez, JA., Egea-Gilabert, C., 2017. Selecting vegetative/ generative/dwarfing rootstocks for improving fruit yield and quality in water stressed sweet peppers. Science Horticulture; 214:9-17.
- M'hamdi M., N. Boughalleb, N. Ouhaibi, N. Tarchoun, M. Souli and L. Belbahri, 2010. Evaluation of grafting techniques and a new rootstock for resistance of pepper (*Capsicum annum* L.) towards *Phytophthora nicotianae*. Journal of Food, Agriculture and Environment 8, 135–139.
- Mahmoud, M.A. 2014. Grafting as a Tool to Improve TYLCV-Tolerance in Tomato. Journal of Horticultural Science & Ornamental Plants. 6 (3): 109-115.
- Martinez-Rodriguez, MM., Estan, MT., Moyano, E., Garcia-Abellan ,JO., Flores ,FB., 2008 The effectiveness of grafting to improve salt tolerance in tomato when an 'excluder' genotype is used as scion. Environ Exp Bot 63: 392-401.
- Matsuzoe, N., Nakamura, H., Okubo, H., and Fujieda, K. 1993. Growth and yield of tomato plants grafted on *Solanum* rootstocks. Journal of the Japanese Society for Horticultural Science. 61:847-855.
- Michel, V.V., Lazzeri, L., 2010. Green manures and organic amendments to control corky rot of tomato. Acta Hort. 883, 287e294.
- Morra ,L., M, Bilotto .,2006. Evaluation of new rootstocks for resistance to soilborne pathogens and productive behavior of pepper (*Capsicum annum* L.) Journal of Horticultural Science and Biotechnology 81, 518–524.
- Morra, L. and Bilotto, M. 2006 Evaluation of new rootstocks for resistance to soilborne pathogens and productive behavior of pepper (*Capsicum annum* L.) Journal of Horticultural Science and Biotechnology. 81:518-524.
- Nilsen, ET., Freeman, J., Grene ,R., Tokuhisa, J., A rootstock provides water conservation for a grafted commercial tomato (*Solanum lycopersicum* L.) line in response to mild-drought conditions: a focus on vegetative growth and photosynthetic parameters. Plos one 2014;9(12): e115380.
- Onduso, J. N. 2014. Management of bacterial wilt of tomato by use of resistant rootstock. Masters of science Thesis. University of Nairobi, Kenya.
- Otani, T., Seike, N., 2007 Rootstock control of fruit dieldrin concentration in grafted cucumber (*Cucumissativus*). J Pestic Sci 32: 235-242.

- Palada, M.C., D.L. Wu., 2008. Evaluation of chili rootstocks for grafted sweet pepper production during the hot-wet and hot-dry seasons in Taiwan. *Acta Horticulturae (ISHS)* 767, 151–158. [http://www.actahort.org/books/767/767\\_14.htm](http://www.actahort.org/books/767/767_14.htm)
- Pandey., A.K., Rai, M., 2003. Prospects of grafting in vegetables: an appraisal. *Vegetable science*. 30(2): 101-109.
- Paplomatas, E.J., Elena, K., Tsagkarakou, A. and Perdikaris, A.2002. Control of Verticillium wilt of tomato and cucurbits through grafting of commercial varieties on resistant rootstocks. *Acta Horticulturae*. 579:445–449.
- Peddy, P.,1986 Analysis of crop losses in certain vegetables due to nematode. *Int Nematol Net New let* 3: 3-5. 15.
- Penella C, Nebauer SG, San Bautista A, López-Galarza S, Calatayud A. 2014b Rootstock alleviates PEG-induced water stress in grafted pepper seedlings: physiological responses. *Journal of Plant Physiology* 171:842-851.
- Petran, AJ., Interspecific Grafting of Tomato (*Solanum lycopersicum*) Onto Wild Eggplant (*Solanum torvum*) for Increased Environmental Tolerances. M.Sc. Thesis University of Minnesota, Minneapolis, MN 2013.
- Poffley, M. 2003. Grafting tomatoes for bacterial wilt control. *Agnote*, 603, No. B40.
- Pogonyi, A., Peka, Z., Helyes, L., Lugasi, A., 2005. Effect of grafting on the tomato's yield, quality and main fruit components in spring forcing. *Acta Aliment.* 34, 453-462.
- Pulgar, G., Villora, G., Moreno, DA., Romero L (2000) Improving the mineral nutrition in grafted watermelon plants: nitrogen metabolism. *Biol Plant* 43: 607-609.
- Rahman, M.A., Rashid, M.A., Salam, M.A., Masud, M.A.T., Masum, A.S.M.H., and Hossain, M.M. 2002. Performance of some grafted eggplant genotypes on wild *Solanum* root stocks against rootknot nematode. *Journal of Biological Sciences* 2:446-448.
- Rivard, C. L., O'Connell, S., Peet, M M., and Louws, F. J. 2010. Grafting Tomato with Interspecific Rootstock to Manage Diseases Caused by *Sclerotium rolfsii* and Southern Root-Knot Nematode. *Plant Disease*. 94(8): 1015- 1021.
- Rivard, C.S. and F.J. Louws. 2008. Grafting to Manage Soilborne Disease in Heirloom Tomato Production. *HortScience*. 43(7): 2104-2111.
- Rivero, RM., Ruiz, JM., Romero, L., 2003 Role of grafting in horticultural plants under stress conditions. *J Food Agric Environ* 1: 70-74.
- Romano Daniela and A. Paratore (2001). Effects of grafting on tomato and eggplant. *Acta Horticulturae* 559(559):149-154
- Rouphael Youssef, Schwarz Dietmar, Krumbain Angelika and Colla Giuseppe (2010). Impact of grafting on product quality of fruit vegetables. *Scientia Horticulturae* 127(2):172-179
- Rouphael ,Y., Cardarelli ,M., Colla, G., Rea, E., 2008 Yield, mineral composition, water relations, and water use efficiency of grafted mini watermelon plants under deficit irrigation. *Hort Sci* 43: 730-736.

- Sanchez-Rodriguez, E., Ruiz, J. M., Ferreres, F. and Moreno, D.A. 2012. Phenolic Metabolism in Grafted versus Nongrafted Cherry Tomatoes under the Influence of Water Stress. *Journal of Agricultural and Food Chemistry*. 59(16):8839-8846.
- Savvas, D., Colla, G., Roupael, Y., Schwarz, D., 2010 Amelioration of heavy metal and nutrient stress in fruit vegetables by grafting. *Sci Hort* 127: 156-161.
- Thakur Diksha., Savita 2020. Role of grafting in vegetable crops: A review. *Journal of Pharmacognosy and Phytochemistry*, 9(6): 1170-1174
- Tuberosa, R., Phenotyping for drought tolerance of crops in the genomics era. *Front. Physiol* 2012; 3:347.
- Turhan, A., Ozmen, N., Serbeci, M.S., Seniz, V., 2011. Effects of grafting on different rootstocks on tomato fruit yield and quality. *Hort. Sci.* 38, 142e149.
- Turhan, A., Ozmen, N., Serbeci, M.S. & Seniz, V. 2011 Effects of grafting on different rootstocks on tomato fruits yield and quality *Hort. Sci. (Prague)* 38 142 149
- Venema, JH., Dijk, BE., Bax, JM., Hasselt, P., Elzenga, JTM., 2008 Grafting tomato (*Solanum lycopersicum*) onto the rootstock of a high-altitude accession of *Solanum habrochaites* improves sub-optimal temperature tolerance. *Environ Exp Bot* 63: 359-367.
- Vitale, A., Rocco, M., Arena, S., Giuffrida, F., Cassaniti, C., Scaloni, A., Lomaglio, T., Guarnaccia, V., Polizzi, G., Marra, M. and Leonardi, C. 2014. Tomato susceptibility to fusarium crown and root rot: effect of grafting combination and proteomic analysis of tolerance expression in the rootstock. *Plant Physiology and Biochemistry*. 83:207- 216.