

### **Impact of Biotic and Abiotic Stresses and their management prospects on Vegetative Growth, Fruit yield and quality of Grafted Solanaceous Vegetables for hi-valued horticultural production**

#### **ABSTRACT**

Grafted veggie seedlings are an exceptional horticultural technique that has been used for a very long time. This approach, along with improved grafting techniques appropriate for commercial production and productivity of grafted vegetable seedlings, was introduced into Europe and other nations in the late 20th century. mainly connected to illness prevalence and abiotic stress tolerance, such as stress from soil and water pollution. Grafting is one of the best techniques to use in organic farming practises, according to a number of study studies on the subject. In order to increase a plant's quality and resilience to both high and low temperatures, grafting is used. The variety of rootstock used has a significant impact on the growth, yield, and fruit quality of scion solanaceous crops. With a focus on the genetics and breeding of the rootstock, this review examines recent literature on vegetable grafting and explores current problems affecting the grafting industry. Fruit quality of solanaceous agricultural plants grafted on *S. sisymbriifolium*, *S. torvum*, and *S. toxicarium* rootstocks—all of which are immune to soil-borne diseases—was compared to that of plants on their own roots.

**Key words:** rootstock, solanaceous crops, tomato, brinjal, chilli and grafting

#### **INTRODUCTION**

"Grafting is a technique for combining two plant components (a rootstock and a scion) through the process of tissue regeneration, which eventually results in the growth of a single plant. Grafting of vegetables is a novel method in olericulture, a field of horticulture that deals with the cultivation, preservation, processing, and marketing of vegetables. Grafting of fruit trees has been done for many years. Vegetable grafting lessens the reliance on agrochemicals in organic agriculture" (Rivardet *et al.*, 2008). Many employees have become aware of the benefits of vegetable grafting. Grafts were used to "improve nutrient uptake, boost endogenous hormone synthesis, and promote resistance against low and high temperatures." increase water usage effectiveness decrease the uptake of persistent organic contaminants from farming soils, increase salt and alkalinity tolerance, and lessen the toxic effects of heavy metals Rivero *et al.*, (2003), Venema *et al.*, (2008), Pulgar *et al.*, (2000), Colla *et al.*, (2010a), Rouphe *et al.*, (2008), Otani and Seike (2007) Colla *et al.*, (2010), Martinez-Rodriguez *et al.*, (2008), Savvas *et al.*, (2010). Size, yield, and fruit quality are influenced by scion variety in grafted plants, but rootstock impacts have a significant impact on these quality traits as well. Davis *et al.*, (2008). Grafting may alter the quality traits by transferring metabolites linked to fruit quality to the scion through the xylem and/or altering the physiological functions of the scion. Rouphe *et al.*, (2008). By promoting improved nutrient uptake and supplying essential resistance against pests and diseases, rootstocks play a significant part

in increasing fruit output. The genetic diversity of rootstocks led to interactions with the soil's quality, replanting circumstances, and pH that resulted in varying nutrient concentrations in the scions' fruits and foliage, which allowed for the observation of the differences between rootstocks. Dowarahet *al.*, (2022). According to a study from India, bacterial wilt can result in a 27% loss of eggplant. P. Peddy (1986). An efficient technique to stop the bacterial wilt is to use rootstock that is resistant to the causative organism. According to a study from Bangladesh, grafting eggplant on rootstocks that are resistant to bacterial wilt can cause the disease. In the 1960s, grafting tomatoes were made available for purchase. (Lee and Oda, 2003). Vegetable grafting evolved into a critical tool to combat soil-borne diseases and other pests along with the quick development of intensive protected cultivation technologies that prohibited farmers from continuing traditional crop rotation. Vegetable grafting is regarded as an innovative method and is in growing demand by farmers in the Mediterranean, where continuous cropping is a prevalent practice and land use is very intensive. (Khahet *al.* 2006). Tomato grafting on suitable rootstocks has been shown to improve cultivation performance, particularly in greenhouse conditions, according to Khahet *et al.* (2006). Vegetable grafting, mainly for tomatoes, has recently increased (Atasayar 2006). Recently, chilli peppers (*C. annum L.*) have been grafted onto rootstocks that have resilience to nematodes and fungi that cause soilborne diseases (Morra and Bilotto, 2006, in Rodriguez and Bosland, 2010). For sweet peppers (*C. annum L.*) grafting, foliar applications of the phytohormone abscisic acid, to lessen defoliation, and the vitamin ascorbic acid, to hasten callus development at the cut stem surfaces, were recommended (Johkanet *al.*, 2008, 2009). Colla *et al.* (2008), Palada and Wu (2008), López-Marn *et al.* (2009), Gisbert *et al.* (2010), and others have shown that pepper grafting can affect plant growth or yield traits positively or negatively (M'hamdi *et al.* (2010) and Garcá-Rodríguez *et al.* (2010)).

**Challenges in grafting** As was seen when tomatoes were grafted onto jimson weed, a small quantity of alkaloids from the rootstock may transfer to the fruits (Lowman and Kelly, 1946). There were varying degrees of improvement in resistance and production with grafted plants. Rootstock-scion compatibility requires knowledge that may not be accessible to or correctly understood by rural farmers.

### **Grafting method**

Crops are chosen for grafting with the aid of the farmer's experience, the number of grafts needed, the purpose of the graft, and the availability of the facilities needed for the machinery and infrastructure (Lee *et al.*, 2010). As opposed to using machinery or grafting robots, manual grafting is preferable and more common (Lee *et al.*, 2010).

#### **1. Tongue / approach grafting**

For grafting, equal-sized stem and scion are used. Small-scale farmers most frequently use this technique, which requires more work and room but has a high percentage of seedling survival. In rootstocks with empty hypocotyls, this technique is not applied. both Thakur and Savita (2020).

#### **2. Cleft grafting**

Another name for it is wedge or apical anastomosis. This technique involves making a slant angle in the lower stalk, pruning the scion with 1-3 true leaves, splitting the scion, and attaching a clip between the scion and the rootstock (Johnson *et al.*, 2011). Primarily solanaceous products are used for this.

#### **3. Slant grafting:**

One cotyledon anastomosis is another name for this. Commercial seedling nurseries have just lately started using it. It applies to the majority of veggies. It was made specifically for autonomous grafting. To guarantee

uniform hypocotyl diameter and securely retain the scion on the rootstock, rootstock should be sown 7–10 days before scion sowing. Lakshmi and co. (2019).

#### 4.Hole insertion grafting:

Another name for this is top insertion grafting. This is the most well-liked in cucurbit. This approach is favored when both the rootstock and the scion have hollow hypocotyls. According to this technique, grafting can be done by creating a hole on top of the root stock and inserting the scion into that hole, which should be prepared to have a comparable diameter to the measurements of the stock hole (Lakshmi et al., 2016).

#### 5.Splice grafting

The majority of producers use this technique. In the majority of Solanaceous Vegetable Crops, it can be done manually or mechanically Savita and Thakur (2020). One cotyledon was grafted together. The majority of producers and commercial graded transplant businesses use and favour this technique. In most veggies, it can be done manually or mechanically. This approach is common for solanaceous vegetable products Hashim Ahmed (2019).

#### 6.Pin grafting

Instead of using grafting clips, specially designed pins are used in this technique to hold the grafted position. It is a technique akin to the patch grafting Thakur and Savita used (2020). The ceramic pin has a hexagonal cross-section with a diagonal diameter of 0.5 mm and is almost 15 mm long. Because the pins are natural ceramic, they can be kept on the plant without any issues. Lakshmi and co. (2019). Experimental findings showed that bamboo pins with a rectangular cross-section could effectively substitute costly ceramic pins for a much lower cost.

**Table 1: Grafting methods and rootstocks used in Solanaceous crops**

Scion Plant	Rootstock	Method
Tomato	<i>Lycopersicon pimpinellifolium</i>	Cleft grafting (Ahmed Hashim (2019)
	<i>Solanum nigrum</i>	Tongue and cleft grafting (Ahmed Hashim (2019)
Brinjal	<i>Solanum torvum</i>	Tongue and cleft grafting (Ahmed Hashim (2019)
	<i>Solanum sissymbriifolium</i>	Cleft method (Ahmed Hashim (2019)
	<i>Solanum khasianum</i>	Tongue and cleft method (Ahmed Hashim (2019)

**Table.2: Potential rootstocks of solanaceous vegetable for biotic stress tolerance**

Crop	Biotic stress	Potential root stocks
Tomato	Fusarium wilts	Efialto (Bolandnazaret <i>al.</i> ,2014)
	Bacterial wilt	Dai Honmei, RST-04-105-T, VI046103 (EG195), VI045276 (EG203), Shin cheong gang F1, and Nordox (Rivardet <i>al.</i> , 2012)
	Fusarium crown and root rot	Natalia, Cuore di Bue F1 and He-Man F1(Keatingeet <i>al.</i> , 2014)
	Southern blight	Beaufort, and Maxifort (Onduso, 2014)

	Verticillium wilt	Beaufort (Vitale <i>et al.</i> , 2014)
	Root knot nematode	Big Power, VI046103(EG195) and VI045276 (EG203) (Hibaret <i>et al.</i> , 2006)
<b>Pepper</b>	Phytophthora blight	PR 920', 'PR 921', and 'PR 922 ( Kaskavalci <i>et al.</i> , 2009)
	Corky root	Eldorado, Beaufort, Snooker ( Jang <i>et al.</i> , 2012)
<b>Brinjal</b>	Bacterial Wilt	<i>Solanum toxicarium</i> , <i>S. torvum</i> and <i>S.integrifolium</i> (Rivardet <i>et al.</i> , 2010)
	Nematode	Beaufort (Al- Chaabi <i>et al.</i> , 2009)

**Table 3: Potential rootstocks of solanaceous vegetable for abiotic stress tolerance**

<b>Crop</b>	<b>Abiotic stress</b>	<b>Potential root stocks</b>
<b>Egg plant</b>	High temperature tolerance	<i>Solanum integrifolium</i> x <i>Solanum melongena</i> (Lee <i>et al.</i> , (2010)
	Tolerant to drought	<i>Solanum macrocarpum</i> , <i>Solanum gilo</i> , PKM-1 (Lee <i>et al.</i> , (2010)
	Higher yield even at low temperature	<i>S. integrifolium</i> x <i>S. melongena</i> (Pandey and Rai (2003)
<b>Tomato</b>	Resistant to water-logging	<i>Solanum laciniatum</i> (Pandey and Rai (2003)
	Low and high temperature tolerance	<i>Solanum melongena</i> (Lee <i>et al.</i> , (2010)
	Resistant to drought	<i>Solanum chilense</i> (Pandey and Rai (2003)
	Tolerant to drought	<i>Solanum pennellii</i> (Penella <i>et al.</i> , 2014)
	Tolerance to cold and chilling	<i>Solanum habrochaites</i> (Penella <i>et al.</i> , 2014)
	Tolerant to humidity	<i>Solanum cerasiforme</i> (Penella <i>et al.</i> , 2014)
	Resistant to salt	<i>Solanum cheesmanii</i> (Penella <i>et al.</i> , 2014)
	Tolerance to salt	<i>Solanum galapagense</i> (Penella <i>et al.</i> , 2014)
	Flood- and drought-tolerances	<i>Solanum torvum</i> , 'EG195' o'EG203, 'PP0237-7502, PP0242-62(Penella <i>et al.</i> , 2014)
<b>Chilli</b>	High temperature tolerant	<i>C. annum</i> cv. Toom-1 and 9852-54 (AVRDC 2003 & 2009)
<b>Pepper</b>	Flooding tolerance	Chilli accessions 'PP0237-7502', 'PP0242-62' and 'Lee B' (Pandey and Rai (2003)
	Water stress tolerant	Atlante, C-40, Serrano, PI-152225, ECU-973, BOL-58 and NuMex Conquistador (Penella <i>et al.</i> , 2014)

## **EFFECT OF ROOTSTOCK ON GROWTH, YIELD AND QUALITY OF SOLANACEOUS VEGETABLE CROPS:**

### **GROWTH:**

When grafted under water-stressed circumstances, the 2012 drought responsive trait, according to Tuberosa, affects the crop quality and yield. The tomato hybrid "Beaufort" (*Solanum lycopersicum* L. *Solanum habrochaites* S.) grafted onto rootstock ("Resistar"), according to Altunlu and Gul (2012), inhibited plant development when water stress conditions were present. According to Petran (2013), tomato scions ("Celebrity" and "3212") planted on turkey berry rootstock (*Solanum torvum* S.) in drought-stressed circumstances help delay plant wilting. Nilsen et al. (2014) found that when tomato cv. "BHN602" was grafted onto the tomato rootstock "Jakkung" under well-water conditions, the plant's development and leaf area were reduced. Open field harvests grew more quickly after being grafted with the tomato variety "Cherokee Purple" on "Beaufort" (Buller et al., 2013). These outcomes are consistent with those observed by Pogonyiet al. (2005) who discovered increased productivity in tomato cv. "Lemance" grafted on "Beaufort," as indicated by a higher average fruit weight. According to an experiment done by Khah (2005), grafted plants have greater plant heights, leaf area indices, fresh and dry weights of stems and leaves, and yields that are comparable to those of nongrafted plants.

### **YIELD:**

According to Penella et al. (2014), the rootstocks "Atlante," "PI-15225," and "ECU-973" when grafted onto (*Capsicum annum* L. cv. Verset) under water-stressed conditions aid in improving the marketable yield of crop by keeping the photosynthetic rate. This was mentioned by Ibrahim et al. (2014) and Al-Harbi et al. (2016) for the reigntomato cv when grown under full or deficit irrigation. The output increases when Faridah is grafted onto the tomato hybrid "Unifort" (*Solanum lycopersicum* L. *Solanum pimpinellifolium* L.). According to Lopez-Marn et al. (2017), the stem "Atlante" used as the scion behaves differently depending on whether it is grafted onto Atlante/Verset or Atlante/Herminio. Similarly, since it affects yield and growth components, it is crucial to evaluate it for drought tolerance. Endogenous flowering compounds easily penetrate the graft union in grafted plants.

On the other hand, no increase in tomato cv yield was recorded by Buller et al. in 2013. The yield of the greenhouse-grown tomato variety "Big Red" was found to be increased by "He-man," but not by "Cherokee Purple," which was grafted on "Beaufort" (Khahet al., 2006). These variations may be brought about by the potent effects of rootstock-scion combinations. The higher yield produced by plants grafted on "Nova," "Beaufort," and "Vigomax" and infected with *V. dahliae* is consistent with earlier reports that found the influence of rootstocks on tomato yield to be independent of disease pressure in the case of another soil-borne pathogen, such as *Pyrenochaeta lycopersici* R.W. Schneid & Gerlach (Michel and Lazzeri, 2010). Additionally, elements like pathogen identity and virulence, soil and weather conditions, and the crop cultivar used as a scion all have a significant impact on output (Michel and Lazzeri, 2010). Even when only half of the seedlings were planted, Besri (2003) reports that the grafted tomatoes' yield increased by 15-20%. Grafting improved tomato yield by resulting in larger fruit size (Pogonyiet al., 2005). Under ideal growth circumstances and salinity, grafting is a successful method to enhance fruit quality. The root structure, at least in part, determines the fruit quality of the

shoot (Flores et al., 2010). Grafted plants had better marketable yield and fruit quality in soil-free tomato farming (Gebologluet al., 2011).

#### **QUALITY:**

The root systems of the used rootstocks, which have led to greater water and nutrient uptake, may be the cause of the early flowering in grafted plants. These outcomes are consistent with the conclusions reached by Ibrahim et al (2014). The compatibility of various physiological traits, such as photosynthetic rate, nutrient use efficiency, appropriate water flow, and hormonal response, which also affected plant growth and biomass production, may be the cause of the early harvest in grafted plants. The results are in line with the findings of Khahet *al.* (2006), Gisbertet *al.* (2010) and Ibrahim *et al.* (2014). According to Bletsos and Olympios (2008), most of the rootstocks used are breeding products with well-developed root systems that result in higher biomass in scion, so higher root weight of tomato plants grafted on rootstocks is anticipated. He-man was earlier reported to have no effects on stem length in greenhouse-grown tomato cv. "Big Red" in pathogen-free fields, but to increase this characteristic in open field crops (Khahet al., 2006). Additionally, in another research, this rootstock increased the dry weight of the above-ground parts in tomato cv (Romano and Paratore, 2001). Additionally, 'Yeni Talia,' 'Swanson,' and 'Beril' tomato cultivars grafted on 'Beaufort' rootstock reported increases in fruit size and weight as well as overall yield per plant. Turhanet *al.* (2011). Consumers are very concerned about how tomato and other vegetable fruit traits are affected by grafting on rootstocks, which can vary (Rouphalet al., 2010). Total soluble solids (TSS) and pH, which are related to fruit palatability, flesh firmness and size, which characterise fruit appearance, and pericarp thickness, which is related to postharvest injuries and storage, are among the quality traits influencing consumers' tastes. TSS is a crucial quality measure that reflects the amount of sugar in fruit (Flores et al., 2010). According to Lopez-Galarza et al study 's on grafted watermelon, the lower TSS content observed in tomatoes grown on rootstocks may be due to cytokinins produced by the rootstock (2004). More precisely, these researchers offered proof that, after travelling to the scion, cytokinins lower sugar content. Additionally, according to Bari and Jones (2009), the plant can respond by producing more cytokinins when diseases are present (Bari and Jones, 2009). However, the current research found that tomatoes grafted on specific rootstocks had slightly reduced TSS content (o Brix) and dry matter in their fruit, but these values fell within the normal range for tomatoes and are therefore not regarded as particularly significant (Flores et al., 2010) Additionally, according to Buller et al. (2013), grafting on "Beaufort" had no effect on the fruit TSS of the tomato cultivars "Cuore di Bue," "oxheart," and "Cherokee Purple." Additionally, there were no variations in TSS content between tomato plants grown in open fields and greenhouses that were ungrafted and grafted on the 'Heman' and 'Primavera' rootstocks, respectively (Khahet al., 2006). In contrast to these results, Pogonyiet al. (2005) found that tomato plants grafted on the 'Beaufort' variety had lower levels of soluble solids in their fruit. According to Turhan et al. (2011), tomato grafting on 'Beaufort' increased TSS content based on the cultivars used as a scion but had no impact on pH. Harvesting of fruits was also 7 days early for the grafted tomato plants. When they grafted eggplant cultivars onto perennial and wild Solanaceous species, Gisbert et al. (2011) and Rahman et al. (2002) also noted a similar finding. When grafted onto robust rootstock, tomatoes' stomatal conductance increased, enhancing water and nutrient absorption (Fernandez- Garcia et al., 2004a). Crop yield is raised by grafted plants' increased photosynthesis rates (Matsuzoeet al., 1993). When eggplants were grafted onto *S. torvum*, the fruit size grew without affecting the quality or yield. The variety of rootstock used and

grafting can have an impact on the amount of sugar, flavour, colour, and carotene as well as on the texture (Davis et al., 2008). Contrarily, grafting eggplant on *Solanum torvum* and *Solanum sisymbriifolium* had a detrimental impact on the amount of vitamin C, the firmness, and some sensory characteristics, but the general impression was unaffected (Arvanitoyannis et al., 2005). By grafting tomato "Oxheart" onto two inter-specific *S. lycopersicum* *S. habrochaites*, Di-Gioia et al. (2010) found no significant differences in total soluble solids. They also discovered that the vitamin C content of tomato plants grafted onto Beaufort F1 and Maxifort F1 dropped by 14–20%. In order to enhance qualitative traits using grafting techniques, more study is required. In domesticated tomatoes, there is little variation for abiotic stresses, particularly salt tolerance. However, wild species of the genus *Solanum*, including *S. pimpinellifolium*, *S. peruvianum*, *S. cheesmaniae*, *S. habrochaites*, *S. chmielewskii*, and *S. pennellii*, had sources of tolerance that were used as rootstocks to increase productivity under a variety of environmental stress conditions. However, introgression with commercial tomato is challenging because of the issue of crossability. In conclusion, grafting is a very effective method for removing various difficult biotic and abiotic stressors worldwide, particularly in solanaceous vegetable crops. For greater fruit production and quality, proper scion and rootstock selection is essential. This method could be extremely helpful in increasing vegetable productivity in a changing environment. Therefore, it is necessary to find more resistant sources by utilising wild and exceptional germplasm.

#### **ROLE OF GRAFTING ON BIOTIC STRESS TOLERANCE:**

For protected tomato products, soil-borne diseases such as *Fusarium* (*Fusarium oxysporum* f. sp. *lycopersici*, races 1 and 2) and verticillium wilts (*Verticillium dahliae*, races 1 and 2), bacterial speck (*Pseudomonas syringae* v. *tomato*), root knot nematodes (*Melodogyne* spp) (Besri, 2002; Poffley, 2003). After the Copenhagen Amendment to the Montreal Protocol in 1992, most nations around the world phased out methyl bromide by 1 January 2005 because it was added to the list of substances that deplete the ozone layer. Previously, these were regulated by fumigants, primarily methyl bromide (Anon, 2014). Grafting has become an essential instrument with the rapid growth of intensive protected cultivation. This type of resistance developed in the grafted plants may be the result of limited colonisation of bacteria or pathogen in the lower stem and preventing them from invading xylem tissues. Root stocks of certain species have excellent tolerance to various soil-borne diseases like *Fusarium*, *Verticillium*, *Phytophthora*, nematodes, and other pests (Grimault and Prior, 1994). The inherent resistance of the rootstocks and better nutrient uptake, which improve disease control in grafted vegetable plants, could be another factor. When susceptible scions were grafted on particular root stocks, a few trials also claimed to offer some protection from viruses. Mahmoud, (2014) discovered that grafting increased tomato yield components compared to non-grafted plants, delayed the onset of TYLCV symptoms, and raised TYLCV tolerance in susceptible plants. Iouannou (2001) noted that grafted eggplants had higher yields and larger fruits, as well as reduced disease incidences. Similar findings were made when tomatoes (*S. lycopersicum* and *S. habrochaites*) were transplanted onto Beaufort's rootstocks (Hasna et al., 2009). After being grafted onto commercial rootstock and infected with *Verticillium dahliae*, tomato plants (*Solanum lycopersicum* L.) showed resilience to the verticillium wilt disease (Paplomatas et al., 2002). *Capsicum annum* L. scions are grafted onto *C. annum* rootstocks that have resistance to nematodes and soilborne disease in a recent technique known as chilli pepper grafting (Morra and Biloto 2006). When grafted onto resistant CRA 66 or Hawaii 7996 tomato rootstocks, 'German Johnson' heritage tomatoes had 0% fusarium wilt incidence in



- Arvanitoyannis Ioannis, Khah Ebrahim M, Christakou Evangelia C. and Bletsos Fotios A. (2005). Effect of grafting and modified atmosphere packaging on eggplant quality parameters during storage. *International Journal of Food Science & Technology* 40(3):311 – 322.
- Atasayar ,A., 2006. The usage of grafted watermelon seedling in Turkey. *Hasad Horticulture Magazine*, 252: 87–91.
- AVRDC 2003. Guide. Grafting tomatoes for the production in the hot-wet season. Asian Vegetable Research and Development Center, Pub. No 03-551, Shanhua, Tainan, Taiwan, 6pp.
- AVRDC 2009. Guide. Grafting sweet peppers for the production in the hotwet season. Asian Vegetable Research and Development Center, Pub. No 09- 722, Shanhua, Tainan, Taiwan, 8pp.
- Bari, R., Jones, D.G., 2009. Role of plant hormones in plant defence responses. *Plant Molecular Biology*. 69, 473e488.
- Besri, M. 2002. Tomato grafting as an alternative to methyl bromide in Morocco. Institute Agronomieque et Veterinaire Hasan II. Morocco.
- BletsosFotios A. and OlympiosChristos.,2008. The European journal of Plant Science and Biotechnology 2 (1) 62-73.
- Bletsos, F.A., 2005. Use of grafting and calcium cyanamide as alternatives to methyl bromide soil fumigation and their effects on growth, yield, quality and fusarium wilt control in melon. *J. Phytopathol.* 153, 155–161
- Bolandnazar, S., Moghbeli, E.M., Panahandeh, J. and Arzanlou, M. 2014. Biological control of fusarium wilt in greenhouse tomato by Mycorrhizal fungi and resistant rootstock. *Acta Horticulturae*. 1041:127-132.
- Buller, S., Inglis, D., Miles, C., 2013. Plant growth, fruit yield and quality, and tolerance to Verticillium wilt of grafted watermelon and tomato in field production in the Pacific Northwest. *Horticulture Science* 48.
- Buller, S., Inglis, D. & Miles, C. 2013 Plant growth, fruit yield and quality, and tolerance to verticillium wilt of grafted watermelon and tomato in field production in the Pacific Northwest *HortScience* 48 1003 1009
- Colla ,G., Suarez, CMC., Cardarelli ,M., Roupheal ,Y., 2010 Improving nitrogen use efficiency in melon by grafting. *Horticulture Science* 45: 559-565.
- Colla, G., Roupheal Y, Cardarelli M, O, Temperini., E, Rea., Salern., A, Pierandrei F, 2008. Influence of grafting on yield and fruit quality of pepper (*Capsicum annum* L.) grown under greenhouse conditions. *Acta Horticulturae* (ISHS) 782, 359–364.
- Colla, G., Roupheal, Y., Cardarelli, M., Salerno, A., Rea, E., 2010a The effectiveness of grafting to improve alkalinity tolerance in watermelon. *Environment Experiment Botany* 68: 283-291.
- Davis, AR., Perkins-Veazie, P., Hassell, R., Levi, A., King, SR., Zhang, X., 2008. Grafting Effects on Vegetable Quality. *Horticulture Science* 43(6): 1670-1672.
- Di-Gioia F, Serio F, Buttaro D, Ayala O, Santamaria P. Influence of rootstock on vegetative growth, fruit yield and quality in 'Cuore di Bue', an heirloom tomato *Journal of Horticultural Sciences and Biotechnology* 2010;85(6):477-482.
- Dowarah, Buddhisatya & Pal, Subhajit. (2022). Role of Rootstock in Crop Improvement: A Review. 10.30954/ndp/intpest.26.

- Fernández-García, N., Carvajal M., and Olmos. E. 2004. graft union formation in tomato plants: peroxidase and catalase involvement. *Annals of Botany* 93(1):53-60.
- Fernandez-Garcia, N., Martinez, V., Cerda, A., and Carvajal, M. 2004b. Fruit quality of grafted tomato plants grown under saline conditions. *Journal of Horticultural Science and Biotechnology*. 79: 995–1001.
- Flores F.B., Sanchez-Bel P., Estan M.T., Martinez-Rodriguez M.M., Moyano E., Morales B., Campos J.F., Garcia-Abellan J.O., Egea M.I., Fernandez-Garcia N., Romojaro F., Bolarín M.C., 2010. The effectiveness of grafting to improve tomato fruit quality. *Scientia Horticulturae* 125: 211-217.
- García-Rodríguez M.R., Chiquito-Almanz., E, LoezaLara., P.D, Godoy-Hernández., H, Villordo-Pineda., E, PonsHernández., J.L, González-Chavira., M.M, AnayaLópez., J.L, 2010. Production of ancho chili graft on criollo de morelos 334 for the control of *Phytophthora capsici*. *Agrociencia* 44, 701–709.
- Gebologlu N, Yilmaz E, Cakmak P, Aydin M and Kasap Y 2011. Determining of the yield, quality and nutrient content of tomatoes grafted on different rootstocks in soilless culture. *Scientific Research and Essays* 6(10): 2147- 2153.
- Gisbert C., Sánchez-Torres, P. RaigónM.D. and F. Nuez 2010. *Phytophthora capsici* resistance evaluation in pepper hybrids: Agronomic performance and fruit quality of pepper grafted plants. *Journal of Food, Agriculture and Environment* 8, 116–121.
- Gisbert, C., Prohens, J., Raigón, M.D., Stommel, J.R. and Nuez, F. 2011. Eggplant relatives as sources of variation for developing new rootstocks: Effects of grafting on eggplant yield and fruit apparent quality and composition. *Scientific Horticulturae*.128(1): 14-22.
- Grimault, V. and Prior, P. 1994. Grafting tomato cultivars resistant or susceptible to bacterial wilt: Analysis of resistance mechanisms. *Journal of Phytopathology*. 141: 330–334.
- Hasna, M.K., Ogren, E., Persson, P., Ma´rtensson, A. and Ramert, B. 2009. Management of corky root disease of tomato in participation with organic tomato growers. *Crop Protection*. 28: 155– 161.
- Ibrahim, A., Wahb-Allah,M., Abdel-Razzak ,H., Alsadon, A., 2014. Growth, yield, quality and water use efficiency of grafted tomato plants grown in greenhouse under different irrigation levels. *Life ScienceJournal* 11:118-126
- Jang, Y., Yang, E., Cho, M., Um, Y., Ko, K., Chun, C. 2012. Effect of grafting on growth and incidence of phytophthora blight and bacterial wilt of pepper (*Capsicum annum* L.). *Horticulture, Environment, and Biotechnology*. 53(1): 9–19
- JohkanMasahumi,Mitukuri Kazuhiko,Yamasaki Satoshi and MoriGenjirou(2008a). Causes of defoliation and low survival rate of grafted sweet pepper plants. *Scientia Horticulturae* 119(2):103-107
- Johkan M., K. Mitukuri, S. Yamasaki, G. Mori and M. Oda, (2009b). Causes of defoliation and low survival rate of graft -ed sweet pepper plants. *Scientia Horticulturae* 119, 103–107.
- Johnson, S., Kreider, P., Miles C. 2011. Vegetable Grafting Eggplants and Tomatoes.Washington State University 4.
- Keatinge, J.D.H., Lin, L.-J., Ebert, A.W., Chen, W.Y., Hughes, J.d'A., Luther, G.C., Wang, J.-F. Ravishanka, M., 2014. Overcoming biotic and abiotic stresses in the Solanaceae through grafting: current status and future perspectives. *Biological Agriculture & Horticulture*. 30(4): 272-287.

- Khah E.M., Kakava E., Mavromatis A., Chachalis D., Goulas C., 2006. Effect of grafting on growth and yield of tomato (*Lycopersicon esculentum* Mill.) in greenhouse and open-field. *Journal of Applied Horticulture*, 8: 3–7.
- Khah, E. M. 2005. Effect of grafting on growth, performance and yield of aubergine (*Solanum melongena* L.) in the field and greenhouse. *Journal of Food, Agriculture & Environment*. 3(3&4): 92 - 94.
- Lakshmi, T., Tirupathamma., Venkata,C.,Ramana, Naram ., L, Naidu ., Sasikala ,K., 2019. Vegetable Grafting: A Multiple Crop Improvement Methodology. *Current Journal of Applied Science and Technology*. 33(3): 1-10.
- Lee JM, Oda M (2003) Graing of herbaceous vegetable and ornamental crops. *Hort Rev* 28: 61-124.
- Lee, Jung-Myung., Kubota, C., Tsao, S.J., Bie, Z., Echevarria, H., Morra, P.L. and Oda, M.2010. Current status of vegetable grafting: Diffusion, grafting techniques, automation. *Scientia Horticulturae*. 127(2,8): 93-105.
- Lopez-Galarza S.,San Bautista A.,Perez D. M. and Miguel A. (2004). Effects of grafting and cytokinin – Induced fruit setting on colour and sugar – Content traits in glasshouse – Grown triploid watermelon. *Journal of Horticultural Science and Biotechnology* 79(6):971-976
- López-Marín J., A. Gálvez, A. González and J.A. Fernández, 2009. Agronomic behaviour of grafted sweet pepper grown in a greenhouse in Mediterranean area. *Acta Horticulturae (ISHS)* 807, 655–660.
- López-Marín, J., Gálvez, A., Del Amol, FM., Albacete, A., Fernandez, JA., Egea-Gilabert, C., 2017. Selecting vegetative/ generative/dwarfing rootstocks for improving fruit yield and quality in water stressed sweet peppers. *Science Horticulture*; 214:9-17.
- M'hamdi M., N. Boughalleb, N. Ouhaibi, N. Tarchoun, M. Souli and L. Belbahri, 2010. Evaluation of grafting techniques and a new rootstock for resistance of pepper (*Capsicum annum* L.) towards *Phytophthora nicotianae*. *Journal of Food, Agriculture and Environment* 8, 135–139.
- Mahmoud, M.A. 2014. Grafting as a Tool to Improve TYLCV-Tolerance in Tomato. *Journal of Horticultural Science & Ornamental Plants*. 6 (3): 109-115.
- Martinez-Rodriguez, MM., Estan, MT., Moyano, E., Garcia-Abellan ,JO., Flores ,FB., 2008 The effectiveness of grafting to improve salt tolerance in tomato when an 'excluder' genotype is used as scion. *Environ Exp Bot* 63: 392-401.
- Matsuzoe, N., Nakamura, H., Okubo, H., and Fujieda, K. 1993. Growth and yield of tomato plants grafted on *Solanum* rootstocks. *Journal of the Japanese Society for Horticultural Science*. 61:847-855.
- Michel, V.V., Lazzeri, L., 2010. Green manures and organic amendments to control corky rot of tomato. *Acta Hortic*. 883, 287e294.
- Morra ,L., M, Bilotto .,2006. Evaluation of new rootstocks for resistance to soilborne pathogens and productive behavior of pepper (*Capsicum annum* L.) *Journal of Horticultural Science and Biotechnology* 81, 518–524.
- Morra, L. and Bilotto, M. 2006 Evaluation of new rootstocks for resistance to soilborne pathogens and productive behavior of pepper (*Capsicum annum* L.) *Journal of Horticultural Science and Biotechnology*. 81:518-524.

- Nilsen, ET., Freeman, J., Grene ,R., Tokuhisa, J., A rootstock provides water conservation for a grafted commercial tomato (*Solanum lycopersicum* L.) line in response to mild-drought conditions: a focus on vegetative growth and photosynthetic parameters. *Plos one* 2014;9(12): e115380.
- Onduso, J. N. 2014. Management of bacterial wilt of tomato by use of resistant rootstock. Masters of science Thesis. University of Nairobi, Kenya.
- Otani, T., Seike, N., 2007 Rootstock control of fruit dieldrin concentration in grafted cucumber (*Cucumissativus*). *J Pestic Sci* 32: 235-242.
- Palada, M.C., D.L. Wu., 2008. Evaluation of chili rootstocks for grafted sweet pepper production during the hot-wet and hot-dry seasons in Taiwan. *Acta Horticulturae (ISHS)* 767, 151–158. [http://www.actahort.org/books/767/767\\_14.htm](http://www.actahort.org/books/767/767_14.htm)
- Pandey., A.K., Rai, M., 2003. Prospects of grafting in vegetables: an appraisal. *Vegetable science*. 30(2): 101-109.
- Paplomatas, E.J., Elena, K., Tsagkarakou, A. and Perdikaris, A.2002. Control of *Verticillium* wilt of tomato and cucurbits through grafting of commercial varieties on resistant rootstocks. *Acta Horticulturae*. 579:445–449.
- Peddy, P.,1986 Analysis of crop losses in certain vegetables due to *Meloidogyne incognita*. *Int Nematol Net New let* 3: 3-5. 15.
- Penella C, Nebauer SG, San Bautista A, López-Galarza S, Calatayud A. 2014b Rootstock alleviates PEG-induced water stress in grafted pepper seedlings: physiological responses. *Journal of Plant Physiology* 171:842-851.
- Petran, AJ., Interspecific Grafting of Tomato (*Solanum lycopersicum*) Onto Wild Eggplant (*Solanum torvum*) for Increased Environmental Tolerances. M.Sc. Thesis University of Minnesota, Minneapolis, MN 2013.
- Poffley, M. 2003. Grafting tomatoes for bacterial wilt control. *Agnote*, 603, No. B40.
- Pogonyi, A., Peka, Z., Helyes, L., Lugasi, A., 2005. Effect of grafting on the tomato's yield, quality and main fruit components in spring forcing. *Acta Aliment.* 34, 453e462.
- Pulgar, G., Villora, G., Moreno, DA., Romero L (2000) Improving the mineral nutrition in grafted watermelon plants: nitrogen metabolism. *Biol Plant* 43: 607-609.
- Rahman, M.A., Rashid, M.A., Salam, M.A., Masud, M.A.T., Masum, A.S.M.H., and Hossain, M.M. 2002. Performance of some grafted eggplant genotypes on wild *Solanum* root stocks against rootknot nematode. *Journal of Biological Sciences* 2:446-448.
- Rivard, C. L., O'Connell, S., Peet, M M., and Louws, F. J. 2010. Grafting Tomato with Interspecific Rootstock to Manage Diseases Caused by *Sclerotium rolfsii* and Southern Root-Knot Nematode. *Plant Disease*. 94(8): 1015- 1021.
- Rivard, C.S. and F.J. Louws. 2008. Grafting to Manage Soilborne Disease in Heirloom Tomato Production. *HortScience*. 43(7): 2104-2111.
- Rivero, RM., Ruiz, JM., Romero, L., 2003 Role of grafting in horticultural plants under stress conditions. *J Food Agric Environ* 1: 70-74.

- Romano Daniela and A. Paratore (2001). Effects of grafting on tomato and eggplant. *Acta Horticulturae* 559(559):149-154
- Rouphael Youssef, Schwarz Dietmar, Krumbein Angelika and Colla Giuseppe (2010). Impact of grafting on product quality of fruit vegetables. *Scientia Horticulturae* 127(2):172-179
- Rouphael, Y., Cardarelli, M., Colla, G., Rea, E., 2008 Yield, mineral composition, water relations, and water use efficiency of grafted mini watermelon plants under deficit irrigation. *Hort Sci* 43: 730-736.
- Sanchez-Rodriguez, E., Ruiz, J. M., Ferreres, F. and Moreno, D.A. 2012. Phenolic Metabolism in Grafted versus Nongrafted Cherry Tomatoes under the Influence of Water Stress. *Journal of Agricultural and Food Chemistry*. 59(16):8839-8846.
- Savvas, D., Colla, G., Rouphael, Y., Schwarz, D., 2010 Amelioration of heavy metal and nutrient stress in fruit vegetables by grafting. *Sci Hort* 127: 156-161.
- Thakur Diksha., Savita 2020. Role of grafting in vegetable crops: A review. *Journal of Pharmacognosy and Phytochemistry*, 9(6): 1170-1174
- Tuberosa, R., Phenotyping for drought tolerance of crops in the genomics era. *Front. Physiol* 2012; 3:347.
- Turhan, A., Ozmen, N., Serbeci, M.S., Seniz, V., 2011. Effects of grafting on different rootstocks on tomato fruit yield and quality. *Hort. Sci.* 38, 142e149.
- Turhan, A., Ozmen, N., Serbeci, M.S. & Seniz, V. 2011 Effects of grafting on different rootstocks on tomato fruits yield and quality *Hort. Sci. (Prague)* 38 142 149
- Venema, JH., Dijk, BE., Bax, JM., Hasselt, P., Elzenga, JTM., 2008 Grafting tomato (*Solanum lycopersicum*) onto the rootstock of a high-altitude accession of *Solanum habrochaites* improves sub-optimal temperature tolerance. *Environ Exp Bot* 63: 359-367.
- Vitale, A., Rocco, M., Arena, S., Giuffrida, F., Cassaniti, C., Scaloni, A., Lomaglio, T., Guarnaccia, V., Polizzi, G., Marra, M. and Leonardi, C. 2014. Tomato susceptibility to fusarium crown and root rot: effect of grafting combination and proteomic analysis of tolerance expression in the rootstock. *Plant Physiology and Biochemistry*. 83:207- 216.