

Original Research Article

Effect of Organic and Inorganic Mulches on Quality and Soil parameters on ridge gourd

Formatted: Centered

Abstract

Aim: To study the effect of different mulches on quality and soil parameters on ridge gourd

Study design: The experiment was carried out with 7 treatments in Randomized Block Design (RBD) with three replications.

Place and duration of study: The research Research trial was carried out at P.G block, College of Horticulture, Rajendranagar, SKLTSU, Hyderabad during *Kharif*, 2021 and Summer, 2022.

Results: Among the organic and inorganic mulch mulches treatments, the results reported that the silver-silver-black polyethylene mulch recorded the highest highest in quality parameters and benefit-cost ratio compared to other mulches.

Comment [K1]: Please compared factors with data or percent

Keywords: Ridge gourd; Organic mulches; Inorganic mulches; Quality, Soil

1. INTRODUCTION

Comment [K2]: Need to be complete with the references about mulchs

Ridge gourd (*Luffa acutangula*) is a popular warm-season vegetable that may be grown commercially from seeds. It can be found throughout the tropical and subtropical regions and is cultivated for its green tender fruits during spring-summer and rainy seasons when the temperature ranges between 20 - 32°C. The unripe fruits are consumed as cooked vegetables a cooked vegetable and nutritionally the fruits are rich in several minerals and vitamins. Each 100 g of edible fresh fruit contains around 5 mg vitamin C, 0.01mg riboflavin, 33 µg carotene, 92.5% moisture, 0.5g protein, 0.5 g fat, 3.4 g carbohydrate, 18 mg calcium, 26 mg phosphorous and 0.5 mg iron (1). The application of different doses of nitrogen (N) fertilization increased plant growth and yield (2) Phosphorus is necessary for cellular preparation and in the metabolism of starch, protein and fats. One of the most important effects of phosphorus on plants is the stimulation of early root formation and growth. Low available phosphorus content in the soil means -delayed delay in maturity and poor plant growth (3). Potassium (K) is the nutrient having the strongly influences strongest influence on plant growth, yield and quality attributes that determine fruit marketability (4). Therefore, considering the importance of different mulching in various vegetable crops, data and the facts, the present investigation entitled Studies on the effect of mulches on quality and soil parameters on ridge gourd (*Luffa acutangula* L.)” was planned to be carried out in the growing region of Hyderabad, Telangana, India.

Comment [K3]: Please add more references in this subject

2. MATERIALS AND METHODS

2.1 The experimental site

Comment [K4]: Need to be complete

The present investigation was carried out during *Kharif*, 2021 and ~~Summersummer~~, 2022 at P.G block, College of Horticulture, Rajendranagar, SKLTSHU, ~~Hyderabadand~~ Hyderabad, India.

2.2 The experimental ~~ly~~ variants and the experiment design

The plots were demarcated into three (3) replications, each replication with seven treatments and the experimental design ~~followed iswas~~ Randomized Block Design (RBD) replicated thrice consisting of Paddy straw- T₁, Paddy husk- T₂, Groundnut shell- T₃, Leaf mould-T₄, Silver black polyethylene sheet-T₅, Clear transparent polyethylene mulch-T₆ and Control-T₇ ~~which with three~~ replication ~~ioned thrices~~.

2.3 The determined parameters ~~determined~~

2.3.1 Quality parameters

The recorded data ~~reecorded~~ on quality parameters like TSS (° Brix), ascorbic acid content (mg/100g), reducing sugars (%), ~~non-non~~-reducing sugars (%), total sugars (%), chlorophyll content (DA meter reading) and crude fibre content (%) were ~~analysed~~ analyzed.

2.3.2 Soil parameters

The data recorded on soil parameters like pH, EC (dS/m), organic carbon (%), available nitrogen (kg/ha), available phosphorus (kg/ha), available potassium (kg/ha) were ~~analysed~~ analyzed.

2.4 Statistical analysis

The experimental collected data ~~collected~~ on various quality, soil components of plants were subjected to Fisher's method of "Analysis of variance" (ANOVA) as outlined by (5) were ~~analysed~~ analyzed.

3. RESULTS AND DISCUSSION

3.1 Quality parameters

TSS (° Brix)

The enuniciated data ~~enuniciated~~ on the TSS (° Brix) as influenced by the effect of mulching on quality in ridge gourd ~~is was~~ presented in Table 1.

Comment [K5]: Need to be complete the methods

Comment [K6]: Need to be complete the methods

Table 1. Effect of mulching on TSS (^o Brix), ascorbic acid content (mg/100g) of ridge gourd during *Kharif*, 2021 and ~~Summersummer~~, 2022

Treatments	TSS (^o Brix)		Ascorbic acid content (mg/100g)	
	Kharif 2021	Summer 2022	Kharif 2021	Summer 2022
T ₁ : Paddy straw	3.89	4.23	10.21	10.43
T ₂ : Paddy husk	3.55	3.49	9.49	9.75
T ₃ : Groundnut husk	3.63	3.56	9.56	9.89
T ₄ : Leaf mould	3.60	3.35	9.41	9.64
T ₅ : Silver black polythene sheet	4.64	4.82	12.82	13.39
T ₆ : Clear transparent polythene sheet	3.97	4.58	11.58	12.64
T ₇ : Without mulch	2.98	3.27	9.25	9.47
S.Em±	0.05	0.05	0.12	0.13
CD at 5%	0.15	0.15	0.38	0.40

Formatted: Left

Formatted: Left

From the data, it is clear that there ~~was-were~~ significant differences observed among the treatments with respect to TSS during the *Kharif* season. Significantly, the maximum TSS (4.64) was recorded in T₅ (Silver black polyethylene mulch) followed by T₆ (clear transparent polyethylene mulch) (3.97), while the minimum TSS (2.98) was recorded with T₇ (without mulch).

During ~~Summersummer~~ season among all the treatments, T₅ (Silver black polyethylene mulch) recorded maximum TSS (4.82) followed by T₆ (clear transparent polyethylene mulch) (4.58), while the minimum TSS (3.27) was recorded ~~with-in~~ T₇ (without mulch).

Ascorbic acid content (mg/100g)

The data presented in Table 1 revealed that during *Kharif* season, significantly, the maximum ascorbic acid content (12.82) was observed with T₅ (Silver black polyethylene mulch) followed by T₆ (clear transparent polyethylene mulch) recording the ascorbic acid content (11.58). The minimum ascorbic acid content (9.25) was recorded with T₇ (without mulch).

During ~~Summers~~summer season, significantly maximum ascorbic acid content (13.39) was observed with T₅ (Silver black polyethylene mulch) and was at par with T₆ (clear transparent polyethylene mulch) recording the ascorbic acid content of 12.64. The minimum ascorbic acid content (9.47) was recorded with T₇ (without mulch).

Reducing sugars (%)

The ~~enunciated~~ data ~~enunciated~~ on the reducing sugar as influenced by the effect of mulching on quality in ridge gourd is presented in Table 2.

From the data it is clear that there was significant differences observed among the treatments with respect to reducing sugars during *Kharif* season. Significantly maximum reducing sugars (4.86%) was recorded in T₅ (Silver black polyethylene mulch) followed by T₆ (clear transparent polyethylene mulch) (4.51%), while the minimum reducing sugars (3.45%) was recorded with T₇ (without mulch).

During ~~Summers~~summer season among all the treatments, T₅ (Silver black polyethylene mulch) recorded maximum reducing sugars (4.98%) followed by T₆ (clear transparent polyethylene mulch) (4.54%), while the minimum reducing sugars (2.39%) was recorded with T₇ (without mulch).

~~Non-Non~~-reducing sugars (%)

The ~~presented~~ data ~~presented~~ in Table 2 revealed that during *Kharif* and Summer season, significantly minimum non reducing sugars (1.45 and 1.47%) was observed with T₅ (Silver black polyethylene mulch), while the maximum reducing sugars (1.73 and 2.50 %) were recorded with T₇ (without mulch) during *Kharif* and ~~Summer~~-summer season, respectively.

Total sugars (%)

The ~~enunciated~~ data ~~enunciated~~ on the total sugars as influenced by the effect of mulching on quality in ridge gourd is presented in Table 2.

From the data it ~~is~~-was clear that there was significant differences observed among the treatments with respect to total sugars during *Kharif* season. Significantly maximum total sugars (6.31%) was recorded in T₅ (Silver black polyethylene mulch) followed by T₆ (clear transparent polyethylene mulch) (6.03%), while the minimum total sugars (5.18%) was recorded with T₇ (without mulch).

During ~~Summers~~summer season among all the treatments, T₅ (Silver black polyethylene mulch) recorded maximum total sugars (6.45%) and was at par with T₆ (clear transparent polyethylene mulch) (6.33%), while the minimum total sugars (4.89%) was recorded with T₇ (without mulch).

Table 2. Effect of mulching on reducing sugars (%), non-reducing sugars (%), total sugars (%) of ridge gourd during *Kharif*, 2021 and *Summersummer*, 2022

Reatments	Reducing sugars (%)		Non-reducing sugars (%)		Total sugars (%)	
	Kharif 2021	Summer 2022	Kharif 2021	Summer 2022	Kharif 2021	Summer 2022
T ₁ : Paddy straw	3.74	4.27	1.72	1.96	5.41	6.23
T ₂ : Paddy husk	3.69	3.65	1.57	1.52	5.30	5.17
T ₃ : Groundnut husk	3.73	3.60	1.58	2.14	5.32	5.74
T ₄ : Leaf mould	3.56	3.42	1.64	1.76	5.20	5.18
T ₅ : Silver black polythene sheet	4.86	4.98	1.45	1.47	6.31	6.45
T ₆ : Clear transparent polythene sheet	4.51	4.54	1.52	1.79	6.03	6.33
T ₇ : Without mulch	3.45	2.39	1.73	2.50	5.18	4.89
S.Em±	0.05	0.05	0.02	0.03	0.07	0.08
CD at 5%	0.16	0.15	0.07	0.08	0.23	0.23

Formatted: Left

Formatted: Left

Chlorophyll content (DA meter reading)

The data presented in Table 3 revealed that during *Kharif* season, significantly maximum chlorophyll content (1.49) was observed with T₅ (Silver black polyethylene mulch) followed by T₆ (clear transparent polyethylene mulch) recording the chlorophyll content of 1.26. The minimum chlorophyll content (0.87) was recorded with T₇ (without mulch).

During *Summersummer* season significantly maximum chlorophyll content (1.68) was observed with T₅ (Silver black polyethylene mulch) and was at par with T₆ (clear transparent polyethylene mulch) recording the chlorophyll content of 1.52. The minimum chlorophyll content (0.99) was recorded with T₇ (without mulch).

Crude fibre content (%)

The data presented in Table 3 revealed that during *Kharif* season, significantly minimum crude fibre (2.32) was observed with T₅ (Silver black polyethylene mulch) followed by T₆ (clear transparent polyethylene mulch) recording the crude fibre content of 2.53. The maximum crude fibre (2.96) was recorded with T₇ (without mulch).

During *Summersummer* season significantly minimum crude fibre (2.21) was observed with T₅ (Silver black polyethylene mulch) followed by T₆ (clear transparent polyethylene mulch) recording the crude fibre of 2.50. The maximum crude fibre (2.87) was recorded with T₇ (without mulch).

The improvement in the quality parameters by using silver black polyethylene sheet might be due to better availability and uptake of nutrients which in turn might have lead to more nitrogenous compounds in plant tissues and ultimately resulted in their efficient metabolism. (6,7,7)

Table 3. Effect of mulching on chlorophyll content, crude fibre content of ridge gourd during *Kharif*, 2021 and *Summersummer*, 2022

Treatments	Chlorophyll content		Crude fibre content	
	Kharif 2021	Summer 2022	Kharif 2021	Summer 2022
T ₁ : Paddy straw	1.13	1.57	2.58	2.53
T ₂ : Paddy husk	0.99	1.17	2.64	2.60
T ₃ : Groundnut husk	1.14	1.21	2.67	2.58
T ₄ : Leaf mould	0.94	1.06	2.80	2.72
T ₅ : Silver black polythene sheet	1.49	1.68	2.32	2.21
T ₆ : Clear transparent polythene sheet	1.26	1.52	2.53	2.50
T ₇ : Without mulch	0.87	0.99	2.96	2.87
S.Em±	0.01	0.02	0.04	0.04
CD at 5%	0.04	0.05	0.12	0.11

3.2 Soil analysis

pH

It ~~is~~ was clear from Table 4 that pH did not vary significantly with different treatments during *Kharif* and *Summersummer* seasons, respectively.

Table 4. Effect of mulching on P^H, Electrical conductivity (dS/m) and Organic carbon (%) of ridge gourd during *Kharif*, 2021 and ~~Summersummer~~, 2022

Treatments	p ^H		Electrical conductivity (dS/m)		Organic carbon (%)	
	Kharif 2021	Summer 2022	Kharif 2021	Summer 2022	Kharif 2021	Summer 2022
T ₁ : Paddy straw	7.17	7.22	0.287	0.288	0.60	0.64
T ₂ : Paddy husk	7.26	7.28	0.283	0.286	0.55	0.56
T ₃ : Groundnut husk	7.23	7.23	0.287	0.285	0.57	0.60
T ₄ : Leaf mould	7.28	7.35	0.289	0.287	0.51	0.53
T ₅ : Silver black polythene sheet	7.11	7.17	0.296	0.298	0.67	0.68
T ₆ : Clear transparent polythene sheet	7.14	7.20	0.290	0.291	0.64	0.65
T ₇ : Without mulch	7.20	7.22	0.279	0.280	0.49	0.51
S.Em±	0.10	0.10	0.004	0.004	0.01	0.01
CD at 5%	NS	NS	NS	NS	0.02	0.02

EC (dS/m)

The ~~presented~~ data ~~presented~~ in Table 4 there was no significant difference observed between the treatments during *Kharif* and ~~Summersummer~~ seasons, respectively.

Organic carbon (%)

The organic carbon of the soil as influenced by the effect of mulching by different treatments was observed and the relevant data is presented in Table 4. The highest organic carbon (0.67 and 0.68) was recorded in T₅ (Silver black polyethylene mulch) followed by T₆ (clear transparent polyethylene mulch) recording the organic carbon of 0.64 and 0.65 during *Kharif* and ~~Summersummer~~ respectively. The organic carbon was lowest in T₇ (without mulch) (0.49 and 0.51) during *Kharif* and ~~Summersummer~~ seasons, respectively.

The organic carbon progressively increased progressively with the advancement of crop growth. The positive response of mulches on organic carbon might be due to increased microbial activity and decomposition of organic matter in the rhizosphere of the crop. (8, 9)

Available nitrogen (kg/ha)

The data pertaining to available nitrogen in the soil after harvest as influenced by the effect of mulching are presented in Table 5.

The data indicated that among the treatments evaluated, significantly maximum available nitrogen content in the soil (267.85 and 269.75) was observed with T₅ (Silver black polyethylene mulch) and was at par with T₆ (clear transparent polyethylene mulch) (255.91 and 258.64), while the minimum nitrogen content in the soil was recorded with T₇ (without mulch) (244.76 and 246.45) -during *Kharif* and Summer seasons, respectively.

It was also observed that available nitrogen increased in the mulched plot over compare to the non-mulch plot. It might be due to less depletion of nutrients and moisture, improved microbial activity, solarization effect and favourable soil characteristics.

Available phosphorus (kg/ha)

The data about available phosphorus in the soil after harvest as influenced by the effect of mulching are presented in Table 5.

The data indicated that among the treatments evaluated, significantly maximum available phosphorus content in the soil (35.59 and 35.90) was observed with T₅ (Silver black polyethylene mulch) followed by T₆ (clear transparent polyethylene mulch) (33.51 and 34.25), while the minimum phosphorus content in the soil was recorded with T₇ (without mulch) (30.63 and 30.43) during *Kharif* and ~~Summer~~ summer seasons, respectively.

It was observed that available phosphorus increased in mulched plot over the non mulch plot. It might be due to microbial activity, less evaporation of soil moisture content and favourable soil characteristics.

Available potassium (kg/ha)

The data about available potassium in the soil after harvest as influenced by the effect of mulching are presented in Table 5.

The data indicated that among the treatments evaluated, significantly maximum available potassium content in the soil (215.81 and 217.23) was observed with T₅ (Silver black polyethylene mulch) and was at par with T₆ (clear transparent polyethylene mulch) (210.70 and 212.67), while the minimum potassium content in the soil was recorded with T₇ (without mulch) (189.73 and 195.75) ~~during~~ during *Kharif* and Summer seasons.

There was slight increase of available K initially but gradually it decreased consequent to the utilization of K for the crop and also due to the fixation of K in the soil complex.

In general, the uptake of nutrients was comparatively more in mulched plots than non mulched plots. This might be due to better crop development under mulched plots, consequent of soil moisture content and better utilization of soil nutrients- (10, ~~11~~, 11).

Table 5. Effect of mulching on N, P and K of ridge gourd during *Kharif*, 2021 and *Summer*, 2022

Treatments	Nitrogen (kg/ha)		Phosphorous (kg/ha)		Potassium (kg/ha)	
	Kharif 2021	Summer 2022	Kharif 2021	Summer 2022	Kharif 2021	Summer 2022
T ₁ : Paddy straw	257.75	259.32	33.04	33.15	205.50	207.82
T ₂ : Paddy husk	253.07	255.60	32.56	32.88	199.33	201.15
T ₃ : Groundnut husk	250.65	251.53	32.11	32.53	202.95	204.74
T ₄ : Leaf mould	249.22	250.63	31.54	31.75	193.74	199.45
T ₅ : Silver black polythene sheet	267.85	269.75	35.59	35.90	215.81	217.23
T ₆ : Clear transparent polythene sheet	255.91	258.64	33.51	34.25	210.70	212.67
T ₇ : Without mulch	244.76	246.45	30.63	30.43	189.73	195.75
S.Em±	2.89	2.93	0.41	0.44	2.72	2.78
CD at 5%	8.91	9.03	1.27	1.36	8.38	8.56

Benefit cost ratio

The economics involved as influenced by the effect of mulching has been calculated and presented in Table 6.

Among the treatments during *Kharif* season, significantly T₅ (RDF + Silver black polyethylene mulch) recorded the highest gross returns (Rs. 3,588,00), net returns (Rs. 2,459,49.60) with benefit cost ratio of 2.18:1, Whereas the T₇ (RDF + without mulch) recorded the lowest gross returns per hectare (Rs. 2,265,00), net returns per hectare (Rs. 1,276,49.60) with benefit cost ratio of 1.29:1.

During Summer season, T₅ (RDF + Silver black polyethylene mulch) recorded the highest gross returns (Rs. 3,711,00), net returns (Rs. 2,576,99.60) with benefit cost ratio of 2.27:1, Whereas the T₇ (RDF + without mulch) recorded the lowest gross returns per hectare (Rs. 2,398,50), net returns per hectare (Rs. 1,404,49.60) with benefit cost ratio of 1.41:1. (12)

Table 6. Effect of mulching on benefit: cost ratio of ridge gourd during *Kharif*, 2021 and *Summersummer*, 2022

Treatments	Cost of cultivation (Rs/ha)		Gross income (Rs/ha)		Net returns (Rs/ha)		B:C ratio	
	Kharif 2021	Summer 2022	Kharif 2021	Summer 2022	Kharif 2021	Summer 2022	Khari f 2021	Sum mer 2022
T ₁ : Paddy straw	102600.40	103150.40	283800	308700	181199.60	205549.60	1.77	1.99
T ₂ : Paddy husk	101350.40	101900.40	266550	282450	165199.60	180549.60	1.63	1.77
T ₃ : Groundnut husk	102850.40	103400.40	258900	273450	156049.60	170049.60	1.52	1.64
T ₄ : Leaf mould	101350.40	101900.40	247800	258450	146449.60	156549.60	1.44	1.54
T ₅ : Silver black polythene sheet	112850.40	113400.40	358800	371100	245949.60	257699.60	2.18	2.27
T ₆ : Clear transparent polythene sheet	118850.40	119400.40	347550	360900	228699.60	241499.60	1.92	2.02
T ₇ : Without mulch	98850.40	99400.40	226500	239850	127649.60	140449.60	1.29	1.41

4. CONCLUSION

From the study it ~~can~~ could be concluded that silver black polyethylene mulch had shown the positive effects on fruit quality, soil parameters and ~~benefit~~-~~benefit~~-cost ratio in ridge gourd compared to other treatments, hence it was proved to be the best treatment in ridge gourd.

REFERENCES

1. Sheshadri VS, Parthasarthy UA. Cucurbits in vegetable crops In: Bose TK, Kabir J, Maity T K, Parthasarthy VA, Som MG, (editors.). Vegetable crops. 1980; 496-497.
2. Wahocho NA, Maitlo AA, Baloch QB, Kaleri AA, Rajput LB, Talpur NA, et al. Effect of varying levels of nitrogen on the growth and yield of muskmelon (*Cucumis melo* L.), Journal of Basic & Applied Sciences. 2017;13:448-453

Comment [K7]: Need to be discuss the results with the references

Formatted: Font: Italic

Formatted: Font: Italic

3. Meena OP, Meena RK, Dhaka RS, Meena NK, Sharma A. Effect of Nitrogen and Phosphorous Levels on Growth and Yield of Bottle gourd [*Lagenaria siceraria* (Mol.) Standl.] cv. Pusa Naveen. *Int. J Pure App. Biosci.* 2017;5(4):1178-1184.
4. Lester GE, Jifon JL, Stewart WM. Foliar potassium improves cantaloupe marketable and nutritional quality. *Better Crops.* 2007; 91:24-25.
5. Gomez KH, Gomez AA. *Statistical Procedures for Agriculture Research.* John Willy and Sons, Inc., New York. 1984.
6. Lourduraj C, Padmini AK, Rajendran R, Ravi V, Pandiarajan T, Sreenarayanan VV. Effect of plastic mulching on bhendi *Abelmoschus esculentus* (L.) Moench. *South Indian Horticulture.* 1997; 45(3&4): 128-133.
7. Hallidri M. Comparison of different mulching materials on growth, yield and quality of cucumber (*Cucumis sativus* L.). *Acta Horticulturae.* 2001; 559: 49-53.
8. Khokhar UV, Sharma MK, Singh RR. Changes in some physicochemical and microbiological properties of the soil under various systems of floor management in almond (*Prunus amygdalus* Batsch) orchard. *Journal of the India Society of Soil Science.* 2001; 49 (1): 213-215.
9. Mishra LN, Pathak RA, Pandey AK, Bhanu P, Singh SK. Effect of various mulches on the physico-chemical properties of the sodic soil under aonla + guava cropping system. *Acta Horticulture.* 2002; 54 (4): 280-282.
10. Hundel IS, Sandhu KS, Daljeet S, Sandhu MS. Effect of different types of mulches and herbicidal treatments in nutrient uptake in tomato. *Haryana Journal of Horticulture Science.* 2000; 29 (3 & 4): 242-244.
11. Vethamoni PI, Balakrishnan R. Studies on the influence of herbicide, nitrogen and mulching on the nutrient uptake of okra. *Indian Journal of Horticulture.* 1990; 47(2): 233-238.
12. Sushant K, Satish P, Subash S, Shrestha, S. Effect of mulching materials and plant spacing on growth, sex expression and yield of bitter gourd (*Momordica charantia*) cv. Paalee in Chitwan, Nepal. *Azarian Journal of Agriculture.* 2020; 7(1): 1-7.

Formatted: Font: Italic

Formatted: Font: Italic

UNDER PUBLICATION