

Studies on effect of different cutting and varieties on growth parameter of spinach beet (*Beta vulgaris* var. *bengalensis* L.)

Abstract

During the Rabi season, the Department of Horticulture at Babasaheb Bhimrao Ambedkar University Lucknow (U.P.) conducted research on the effect of different cuttings and varieties on the Horticulture Research Farm. The experiment is conducted in a RBD design with 16 treatments and two factor varieties (V₁ Chandrika, V₂ Pusa Jyoti, V₃ All Green, and V₄ Palak Katadar) and cutting (C₀ no cutting, C₁ cutting 1, C₂ cutting 2, and C₃ cutting 3). We found that the effect of cutting on vegetative growth parameters, viz., plant height, number of leaves, leaf length, leaf width, leaf petiole length, and germination percentage, is significant. We found that cutting C₀ had a significant influence on the growth parameters (plant height, number of leaves, leaf length, leaf width, and petiole length), with the minimum observed at C₃, the maximum germination observed at C₁ as compared to C₂ treatment, and the least germination recorded at C₀ treatment. While plant height, number of leaves, leaf length, leaf width, petiole length, and germination percentage have a greater influence on Pusa jyoti (V₂) than on All Green (V₃), minimums are reported for Chandrika (V₁).

Keywords:- Cutting, Varieties, spinach beet, Experiment, RBD.

Introduction

Spinach beet (*Beta vulgaris* var. *bengalensis* L.) is one of the most important leafy vegetables in the winter. Indian spinach belongs to the "Chenopodiaceae" family and originated in the Indo-Chinese region. The leafy vegetables have the highest nutritional value of other leafy vegetable (Fageria et al. 2022). They are rich in minerals and hence can be called "mines of minerals." It is one of the most nutritious leafy vegetable crops in tropical and sub-tropical regions. Spinach is a rich source of Vitamin A (9770 IU) and Vitamin C (70 mg/100 g). Spinach is high in nutrients. It has 3.4 g of protein, 0.8 g of fat, 380mg of calcium, 30 mg of potassium, and 16.2 mg of iron (Singh 2019). The oxalic acid content of spinach beet is low. Palak is closely related to beet root, Swiss chard (*Beta vulgaris* var. *cicla*), and sugar beet. The popular spinach beet growing states include Uttar Pradesh, West Bengal, Maharashtra, and Gujarat. However, spinach beet is not very popular in south India. It is primarily used as a pot herb. Spinach leaves are valued for their medicinal properties. The leaves are used to treat inflammation, paralysis, headaches, and earaches, as well as spleen and liver diseases. Leaves of this crop might have been first used in Bengal, and hence it is known as *B. vulgaris* var. *bengalensis*. Indian spinach is one of the leafy vegetables that should be considered for future development. It is fairly tolerant and can be grown successfully on saline soil (Mishra et al. 1973).

It is cultivated for fresh, green leaves, which become ready to harvest (cut) in about 30 to 35 days from sowing. Because most leafy vegetables can be cut multiple times, they require a lot of fertilizer to keep up with their fast growth. To increase yield per unit area and per unit time, a farmer discovered maximum leaves by increasing cutting in a specific variety.

The goal of our research is to find the best combination of varieties that produce the most cutting in order to maximize yield.

Materials and methods

The experiment is conducted on Horticultural Research Farm-1, Babasaheb Bhimrao Ambedkar University, Lucknow, 226 025 (U.P.), during the Rabi season of 2019-20. The experimental site is located at 26° 50' N latitude, 80° 52' E longitude, and 123° MSL. Weather data are reported by the IISR. In general, the maximum temperature in the summer is (43⁰C) and the minimum temperature in the winter is (2⁰C). The soil at the experiment field is sandy loam that is slightly alkaline. Used four varieties (V₁ Chandrika, V₂ Pusa Jyoti, V₃ All Green, and V₄ Palak Katadar) and three cuttings (C₀ no cutting, C₁ cutting 1, C₂ cutting 2, and C₃ cutting 3) in a factorial RBD design with 16 treatments and 3 replications. We used that treatment combination: zero cutting + Chandrika, zero cutting + Pusa Jyoti, zero cutting + All Green, zero cutting + Palak Katadar, one cutting + Chandrika, one cutting + Pusa Jyoti, one cutting + All Green, one cutting + Palak Katadar, two cutting + Chandrika, two cutting + Pusa Jyoti, two cutting + All Green, two cutting + Palak Katadar, three cutting + Chandrika, three cutting + Pusa Jyoti Crops are sown on 10 November 2019 during the Rabi season. Data is recorded in growth attributes such as plant height (cm), number of leaves, leaf length (cm), leaf width (cm), leaf petiole length (cm), and germination percentage (%). A statistical analysis of data obtained in different sets of experiments was calculated following the standard procedure of **Panase and Sukhatme (1985)**.

Result and discussion

Effect of cutting on varieties considering in following group.

Effect of cutting:

On growth parameter

Data recorded for growth attributes at regular intervals of 30, 50, and 70 DAS are shown in Table-1. Growth attributes that significantly influence cutting at 70 DAS, viz., plant height, number of leaves, leaf length, width of leaves, and length of petiole. maximums recorded (plant height, number of leaves, leaf length, width of leaves, and length of petiole) in the control (no cutting) C₀ respectively (49.11 cm), (24.33), (12.49 cm), (7.09 cm), (7.09 cm), and minimum data reported at growth attributes (plant height, number of leaves, leaf length, width of leaves, and length of petiole) C₃ (three cutting) respectively (18.29 cm), (14.17 cm), (8.78 cm), (4.99 cm), and (4.99 cm) its result is due to maximum C₀ allowed to thought out the life period undisturbed without cutting but C₃ treatment plot has to faced three cutting conformed **Singh et al. (2015) Dadiga et al. (2015), Naik et al. (2010)**. While in case of germination (%) maximum germination is observed C₁ (78.87 %) treatment followed by the C₂ (78.04 %) treatment and minimum germination reported C₀ (74.96%) treatment similar result are reported by **Bhore et al (2000)**.

Effect of varieties

Data recorded for growth attributes at regular intervals of 30, 50, and 70 DAS are shown in Table -1. varieties also significantly influence growth attributes viz., plant height, number of leaves, leaf length, width of leaves, and length of petiole, showed that V₂ (32.17 cm) recorded maximum height and minimum height was recorded in variety V₁ (29.13 cm) at 70DAS, where in number of leaves per plant maximum V₂ (19.58) and followed by V₃ (18.33) and

minimum are reported in V₁ (16.83) at 70 DAS, in case of length of leaves V₂ (Pusa joyti) (11.31 cm) is maximum as compare to V₃ (10.71 cm) and minimum recorded V₁ (9.75 cm) at 70DAS, width of leaves highest recorded in variety V₂ (6.51 cm) followed by V₃ (6.05 cm) and minimum reported in V₁ (5.49 cm) at 70DAS, while in case of length of petioles V₂ (Pusa Jyoti) are recorded in maximum length of petiole 6.51 cm and minimum recorded in V₁ (Chandrika) 5.49 cm result corroborate the finding of **Thakral et al. (1991)** **Waseem et al. (2000)**. While in germination (%) Pusa Jyoti V₂ (78.49%) having highest germination and Chandrika V₁ (75.86%) least germination percentage was recorded closely result reported by **Singh et al. (2013)** and **Waseem et al. (2000)**.

Interaction

Plant height effects of combination maximum plant height (52.14 cm) at 70 DAS with C₀V₂ minimum plant height C₂V₁ (17.70 cm). Combination C₀V₂ had the highest number of leaves per plant (27.67), while combinations C₂V₁ and C₃V₁ had the lowest number of leaves (14.00 leaves per plant) at 70 DAS. The maximum length of leaves (14.68 cm) with C₀V₂ and minimum length of leaves (8.58cm) of the combination C₃V₁ were reported in length of leaves 70DAS. C₀V₂ (8.03 cm) is the maximum width of leaves, followed by C₀V₃ (7.22 cm), and the minimum width of leaves is recorded in conjunction with C₃V₁ (4.80 cm). In 70 DAS, combination C₀V₂ had the longest petiole at 8.03 cm, and combination C₃V₁ had the shortest at 4.80 cm. The C₁V₂ (80.04 cm) combination recorded the maximum germination percentage, while the minimum germination percentage was recorded in the C₀V₁ (72.71 cm) combination Similar result were also reported by **Bradely et al. (1971)** and **Singh et al. (2015)**.

Table-1Effect of different cutting and varieties on growth parameter.

Treatments	Plant height (cm)			Number of leaves			Length of leaves (cm)		
	30DAS	50DAS	70DAS	30DAS	50DAS	70DAS	30DAS	50DAS	70DAS
V ₁	17.28	22.32	29.13	10.50	14.17	16.83	7.66	10.59	9.57
V ₂	20.78	25.29	32.17	12.17	17.25	19.58	8.42	11.99	11.31
V ₃	19.85	23.86	31.44	11.58	16.00	18.33	8.09	11.54	10.71
V ₄	19.14	23.42	30.46	11.00	15.67	17.83	7.99	11.13	10.49
SEM±	0.23	0.16	0.19	0.25	0.19	0.21	0.14	0.19	0.19
CD at 5%	0.68	0.46	0.54	0.73	0.56	0.60	0.42	0.55	0.56
Cutting									
C ₀	18.25	35.30	49.11	10.67	18.92	24.33	7.39	12.13	12.49
C ₁	19.84	19.95	37.41	11.83	14.33	19.83	8.68	10.83	11.16
C ₂	19.45	20.05	18.38	11.67	15.00	14.25	8.43	11.38	9.82
C ₃	19.48	19.60	18.29	11.08	14.83	14.17	7.65	10.83	8.78
SEM±	0.23	0.16	0.19	0.25	0.19	0.21	0.14	0.19	0.19
CD at 5%	0.68	0.46	0.54	0.73	0.56	0.60	0.42	0.55	0.56

Table-2Effect of different cutting and varieties on growth parameter.

Treatments	Width of leaves (cm)			leaf petiole length (cm)			Germination (%)
	30DAS	50DAS	70DAS	30DAS	50DAS	70DAS	
V ₁	5.16	5.50	5.49	5.30	6.24	5.49	75.86

V ₂	5.82	6.57	6.51	6.38	7.30	6.51	78.49
V ₃	5.47	6.32	6.05	6.03	6.83	6.05	77.41
V ₄	5.35	6.16	5.85	5.72	6.54	5.85	76.73
SEm±	0.14	0.19	0.18	0.15	0.16	0.36	0.17
CD at 5%	0.40	0.54	0.52	0.42	0.45	0.53	0.50
Cutting							
C ₀	4.70	6.61	7.09	4.67	7.40	7.09	74.96
C ₁	5.81	5.76	6.21	6.84	6.16	6.21	78.87
C ₂	5.73	6.53	5.59	6.12	7.20	5.59	78.04
C ₃	5.57	5.64	4.99	5.80	6.15	4.99	76.62
SEm±	0.14	0.19	0.18	0.15	0.16	0.36	0.17
CD at 5%	0.40	0.54	0.52	0.42	0.45	0.53	0.50

Table-3 Interaction effect different cutting and varieties on growth of palak.

Treatments	Plant height (cm)			Number of leaves			Length of leaves (cm)		
	30DAS	50DAS	70DAS	30DAS	50DAS	70DAS	30DAS	50DAS	70DAS
C ₀ V ₁	14.04	32.29	44.09	8.67	14.33	19.33	6.83	10.86	9.89
C ₀ V ₂	21.39	37.77	52.14	11.67	22.67	27.67	7.75	13.01	14.68
C ₀ V ₃	19.60	36.02	51.71	11.33	19.67	25.33	7.53	12.35	13.06
C ₀ V ₄	17.98	35.12	48.51	11.00	19.00	25.00	7.46	12.30	12.34
C ₁ V ₁	17.97	19.03	36.84	11.67	14.00	20.00	8.26	10.06	10.83
C ₁ V ₂	21.28	21.01	38.25	12.33	14.67	20.33	9.23	11.69	11.58
C ₁ V ₃	20.33	20.10	37.56	12.00	14.33	19.67	8.68	11.39	11.15
C ₁ V ₄	19.78	19.65	37.01	11.33	14.33	19.33	8.57	10.19	11.09
C ₂ V ₁	18.29	19.12	17.70	10.67	14.33	14.00	8.24	10.93	9.68
C ₂ V ₂	20.38	21.39	19.13	12.67	16.00	15.33	8.85	11.78	9.92
C ₂ V ₃	19.85	20.03	18.37	12.00	15.00	14.00	8.34	11.60	9.82
C ₂ V ₄	19.29	19.64	18.33	11.33	14.67	13.67	8.29	11.22	9.87
C ₃ V ₁	18.85	18.85	17.88	11.00	14.00	14.00	7.31	10.49	8.58
C ₃ V ₂	20.06	21.00	19.16	12.00	15.67	15.00	7.86	11.18	9.06
C ₃ V ₃	19.61	19.29	18.13	11.00	15.00	14.33	7.76	10.84	8.82
C ₃ V ₄	19.41	19.26	18.00	10.33	14.67	13.33	7.66	10.80	8.65
SEm±	0.46	0.32	0.37	0.50	0.39	0.42	0.29	0.38	0.39
CD at 5%	1.35	0.93	1.08	NS	1.12	NS	NS	NS	1.12

Table-4 Interaction effect different cutting and varieties on growth of palak.

Treatments	Width of leaves (cm)			leaf petiole length (cm)			Germination (%)
	30DAS	50DAS	70DAS	30DAS	50DAS	70DAS	

C ₀ V ₁	4.32	5.81	6.28	3.71	6.59	6.28	72.71
C ₀ V ₂	4.95	7.06	8.03	5.24	8.19	8.03	72.27
C ₀ V ₃	4.81	6.89	7.22	5.18	7.52	7.22	75.45
C ₀ V ₄	4.72	6.68	6.83	4.56	7.29	6.83	75.41
C ₁ V ₁	5.62	5.23	5.55	6.09	5.92	5.55	78.14
C ₁ V ₂	6.41	6.15	6.82	7.38	6.46	6.82	80.04
C ₁ V ₃	5.67	5.95	6.32	7.06	6.22	6.32	79.04
C ₁ V ₄	5.55	5.72	6.16	6.85	6.03	6.16	78.23
C ₂ V ₁	5.25	5.78	5.31	5.67	6.69	5.31	77.06
C ₂ V ₂	6.25	7.08	5.93	6.66	7.98	5.93	79.31
C ₂ V ₃	5.79	6.66	5.59	6.26	7.22	5.59	78.15
C ₂ V ₄	5.61	6.61	5.54	5.88	6.90	5.54	77.64
C ₃ V ₁	5.47	5.15	4.80	5.74	5.76	4.80	75.53
C ₃ V ₂	5.68	5.98	5.25	6.25	6.56	5.25	78.32
C ₃ V ₃	5.62	5.78	5.05	5.62	6.37	5.05	77.01
C ₃ V ₄	5.51	5.63	4.87	5.57	5.92	4.87	75.64
SEm±	0.28	0.38	0.36	0.29	0.31	0.36	0.35
CD at 5%	NS	NS	NS	NS	NS	NS	1.01




Figure-1. Showed effect of cutting and varieties on plant height and length of leaves.




Figure-2. Showed effect of cutting and varieties on width of leaves and germination (%).

Conclusion:

For the result of experiment it can be concluded that for getting maximum green yield of Palak var. Pusa Jyoti at three leaf cutting is better means V_2C_3 combination is best for production per unit area and per unit time.

References:-

1. Fageria MS, Choudhary, BR, Dhaka RS. Vegetable crops production technology text book vol.2 Kalyani Publication 2022.
2. Singh, D. Vegetable science published by new Vishal publication New Delhi reprint (2019).
3. Gautam A, Ram RB. Effect of organic manures and number of cuttings on growth, yield and quality of Indian spinach. The Pharma Innovation Journal 2022; SP-11(9): 1812-1816.
4. Waseem K, Nadeem MA. Enhancement of spinach production by varying sowing dates, row spacing and frequency of cuttings. Online Journal of Biological Science. 2001; 1(10):902-904.
5. Thakral KK, Singh GR, Pandey UC, Srivastava VK. Effect of nitrogen levels and cutting on the production of green leaves and seed yield of coriander cv. Natural selection. Haryana Agr. Univ. J. Res. 1991; 22:35-9.
6. Singh R, Bassi G, Sing D. Influence of sowing time and leaf cuttings on the seed yield and quality components of palak (*Beta vulgaris var. bengalensis*). Indian Journal of Agricultural Sciences. 2013 Apr 1; 83(4):410-.
7. Bradley GA, Sistrunk WA, Loadenlages MA, Cash JA .Effect of close spacing on spinach yield. *Arkansas farm J.*, 1971, 20(4): 7 (*Hort. Abstr.*, 42: 58-25).
8. Baboo R. Effect of cutting management, nitrogen and phosphorus on growth and yield of fenugreek. *Annals Agric. Res.* 1997: 18(3): 380-382.

9. Bhore MR, Prabhu T, Sajindranath AK. Effect of sowing dates and nitrogen levels on growth, leaf yield and quality of palak. Indian Journal of Horticulture. 2000; 53(1-6):340-4.
10. Gill BS, Randhawa GS, Saini SS. Effect of sowing dates and herb-cutting management on growth and yield of fenugreek (*Trigonella foenum-graecum*). Indian Journal of Agronomy. 2001; 46(2):364-7.
11. Kasture VM, Chauhan PS, Patil VN, Shivankar RS, Rathod RR. Effect of cuttings on seed yield and quality of Indian spinach. Bioved. 2002; 13(1/2):109-11.
12. Lal S, Pandey U C, Pandita M L and Singh K. Effect of nitrogen and cuttings on seed yield of *Beta vulgaris*. Seed Research. 1979; 8(2): 136–40.
13. Singh GP, Meena ML, Prakash J. Effect of different levels of nitrogen and cutting on growth, leaf yield and quality of spinach beet (*Beta vulgaris var. bengalensis*) cv. All Green European J. Biotechnol. BioSci. 2015 Jun; 6:38-42.
14. Sinha SK, Singh JN, Jha BN, Mishra M. Influence of nitrogen and cutting on seed production of Palak Seed Res., 1985; 13 (2) : 73-77.
15. Panse VC, Sukhatme, PV. Statistical methods for agricultural workers. ICAR Publications, New Delhi 1985.
16. Dadiga, A, Kadwey, S. and Parjapati, S. Influences of organic and inorganic source of nutrients on growth, yield attributed traits and yield economic of coriander (*Coriandrum sativum* L.) CVJD-1. J. Agri. Res., 2015;46 (6): 577-80.
17. Dange MB, Bhosale AM, Barkule SR. Effect of organic and inorganic fertilizers on growth, yield and quality of palak (*Beta vulgaris* L.) var. Pusa Jyoti. International Journal of Plant Sciences (Muzaffarnagar). 2011; 6(1):91-4.
18. Naik, DM, Patil SB, Jature and Shinde SJ. Effect of sowing dates number of leaf cutting on growth and yield of palk (*Beta vulgaris* L.). Asian J. Hort., 2010;4 (2): 377-79.
19. Nayak PK, Maji S. Response of nutrient and cutting management for quality and green yield of Palak (*Beta vulgaris var. bengalensis*). Journal of Crop and Weed. 2018; 14(1):126-9.
20. Singh R, Bassi G, Sing D. Influence of sowing time and leaf cuttings on the seed yield and quality components of palak (*Beta vulgaris var. bengalensis*). Indian Journal of Agricultural Sciences. 2013 Apr 1; 83(4):410-.
21. Sarkar RK, Jana JC, Datta S. Effect of cutting frequencies and nitrogen levels on growth, green and seed yield and quality of water spinach (*Ipomoea reptans* Poir.). Journal of Applied and Natural Science. 2014 Dec 1; 6(2):545-51.

22. Singh GP, Meena ML, Prakash J. Effect of different levels of nitrogen and cutting on growth, leaf yield and quality of spinach beet (*Beta vulgaris var. bengalensis*) cv. All Green European J. Biotechnol. BioSci. 2015 Jun; 6:38-42.

UNDER PEER REVIEW