

Evaluation of performance for the quantitative traits in tomato (*Solanum lycopersicon* L.) over seasons

Comment [r1]: The first letter of all words except prepositions must be capitalized.

Abstract

Present investigations were carried out in tomato to assess the mean performance, general mean, and range for nineteen quantitative characters. Forty-five tomato hybrids were generated by crossing ten parents in randomized complete block design with three replications at the Main Experiment Station, Department of Vegetable Science, ANDUA&T, Kumarganj, Ayodhya, during the Rabi season 2020-2021 and 2021-2022. The pooled estimates of mean performance for total fruit yield per plant varied from 2.35 to 3.24 kg for parents and 1.87 to 3.43 kg fruit for hybrids. The mean values over the parents and F₁ hybrids were 2.51 and 2.68 kg, respectively. The highest mean performance for the most desirable traits fruit yield per plant was exhibited by P₆(3.24 kg) followed by P₉ (3.23 kg), P₁₀ (3.09 kg), P₅(2.99 kg), and P₄(2.81 kg) for the parent. Among the hybrids, the highest fruit yield per plant was exhibited by P₂× P₅(3.43 kg) followed by P₄× P₉(3.41 kg), P₄× P₇(3.30 kg), P₁× P₅(3.22 kg) and P₄× P₈(3.20 kg) in descending order.

Comment [r2]: Because this is an international journal, first try to express the equivalent of this season in the form of common seasons (for example, summer, winter, etc.) and then use the word Rabi.

Keywords: Mean performance, mean value, tomato.

Comment [r3]: Do you have any suggestions for scientific community or other beneficiaries?

Introduction -A significant impact of globalization on horticulture has been an increasing demand for quality improvement and the wider adoption of quality standards for fruit, vegetable, and salad commodities. Tomato is used as a vegetable worldwide. It is a very popular and important vegetable in the world. Tomato (*Solanum lycopersicum* L.) is a member of the Solanaceae family. Its chromosome no is $2n=2x=24$. It is grown worldwide under outdoor and indoor conditions. It has become an important commercial crop so far as the area, production, industrial values, and ~~its~~ contribution to human nutrition ~~is~~ are concerned. Tomato is known as protective food because of its special nutritive value and also for its high level of antioxidants. Tomatoes contain many health-promoting compounds and are easily integrated as a nutritious part of a balanced diet (Martí *et al.*, 2016). Tomato is one of the most popular and widely grown vegetable crops of the world next to potatoes. The

Comment [r4]: Lack of references to explain the review literature and the importance of research. Only two references are mentioned for the introduction section.

genus Solanum consists of annual or ~~short-short~~-lived perennial herbaceous, typical ~~day-day~~-neutral plants and ~~warm-warm~~-season crops. Tomato fruit ~~are-is~~ eaten raw or cooked called ~~as~~–“Protective Food” ~~which~~ is being extensively grown as ~~an~~ annual plant all over the world. Tomato in large quantities is used to produce several items like, juice, ketchup, paste, syrup, puree, and drinks, etc. It is rich in beta-carotene, folate, vitamin A, vitamin C, vitamin E, flavonoids, potassium, and other minerals. Overall, tomatoes provide approximately 20 mg of vitamin ‘C’ per 100 grams of edible part. Vitamin C is considered ~~as~~–an excellent antioxidant because it donates electrons ~~for-to~~ enzymes or other compounds that are oxidants. ~~The Red-red~~ colour of ~~the~~ tomato is due to the presence of lycopene. India ranks second in terms of ~~the~~ production of tomatoes after China. In India, the leading ~~tomato-tomato~~-growing states are: Andhra Pradesh, Madhya Pradesh, Karnataka, West Bengal, Maharashtra, U.P., Haryana, Punjab, Gujarat, and Bihar. The total area covered under tomato cultivation is 0.85 Mha with ~~the~~ production of 21.001 MT and its productivity is 25.34 tonnes per ha (NHB database, 2020). The mean performance of genotypes may be used as donor parents in ~~the~~ hybridization programmes for developing ~~high-high~~-yielding varieties of respective groups. Some other genotypes exhibiting very high mean performance for characters other than fruit yield per plant may also be used for transferring these traits. These lines ~~merits~~ due consideration as promising parents for ~~the~~ hybridization programme for bringing overall improvement in plant genetic architecture in a component breeding approach ultimately leading to ~~high-high~~-yielding and ~~high-high~~-quality bottle gourd genotypes even if they have moderate or low fruit yield. Keeping in view the above facts the present investigation was conducted to find out ~~the~~ stable genotype of bottle gourd for improvement in ~~the~~ future.

Materials and Methods

The present research work was conducted during ~~the Rabi~~ seasons of 2020 (Y₁) and 2021 (Y₂) to study the mean performance, general means, and range for nineteen characters using diallel mating design at the Main Experiment Station (MES) of the Department of Vegetable Science, Acharya Narendra Deva University of Agriculture and Technology, Narendra Nagar, Kumarganj, Ayodhya (U.P.) India. The

Comment [r5]: How and with what statistical test was the means compared?

Comment [r6]: Because this is an international journal, first try to express the equivalent of this season in the form of common seasons (for example, summer, winter, etc.) and then use the word Rabi.

observations were recorded on nineteen characters. The experimental materials for the present investigation comprised ~~of~~ nineteen promising and diverse inbred lines/varieties of tomato selected ~~on the basis of~~ based on genetic variability from the germplasm stock maintained in the Department of Vegetable Science, Acharya Narendra Deva University of Agriculture and Technology, Kumarganj, Ayodhya (U.P.) India. The selected parental lines *i.e.*; 2013/TODVAR-2, NDT-2, NDT-Sel-1, NDT-5, NDT-7, NDT-4, NDT-P-1, NDT-6, NDT-6-2-1, NDT-Sel-2. to get 45 F₁ seed. Each hybrids and parents were grown in rows spaced ~~at~~ 0.60 meters apart with a ~~plant~~ ~~plant-to-to~~ plant spacing of 0.50 meter. All the recommended agronomic packages of practices and protection measures were followed to raise good crops. Observations were recorded on days to 50% flowering, days to first fruit harvest, plant height (cm), number of primary branches per plant, number of fruits per cluster, number of fruits per plant, average fruit weight (g), pericarp thickness (mm), number of locules per fruit, polar diameter (cm), ~~equitorial-equatorial~~ equatorial diameter (cm), marketable fruit yield per plant (kg), total soluble solids (TSS), titrable acidity (%), ascorbic acid content (mg/100g) Reducing sugar (mg/100g). none - reducing sugar (mg/100g). total sugar (mg/100g) and total fruit yield per plant (kg).

Result and Discussion

Mean performance, general mean, range, coefficient of variation, critical difference, and standard error for nineteen characters 45 F₁'s and their 10 parents (Y₁=2020 and Y₂=2021) and pooled had been presented in Table-1. The highest mean performance for ~~the~~ most desirable traits fruit yield per plant was exhibited by P₄(3.01 kg) followed by P₉(2.84 kg), P₇(2.83 kg), P₃(2.80) and P₁₀(2.62 kg) in Y₁, P₄(3.45 kg) exhibited highest fruit yield per plant followed by P₃(2.97 kg), P₉(2.89 kg), P₁(2.86) and P₇(2.83 kg) in Y₂ and P₆(3.24 kg) exhibited highest fruit yield per plant followed by P₉(3.23 kg), P₁₀(3.09 kg), P₅(2.99 kg) and P₄(2.81 kg) in pooled.

The ~~above~~ above-mentioned genotypes may be used as donor parents in ~~the~~ hybridization programmes for developing ~~high-high~~ high-yielding varieties of respective groups. Some other genotypes exhibiting very high mean performance for characters other than fruit yield per plant are also listed in Table-1. These lines merits due consideration as promising parents for ~~the~~ hybridization programme for bringing

Comment [r7]: Regarding the obtained results, the discussion has not been done completely. Each result should be discussed separately and also your expert opinions should be stated for the reason of reaching such observations and then compared with previous studies.

overall improvement in plant genetic architecture in a component breeding approach ultimately leading to **high-high**-yielding and **high-high**-quality tomato genotypes even if they have moderate or low fruit yield. In this context, the most desirable parents were P₆(31.66 days) which showed earliness for days to fifty present flowering among the parents which was followed by P₁₀ (34.66 days), P₁(35.00 days), P₅(38.66 days) and P₃(39.00 days) in Y₁, Parent P₆(31.33 days) followed by P₄(34.00 days), P₁₀(35.78 days) and P₁(36.12days) in Y₂ and Parent P₆(31.11 days) followed by P₅(31.77 days), P₄(32.00 days), and P₃(33.60 days) in pooled. Among the parents days to first fruit harvest ~~was~~ were observed in P₆(83.00 days) followed by P₅(84.66 days), P₇(87.33 days) and P₄(89.66 days) in Y₁, P₆(80.00 days) followed by P₅ (82.67), P₇(89.67 days) and P₉(90.62 days) in Y₂ and P₆(78.98days) followed by and P₇(79.66 days), P₈(82.06days) and P₂(83.32days) in pooled; highest plant height (cm) was recorded in parent P₁₀(119.58cm) followed by P₃(114.84 cm), P₇(113.73cm), P₃(114.84 cm) and P₇(113.73cm) in Y₁, P₇(120.65 cm) followed by P₃(115.67 cm), P₆(113.87cm), P₁₀(111.12cm) and P₁₀(110.00 cm) in Y₂ and P₇(120.09 cm) followed by P₆ (116.80cm), P₃(114.35 cm), P₄(114.29 cm) and P₁₀(112.77cm) in pooled.

highest primary branches per plant among the parents was recorded in P₅ (6.36 branches) followed by P₇(6.15 branches), P₂(5.36 branches), P₉(5.28branches) ~~And~~ and P₆(5.22branches) in Y₁, P₅(6.98branches) branches) followed by P₇(6.89 branches), P₂(6.67 branches) P₉(6.33 branches) in Y₂ and P₅(6.67branches) followed by P₄(6.60 branches), P₃(6.58branches), and P₆(6.45 branches) in pooled; maximum number of fruit per cluster among the parents was observed in P₃(5.16) followed by (4.45), P₁₀(4.19), P₄(3.97), and P₂(3.81) in Y₁, P₃(5.67) followed by P₅(5.00), P₁(4.69), P₁₀(4.32) and P₄ (4.12) in Y₂ and P₅ (4.90) followed by P₃ (4.56), P₆ (4.34) and P₉(4.20 m) in pooled; maximum number of fruit per plant was found in P₉(35.30), followed by P₄(33.62), P₆(31.26), P₁₀(30.33), and P₅(28.62) in Y₁, P₉(36.34) followed by P₄(35.89), P₃(33.87), P₇(32.78) and P₆(32.67) in Y₂ and P₆(36.52) was found for maximum number of fruit per plant among the parents which was followed by P₉(36.28), P₃(33.68), P₄(32.63) and P₁₀(30.03) in pooled; maximum average fruit weight was observed in parent P₁(79.02g) followed by (76.13g), P₄(75.01g), P₆(73.16g) and P₁₀(71.54 g) in Y₁, P₁(85.40g) followed by P₉(81.67g), P₃ (78.86g), P₄(75.45g)

and P₆(74.76g) in Y₂ and P₆(78.41g) followed by P₃(77.89 g), P₁(76.09g), P₉(76.75g) and P₁₀(73.04g) in pooled; parental line P₉(5.25mm) was found for maximum pericarp thickness followed by P₄ (5.09 mm), P₅(4.72 mm), P₆(4.67 mm) and P₁(4.63mm) in Y₁, P₄(5.67 mm) followed by P₉(5.66 mm), P₈(4.83mm), P₁(4.78mm) and P₆(4.77 mm) in Y₂ and P₄(5.49 mm) exhibited maximum pericarp thickness among the parents which was followed by P₉(4.93 mm), P₁₀(4.96 mm), P₉(4.93mm) and P₆(4.72 mm) in pooled.

Comment [r8]: Please correct the format for all similar cases

Maximum number of locules per fruit among the parents was observed in P₇(6.19) followed by P₄(6.06), P₂ (5.43), P₁₀(5.05) and P₆(4.51) in Y₁, P₇(6.34) followed by P₆(6.14), P₁₀(5.67), P₄(4.97) and P₉ (4.67) in Y₂ and P₄ (5.74) followed by P₂ (5.42), P₃ (5.19) and P₁₀(5.01) in pooled; maximum polar diameter of fruit (cm) was recorded in P₈(6.34) followed by P₃(6.31), P₅ (5.55), P₆ (5.45) and P₂(5.25), Y₁, P₅(6.39) followed by P₈(6.79), P₉(5.87), P₂(5.75) and P₄(4.96) in Y₂ and P₉ (6.49) followed by P₆ (6.16), P₃ (6.00), P₁(5.14) and P₇ (5.08) in pooled; maximum Equatorial diameter (cm) of fruit was recorded in P₅(9.48cm) followed by P₈(9.47cm), P₇ (8.35cm), P₂ (7.96cm) and P₉(7.70cm), in Y₁, P₇(9.76cm) followed by P₈(9.49 cm), P₅(8.09 cm), P₆(8.45cm) and P₂ (7.99 cm) in Y₂ and P₅ (9.17cm) followed by P₆ (8.97cm), P₇ (8.66cm), P₉(8.28cm) and P₁ (7.83 cm) in pooled; maximum marketable fruit yield per plant (kg) was recorded in P₃(2.53 kg) followed by P₁(2.51kg), P₉ (2.43kg), P₃(2.61kg) and P₁ (2.34kg) in Y₁, P₄(2.88 kg) followed by P₃(2.877 kg), P₇(2.78 kg), P₁ (2.73kg) and P₁₀(2.70kg) in Y₂ and P₉ (2.97kg) followed by P₆ (2.91kg), P₁₀ (2.86 kg), P₁(2.34kg) and P₂ (2.27kg) in pooled; parents P₁(5.68) contain highest total soluble solids followed by P₈(5.62°B), P₅ (5.47°B), P₇(5.01°B) and P₆ (4.93°B), in Y₁, P₁(5.70°B) followed by P₆(5.33°B), P₉(5.47°B), P₁ (5.31°B) and P₈(5.24°B) in Y₂ and P₇ (5.49°B) followed by P₆ (5.33°B), P₁ (5.31), P₁₀ (5.10°B) and P₄(5.09°B) in pooled;

maximum titrable acidity was recorded in P₁(0.46%) followed by P₇(0.45%), P₆ (0.40%), P₈(0.43%) and P₁₀ (0.37%) in Y₁, P₁(0.43%) followed by P₇(0.42%), P₄(0.38%), P₃(0.37%) and P₂ (0.35%) in Y₂ and P₁ (0.43%) followed by P₂(0.41%), P₇ (0.40%), P₄(0.36%) and P₈ (0.37%) in pooled; parents P₃(24.57mg/100g) contain highest Ascorbic acid content followed by P₇(23.54

mg/100g), P₆ (22.10mg/100g) , P₂(21.77mg/100g) and P₁₀ (21.38mg/100g)), in Y₁P₃ (24.59mg/100g) followed by P₈(23.14 mg/100g), P₇(22.07mg/100g) , P₁(21.53mg/100g) and P₂(21.85mg/100g)) in Y₂and P₃ (24.06 mg/100g) followed by P₇ (22.07mg/100g), P₉(21.73mg/100g) , P₁ (21.53mg/100g)) and P₄ (21.34mg/100g)in pooled; maximum reducing sugar was recorded in P₇(1.66%) followed by P₉(1.39%), P₁₀ (1.61 %) , P₅(1.54%) and in Y₁, P₉(1.76%) followed by P₇(1.65%), P₆(1.50%) , P₃(1.54%) and P₁ (1.43%) in Y₂and P₂ (01.57%) followed by P₁₀(1.51%) , P₅ (1.47%), P₇ (1.44%)) andP₈ (1.43%) in pooled; maximum non-reducing sugar(mg/100g) was recorded in P₉(2.78%) followed by p₄(2.77%), p₁ (2.68%) , P₂(2.67%) andP₁₀(2.60%) in Y₁, P₈(2.94%) followed by P₁(2.76%) , P₁₀(2.75%), P₂(2.68%) and P₅ (2.64%) in Y₂and P₈ (2.80%) followed by P₂(2.75%),P₄(2.71%),P₉(2.56%)andP₁₀ (2.54%) in pooled; maximum total sugar(mg/100g) was recorded inP₁₀(4.21%) followed byP₉(4.16%), P₅ (4.07%) ,P₈(4.05%) and P₂(3.98%) in Y₁, P₈(4.59%) followed by P₁₀(4.51%) , P₅ (4.39%) , P₅(4.07%) and P₆(3.88 %) in Y₂and P₂ (4.33%) followed by P₈(4.06%) , P₁₀ (4.02%), P₉ (3.83%) and P₆ (3.80%)) in pooled.Among the hybrids, the highest fruit yield per plant was exhibited by P₂× P₄(3.94 kg) followed by p₆× p₉(3.67 kg), p₅× P₁₀(3.59 kg), P₄× P₈(3.56 kg) and P₉× P₁₀(5.13 kg) in Y₁and P₅× P₁₀(4.23 kg) followed by P₂× P₄(4.12 kg), P₁× P₄(3.89 kg), P₄× P₈(3.93 kg) and P₄× P₁₀(3.76 kg) in y₂,P₂× P₅(3.43 kg) followed by P₄× P₉(3.41 kg), P₇× P₉(3.30 kg),P₁× P₅(3.22 kg) and P₄× P₄× P₈(3.20 kg) in pooled descending order.

Thus there ~~was~~ were significant differences for all the traits over seasons and pooled among the parents and F₁. This might be due to the influence of the environment. Similar observations in tomatoes es were also reported by Joshi & Kohli (2005), Jogi *et al.* (2008), Mohammed *et al.* (2012), Narolia *et al.* (2012), kerketta *et al.* (2018) Prakash *et al.* (2019) and Anuradha *et al.* (2021) in tomato.

Table 1: Mean performance, general mean, range, coefficient of variation, critical difference, and standard error for nineteen characters of diallel set of 45 F₁'s and their 10 parents (Y₁=2020 -2021 and Y₂=2021-2022), and pooled.

Table 1: Mean performance, general mean, range, coefficient of variation, critical difference and standard error for nineteen characters of a diallel set of 45 F₁'s and their 10 parents (Y₁=2020 -2021and Y₂=2021-2022) and pooled.

Genotypes	Days to 50% flowering			Days to first fruit harvest			Plant height (cm)			Primary branches per plant		
	Y ₁	Y ₂	Pooled	Y ₁	Y ₂	Pooled	Y ₁	Y ₂	Pooled	Y ₁	Y ₂	Pooled
P ₁	35.00	36.12	33.33	90.67	91.07	86.83	81.13	82.77	95.03	4.74	5.56	4.98
P ₂	41.33	42.00	37.50	93.00	94.23	83.32	94.78	95.78	105.27	5.36	6.67	6.12
P ₃	39.33	40.23	33.60	89.00	90.00	86.66	114.84	115.67	114.35	6.04	7.07	6.58
P ₄	33.00	34.00	32.00	89.66	89.13	83.83	108.03	110.03	114.29	5.23	5.44	6.49
P ₅	38.66	40.44	31.77	84.66	82.67	83.10	73.09	74.87	96.76	6.36	6.98	6.67
P ₆	31.66	31.36	31.11	83.66	80.54	78.98	108.93	113.87	116.80	5.22	6.79	6.45
P ₇	40.33	41.56	41.16	87.33	89.01	79.66	113.73	120.65	120.05	6.15	6.89	5.91
P ₈	38.00	39.00	40.39	87.33	89.00	82.06	91.56	90.77	93.50	4.97	5.89	4.94
P ₉	40.33	41.76	41.61	89.00	90.62	87.98	86.87	90.53	102.21	5.28	6.33	5.92
P ₁₀	34.66	35.78	38.77	90.33	92.67	102.21	119.5	111.67	112.77	4.78	5.69	5.48
P ₁ ×P ₂	32.00	34.00	34.06	84.00	85.90	87.53	86.01	87.67	84.39	5.10	6.90	5.33
P ₁ ×P ₃	31.00	32.56	32.50	72.00	70.67	78.95	95.97	96.79	91.81	5.60	6.75	6.24
P ₁ ×P ₄	29.33	31.78	30.94	81.00	78.78	75.83	104.37	110.78	100.58	4.38	5.98	5.56
P ₁ ×P ₅	35.33	36.56	33.55	84.67	85.89	81.72	100.34	103.85	105.55	6.45	7.65	6.21
P ₁ ×P ₆	33.33	35.00	34.94	75.33	76.56	80.61	114.74	117.89	109.29	5.12	5.56	6.38
P ₁ ×P ₇	30.00	28.56	32.50	87.33	88.45	81.94	107.75	108.89	112.82	4.27	5.87	4.91
P ₁ ×P ₈	33.33	34.67	30.94	75.33	78.45	81.89	84.90	82.90	96.89	5.58	6.70	5.72
P ₁ ×P ₉	28.00	26.21	31.33	87.00	87.85	82.72	91.64	90.67	87.26	6.48	7.45	6.59
P ₁ ×P ₁₀	31.33	33.67	28.77	71.67	73.65	79.75	114.42	115.76	102.54	5.28	6.89	6.36
P ₂ ×P ₃	30.33	31.97	36.16	88.00	98.45	91.11	115.06	116.76	105.42	6.07	6.66	6.37
P ₂ ×P ₄	30.00	31.89	30.98	83.33	76.98	90.89	104.35	108.56	110.55	7.12	8.56	6.89
P ₂ ×P ₅	30.00	32.87	30.94	74.00	75.67	75.49	93.33	93.98	100.94	5.00	6.07	6.78
P ₂ ×P ₆	26.33	24.87	29.60	72.00	68.11	73.83	109.77	110.67	101.87	5.13	5.87	5.60
P ₂ ×P ₇	25.00	25.78	24.93	74.33	75.78	71.22	111.04	112.56	110.85	6.22	6.54	6.04
P ₂ ×P ₈	34.67	35.00	30.22	86.00	80.67	80.89	86.35	85.98	99.45	5.96	6.32	6.25
P ₂ ×P ₉	30.67	31.78	32.83	90.00	93.89	85.33	91.27	92.76	88.62	7.30	7.86	6.81
P ₂ ×P ₁₀	26.00	27.87	28.89	82.67	84.34	88.27	112.08	113.86	102.42	6.40	7.12	7.13

P₃×P₄	24.00	22.89	32.11	81.67	83.89	85.83	117.30	120.67	116.48	5.63	6.32	6.35
P₃×P₅	28.00	27.13	25.44	85.33	79.67	84.61	118.52	125.78	119.59	6.27	7.59	6.29
P₃×P₆	31.67	32.97	29.39	85.33	81.87	82.50	117.78	121.65	121.77	5.41	5.98	6.50
P₃×P₇	32.00	33.67	32.48	84.67	80.54	83.26	129.30	130.45	125.47	5.56	5.43	5.77
P₃×P₈	31.33	32.00	32.50	87.33	84.56	83.93	123.20	128.67	126.82	7.12	8.56	6.27
P₃×P₉	27.00	28.57	29.50	86.67	79.78	85.61	106.25	108.45	117.45	5.55	6.57	7.05
P₃×P₁₀	30.00	31.00	29.28	75.00	78.01	77.39	124.71	120.56	116.57	6.63	7.76	6.60
P₄×P₅	26.33	28.78	30.16	85.33	86.98	87.23	116.00	117.56	113.00	4.67	6.56	5.06
P₄×P₆	25.33	26.89	27.05	72.33	72.89	79.65	118.08	119.56	117.82	4.61	4.99	5.58
P₄×P₇	28.00	29.00	27.44	88.00	89.45	80.44	116.30	117.56	117.93	4.74	5.87	4.86
P₄×P₈	41.00	42.76	35.00	84.67	84.78	87.05	105.78	108.53	111.66	5.15	7.56	5.51
P₄×P₉	37.67	38.56	40.21	90.33	91.52	87.55	90.20	95.45	99.36	5.42	7.67	6.48
P₄×P₁₀	27.33	24.89	32.94	86.00	81.54	88.76	118.86	120.45	107.15	4.48	6.98	6.07
P₅×P₆	31.67	32.21	36.05	91.00	92.56	86.83	115.44	117.56	95.15	4.51	4.99	5.75
P₅×P₇	28.00	29.56	30.10	88.00	89.56	90.28	105.39	105.96	111.47	4.34	4.98	4.66
P₅×P₈	40.67	42.67	35.11	86.67	87.46	88.11	84.64	90.34	95.30	4.48	4.43	4.73
P₅×P₉	29.00	32.56	35.83	87.00	88.76	87.22	89.08	90.67	89.70	7.26	7.67	5.84
P₅×P₁₀	32.67	30.56	32.61	73.33	74.97	81.04	118.19	124.67	104.42	6.71	7.69	7.18
P₆×P₇	33.00	33.67	32.16	73.00	68.78	76.77	117.45	118.34	115.66	5.60	6.98	6.19
P₆×P₈	37.00	37.89	35.33	94.33	95.78	81.55	114.12	115.47	116.22	5.97	6.97	6.47
P₆×P₉	36.67	37.35	37.27	81.00	82.67	88.39	83.75	84.45	99.61	6.54	7.98	6.75
P₆×P₁₀	41.00	42.00	39.17	71.67	72.00	77.16	123.60	126.45	104.02	4.52	5.67	6.25
P₇×P₈	40.33	25.90	34.44	88.67	89.47	88.84	109.41	117.56	115.03	5.56	6.98	6.22
P₇×P₉	29.00	30.67	27.45	72.33	73.67	80.90	111.67	108.53	114.61	6.26	7.56	6.62
P₇×P₁₀	41.00	42.78	35.83	75.33	76.79	74.50	116.82	95.45	112.67	5.62	4.89	6.59
P₈×P₉	39.33	40.14	39.16	87.00	88.62	88.00	85.19	120.45	87.98	6.48	6.50	6.18
P₈×P₁₀	41.33	42.89	40.73	91.66	86.98	90.14	114.77	117.56	117.61	5.96	6.56	6.23
P₉×P₁₀	41.00	42.89	41.38	87.00	88.56	88.81	110.03	105.96	100.28	6.16	6.18	6.25
Mean	32.86	33.82	33.30	83.51	83.66	83.49	105.77	107.85	106.78	5.61	6.58	6.09
C.V.	7.73	6.49	12.65	6.59	5.47	8.41	6.55	5.12	10.50	11.54	8.76	14.43

S.E.±M	1.46	1.27	1.72	3.17	2.64	2.86	4.01	3.19	4.58	0.37	0.33	0.35	
C.D. 5%	4.10	3.55	4.79	8.91	7.41	7.98	11.23	8.93	12.753	1.05	0.93	0.99	
Range	Lowest	24.00	22.89	24.93	71.66	68.11	71.22	74.87	74.87	84.39	4.43	4.43	4.66
	highest	41.33	42.89	41.61	94.33	98.45	91.11	130.45	130.45	126.82	8.56	8.55	7.18

Genotypes	Fruit per cluster			No: of fruit per plant			Average fruit weight (g)			Pericarp thickness (mm)		
	Y ₁	Y ₂	Pooled	Y ₁	Y ₂	Pooled	Y ₁	Y ₂	Pooled	Y ₁	Y ₂	Pooled
P₁	4.55	4.69	4.23	26.44	27.78	28.85	79.02	85.40	76.09	4.63	4.78	4.65
P₂	3.81	3.87	3.91	24.81	26.56	27.39	68.25	70.67	71.01	3.78	3.86	3.82
P₃	5.16	5.67	4.56	32.59	33.87	33.68	76.13	78.86	77.89	4.59	4.87	4.58
P₄	3.97	4.12	3.48	33.62	35.89	32.63	75.01	75.45	71.28	5.09	5.67	5.49
P₅	4.13	5.03	4.90	28.59	29.98	32.02	60.58	64.35	68.67	4.72	4.32	4.23
P₆	3.91	4.12	4.34	31.26	32.77	36.52	73.16	74.67	78.41	4.67	4.77	4.72
P₇	3.88	3.88	3.80	20.83	32.78	27.25	64.93	68.90	74.90	3.81	3.97	4.28
P₈	3.91	3.91	3.75	27.37	28.98	29.02	78.54	79.33	76.75	4.81	4.83	4.31
P₉	3.53	3.66	4.20	35.03	38.34	36.28	67.36	81.67	70.03	5.25	5.67	4.93
P₁₀	4.19	4.32	3.93	30.33	22.86	30.00	71.54	74.87	73.04	4.58	4.45	4.96
P₁×P₂	4.52	4.78	4.61	34.04	37.45	30.91	64.84	69.97	75.12	4.44	4.78	4.60
P₁×P₃	3.39	3.67	4.08	27.23	31.78	32.34	66.27	71.97	68.11	4.64	4.83	4.71
P₁×P₄	3.71	4.72	3.68	26.78	34.23	29.28	72.85	77.34	72.41	4.53	4.59	4.68
P₁×P₅	4.29	4.29	4.50	25.29	29.00	29.75	82.74	85.76	80.03	4.80	4.98	4.69
P₁×P₆	4.27	4.76	4.27	25.60	27.89	27.30	81.33	84.87	83.54	5.27	5.55	5.12
P₁×P₇	4.67	4.65	4.71	35.79	32.89	31.83	75.01	76.54	79.93	4.85	4.87	5.20
P₁×P₈	3.69	3.98	4.17	30.47	35.89	31.68	78.33	82.76	77.43	3.63	4.12	4.24
P₁×P₉	4.54	4.87	4.26	31.78	33.87	33.83	82.46	85.76	82.61	4.58	4.56	4.35
P₁×P₁₀	3.89	3.99	4.38	26.93	29.98	30.40	72.83	73.78	79.29	3.68	3.87	4.12
P₂×P₃	4.59	4.76	4.23	25.49	28.12	26.02	82.49	80.56	76.58	4.75	4.97	4.30
P₂×P₄	3.57	4.56	4.16	29.49	33.45	28.80	75.90	80.56	78.23	4.67	4.77	4.82
P₂×P₅	3.60	3.60	4.08	26.91	28.65	30.18	77.47	79.54	79.01	4.67	4.65	4.72

P₂×P₆	3.74	3.86	3.67	26.59	29.56	27.61	76.24	76.32	77.89	3.83	4.56	4.23
P₂×P₇	3.62	3.56	3.73	32.93	27.45	31.24	69.87	70.54	73.09	4.53	4.68	4.54
P₂×P₈	3.37	3.65	3.46	28.89	30.34	28.17	83.60	87.56	77.06	4.66	4.88	4.67
P₂×P₉	4.63	4.76	4.13	29.52	31.56	29.93	71.49	76.43	79.52	4.83	4.97	4.85
P₂×P₁₀	3.96	3.97	4.36	33.52	34.78	32.54	76.85	79.65	76.64	4.48	4.58	4.72
P₃×P₄	4.49	4.50	5.08	32.37	33.78	33.12	66.17	69.54	72.51	4.65	4.65	4.76
P₃×P₅	3.45	4.87	3.97	33.42	39.67	33.60	65.56	71.87	67.55	3.83	3.95	4.24
P₃×P₆	3.70	4.12	4.28	31.23	34.67	35.45	63.52	66.56	67.69	4.67	4.78	4.31
P₃×P₇	3.67	3.65	3.89	27.93	29.56	31.29	67.52	70.67	67.03	4.83	4.83	4.80
P₃×P₈	3.86	4.65	3.75	31.03	34.67	30.29	83.33	85.45	77.00	5.46	5.76	5.14
P₃×P₉	4.52	4.57	4.58	32.90	35.54	33.78	82.83	83.43	84.14	5.60	5.87	5.68
P₃×P₁₀	2.67	2.99	3.62	28.49	31.65	32.01	64.85	67.56	74.14	4.45	5.89	5.16
P₄×P₅	3.56	3.87	3.84	27.51	29.67	31.70	79.36	83.32	77.40	4.40	4.56	5.03
P₄×P₆	3.72	3.79	3.79	35.46	37.45	32.56	74.26	75.78	78.78	4.82	4.34	4.68
P₄×P₇	3.60	3.87	3.69	28.15	30.56	32.80	75.02	77.56	75.40	3.68	3.35	4.01
P₄×P₈	3.64	4.34	3.75	29.28	37.56	29.92	74.90	79.76	76.23	4.53	4.42	3.93
P₄×P₉	3.63	4.53	3.98	26.18	31.87	31.87	76.04	78.56	77.90	4.50	4.32	4.46
P₄×P₁₀	4.59	5.67	4.56	30.97	35.45	31.41	73.92	76.76	76.24	3.59	3.76	3.95
P₅×P₆	4.14	4.87	4.57	25.23	29.56	27.60	71.98	73.89	68.16	4.90	4.83	4.61
P₅×P₇	4.65	4.48	4.76	30.60	32.45	30.07	71.30	71.98	72.59	5.13	5.65	4.97
P₅×P₈	5.29	5.23	4.88	26.74	30.56	29.59	75.00	76.67	73.49	3.85	3.92	4.75
P₅×P₉	4.74	4.78	4.98	31.89	27.56	31.22	80.92	84.65	78.79	4.68	4.76	4.29
P₅×P₁₀	4.71	4.78	4.74	34.45	41.78	31.00	78.04	83.67	81.34	4.66	4.78	4.71
P₆×P₇	3.78	3.97	3.94	31.41	34.67	32.09	78.90	80.08	76.78	5.16	5.69	4.96
P₆×P₈	4.25	4.35	4.11	33.49	35.67	34.07	82.81	84.78	81.44	4.77	4.81	5.23
P₆×P₉	3.67	3.67	4.01	34.34	36.56	35.00	73.14	74.84	78.96	3.77	3.89	4.28
P₆×P₁₀	3.65	3.73	3.66	31.60	33.67	34.07	80.73	84.87	77.78	4.59	4.75	4.24
P₇×P₈	4.61	4.65	4.25	32.93	30.56	32.85	72.70	75.87	70.84	4.93	4.94	4.45
P₇×P₉	3.44	4.21	4.04	33.20	33.67	31.88	70.96	73.98	73.41	4.76	4.87	4.85
P₇×P₁₀	3.59	3.60	3.90	29.67	30.67	31.68	71.90	74.98	72.94	3.70	3.83	4.28
P₈×P₉	4.37	4.67	4.14	30.48	30.56	29.73	73.65	77.85	76.49	4.67	4.78	4.75

$P_8 \times P_{10}$	4.40	4.87	4.5367	32.86	37.56	31.71	68.35	72.65	73.10	4.63	4.62	4.70	
$P_9 \times P_{10}$	3.96	3.67	3.8133	27.36	29.67	32.85	75.90	74.56	78.63	5.15	5.35	5.41	
Mean	4.02	4.30	4.1613	29.98	32.44	31.28	74.15	77.21	75.66	4.56	4.72	4.64	
C.V.	8.51	7.48	11.75	7.11	6.20	10.32	6.42	5.64	7.89	7.33	8.39	10.74	
S.E.±M	0.19	0.19	0.199	1.23	1.16	1.31	2.75	2.51	2.43	0.19	0.23	0.20	
C.D. 5%	0.55	0.52	0.55	3.45	3.25	3.67	7.71	7.04	6.78	0.54	0.64	0.56	
Range	Lowest	2.99	2.99	3.46	22.87	22.86	26.02	64.35	64.35	67.03	3.35	3.35	3.82
	highest	5.67	5.67	5.08	41.78	41.78	36.52	87.56	87.56	84.14	5.89	5.88	5.68

Genotypes	Locules per fruit			Polar diameter of fruit (cm)			Equatorial diameter of fruit (cm)			Marketable fruit yields per plant (kg)		
	Y ₁	Y ₂	Pooled	Y ₁	Y ₂	Pooled	Y ₁	Y ₂	Pooled	Y ₁	Y ₂	Pooled
P₁	4.63	4.64	4.57	4.83	4.96	5.14	7.57	7.76	7.83	2.53	2.73	2.34
P₂	5.43	5.43	5.15	5.25	5.75	4.97	7.96	7.99	7.70	1.88	2.35	2.27
P₃	3.74	3.77	5.19	6.31	5.48	6.0	6.44	6.67	6.94	2.53	2.87	2.61
P₄	6.06	4.97	5.74	4.80	4.82	4.47	6.47	6.70	6.71	2.08	2.88	2.25
P₅	4.54	4.67	4.60	5.55	6.38	4.71	9.48	8.09	9.17	1.65	2.34	2.47
P₆	4.51	6.14	4.69	5.45	5.55	6.16	8.06	8.45	8.97	2.16	2.29	2.91
P₇	6.19	6.34	5.42	4.40	4.48	5.08	8.35	9.76	8.66	2.41	2.78	2.30
P₈	2.98	3.34	3.91	6.34	6.78	5.61	9.48	9.49	8.67	1.65	1.30	2.15
P₉	4.60	4.67	4.19	5.51	5.87	6.49	7.7	7.78	8.28	2.43	2.67	2.97
P₁₀	5.05	5.67	5.01	3.99	4.23	5.22	5.40	5.36	6.58	2.24	2.70	2.86
P₁×P₂	4.50	4.70	4.57	4.67	4.84	4.81	7.33	7.87	7.54	2.15	2.40	2.44
P₁×P₃	4.75	4.73	4.72	5.64	5.70	5.23	7.46	7.46	7.66	2.13	2.43	2.26
P₁×P₄	4.67	4.87	4.70	6.31	6.34	6.00	8.31	8.45	7.88	2.14	3.70	2.28
P₁×P₅	6.67	6.67	5.76	5.25	4.89	5.79	9.43	8.90	8.94	1.46	1.89	2.58
P₁×P₆	4.60	4.98	5.63	4.86	4.86	4.87	6.71	6.87	7.80	1.16	1.65	1.52
P₁×P₇	5.05	5.45	5.01	5.81	5.87	5.33	8.12	8.78	7.49	2.86	2.21	2.25
P₁×P₈	5.68	5.32	5.56	6.48	6.48	6.17	7.08	7.87	7.93	1.62	2.45	1.91

P₁×P₉	4.66	4.64	4.99	5.75	5.83	6.11	6.78	6.65	7.32	2.56	2.77	2.50
P₁×P₁₀	4.66	4.87	4.65	4.63	4.70	5.23	7.79	7.45	7.22	1.94	2.67	2.35
P₂×P₃	4.56	4.76	4.99	5.49	5.75	5.62	7.31	7.76	7.64	1.54	1.87	1.94
P₂×P₄	3.75	3.87	4.25	3.86	3.97	4.80	6.85	6.97	7.30	2.36	3.89	2.11
P₂×P₅	4.64	4.98	4.25	6.41	6.48	5.18	8.19	8.23	7.57	1.55	2.45	2.72
P₂×P₆	4.71	4.81	4.84	5.71	5.77	6.09	7.57	7.23	7.89	2.52	3.34	2.48
P₂×P₇	5.42	5.67	5.11	5.74	5.34	5.75	7.50	7.55	7.36	2.57	2.21	2.95
P₂×P₈	4.74	4.89	5.20	4.94	5.12	5.13	6.53	6.54	7.04	1.98	2.15	2.09
P₂×P₉	4.52	4.61	4.70	6.66	6.69	5.88	7.52	7.65	7.03	2.69	2.70	2.42
P₂×P₁₀	6.52	6.65	5.56	5.64	5.70	6.16	7.63	7.45	7.63	2.51	2.70	2.60
P₃×P₄	4.55	4.87	4.16	5.58	5.87	5.52	8.84	8.86	7.75	1.86	2.43	2.36
P₃×P₅	4.62	4.32	4.74	4.53	4.59	5.20	8.50	8.87	8.68	2.82	2.90	2.62
P₃×P₆	4.47	4.75	4.39	5.60	5.68	5.09	8.59	8.72	8.73	2.29	2.21	2.59
P₃×P₇	3.49	3.54	4.12	5.54	5.97	5.61	7.60	7.70	8.16	1.43	1.56	1.82
P₃×P₈	4.57	4.21	4.05	4.90	4.97	5.43	7.56	7.78	7.63	2.87	3.43	2.21
P₃×P₉	4.57	4.65	4.39	5.67	5.78	5.31	7.50	7.78	7.64	2.43	2.56	2.92
P₃×P₁₀	5.29	5.43	4.97	3.74	4.14	4.76	6.60	6.95	7.19	1.96	2.43	2.26
P₄×P₅	3.51	3.64	4.24	5.58	5.72	5.20	9.43	9.49	8.06	2.67	2.50	2.77
P₄×P₆	3.80	3.89	3.71	6.77	6.89	6.24	9.44	9.34	9.46	2.93	3.56	2.71
P₄×P₇	4.67	4.84	4.28	5.79	5.46	6.33	9.38	9.45	9.35	2.76	2.76	3.16
P₄×P₈	4.41	4.53	4.62	4.72	4.96	5.08	8.51	8.67	8.98	2.93	4.24	2.84
P₄×P₉	5.34	5.75	4.93	5.44	5.44	5.19	9.01	9.34	8.83	2.84	3.12	3.53
P₄×P₁₀	4.61	4.67	5.18	3.56	3.87	4.50	8.68	8.87	9.01	2.95	3.30	3.03
P₅×P₆	3.58	3.76	4.12	4.56	4.64	5.47	8.38	8.56	8.23	1.48	2.31	1.91
P₅×P₇	5.52	5.34	4.64	5.50	5.89	5.06	8.96	8.45	8.76	1.48	1.48	1.89
P₅×P₈	3.72	3.56	4.52	4.53	4.58	5.21	6.38	6.79	7.41	2.21	2.41	1.84
P₅×P₉	4.70	4.79	4.13	5.64	5.87	5.10	8.62	8.87	7.70	1.80	1.80	2.10
P₅×P₁₀	4.74	4.87	4.76	6.26	6.87	6.06	9.66	9.87	9.26	2.96	3.65	2.38
P₆×P₇	6.34	6.38	6.24	5.64	5.85	5.59	8.18	8.97	8.31	2.60	2.70	2.44
P₆×P₈	4.44	4.65	5.41	4.51	4.98	5.18	7.54	7.87	8.25	1.78	2.30	2.23
P₆×P₉	5.31	5.75	4.97	5.57	5.60	5.27	7.48	7.87	7.67	2.83	3.32	2.56

P₆×P₁₀	4.52	4.65	5.13	5.48	5.78	5.54	8.57	8.97	8.22	1.66	2.21	2.48
P₇×P₈	4.55	4.21	5.44	4.61	4.87	4.54	7.98	7.34	8.87	2.54	3.45	2.65
P₇×P₉	6.61	6.76	5.41	4.63	4.87	4.75	6.86	7.23	7.10	2.44	2.22	2.94
P₇×P₁₀	4.43	4.84	5.59	4.56	4.89	4.71	7.53	7.86	7.38	2.73	2.65	2.47
P₈×P₉	4.46	4.32	3.90	5.83	5.97	6.30	7.44	7.97	8.47	2.43	2.80	1.86
P₈×P₁₀	3.67	3.78	3.99	7.03	7.48	6.50	8.48	8.87	8.22	3.32	3.51	3.06
P₉×P₁₀	4.67	4.98	4.67	5.90	6.45	5.88	7.39	7.76	7.58	2.95	3.48	2.81
Mean	4.74	4.87	4.79	5.34	5.50	5.43	7.89	8.05	7.99	2.27	2.65	2.45
C.V.	8.23	8.09	14.86	6.44	7.67	12.75	6.26	6.17	9.48	11.24	10.96	21.02
S.E.±M	0.22	0.23	0.29	0.19	0.24	0.28	0.28	0.29	0.30	0.14	0.17	0.21
C.D. 5%	0.63	0.64	0.81	0.55	0.68	0.78	0.79	0.80	0.86	0.41	0.47	0.58
Range	Lowest	2.98	3.34	3.71	3.56	3.87	4.47	5.40	5.33	6.58	1.15	1.52
	highest	6.66	6.76	6.24	7.03	7.48	6.50	9.65	9.86	9.46	3.32	4.24

...

Genotypes	TSS			Titration acidity %			Ascorbic acid content			Reducing sugar		
	Y ₁	Y ₂	Pooled	Y ₁	Y ₂	Pooled	Y ₁	Y ₂	Pooled	Y ₁	Y ₂	Pooled
P₁	5.69	5.70	5.31	0.46	0.42	0.43	20.50	20.53	21.29	1.11	1.43	1.18
P₂	4.83	4.83	4.76	0.36	0.35	0.42	21.77	21.85	19.25	1.31	1.33	1.57
P₃	4.78	4.79	5.10	0.34	0.37	0.31	24.56	24.59	24.06	0.97	1.54	1.15
P₄	4.84	5.62	5.09	0.38	0.38	0.36	19.79	19.96	21.34	1.10	1.76	1.13
P₅	5.25	4.96	5.11	0.37	0.35	0.40	20.54	20.87	20.15	1.54	1.11	1.47
P₆	4.93	4.85	5.33	0.40	0.34	0.38	22.10	22.56	23.31	1.24	1.50	1.29
P₇	5.01	5.05	5.49	0.45	0.42	0.42	23.54	23.95	23.49	1.66	1.65	1.44
P₈	5.62	4.97	5.24	0.43	0.40	0.37	21.99	22.32	20.38	1.52	1.40	1.43
P₉	5.47	5.47	5.12	0.35	0.30	0.34	19.49	19.64	21.73	1.39	1.76	1.27
P₁₀	4.76	4.90	5.05	0.37	0.35	0.35	21.38	21.85	20.80	1.61	0.49	1.51
P₁×P₂	5.03	5.12	5.36	0.40	0.37	0.4083	22.15	22.78	21.3417	1.24	1.34	1.33
P₁×P₃	4.60	4.60	4.86	0.38	0.32	0.3750	23.59	23.65	23.1867	1.74	1.84	1.54
P₁×P₄	4.49	4.50	4.54	0.42	0.40	0.3700	19.57	19.87	21.6117	1.34	1.72	1.59
P₁×P₅	5.11	5.43	4.80	0.36	0.36	0.3783	20.59	20.78	20.2300	1.24	1.54	1.47
P₁×P₆	5.22	5.12	5.32	0.38	0.39	0.3683	22.33	21.66	21.5533	1.46	1.56	1.50
P₁×P₇	5.07	5.07	5.09	0.39	0.35	0.39	23.71	23.97	22.68	1.11	1.75	1.33

P₁×P₈	5.18	5.87	5.12	0.42	0.45	0.38	24.59	24.67	24.27	1.55	1.70	1.65
P₁×P₉	4.80	4.80	5.33	0.36	0.32	0.40	21.25	21.45	22.96	1.39	1.40	1.54
P₁×P₁₀	4.69	4.70	4.74	0.45	0.48	0.38	16.41	16.74	18.93	1.83	1.84	1.61
P₂×P₃	4.67	4.69	4.75	0.44	0.41	0.39	22.24	22.12	22.04	1.08	1.23	1.20
P₂×P₄	4.70	4.71	4.69	0.43	0.38	0.42	23.56	23.76	22.84	1.24	1.32	1.23
P₂×P₅	5.63	5.69	5.16	0.35	0.35	0.36	24.44	24.87	24.10	1.22	1.28	1.27
P₂×P₆	5.01	5.02	5.35	0.48	0.42	0.41	22.80	22.65	23.83	1.53	1.48	1.40
P₂×P₇	5.05	5.07	5.03	0.43	0.47	0.42	19.66	19.98	21.15	1.17	1.21	1.32
P₂×P₈	5.47	5.34	5.27	0.35	0.35	0.40	23.44	23.74	21.71	1.17	1.20	1.19
P₂×P₉	5.09	5.06	5.21	0.36	0.33	0.35	23.87	23.95	23.80	1.84	1.87	1.52
P₂×P₁₀	5.21	5.43	5.13	0.35	0.29	0.34	23.07	23.56	23.51	1.30	1.33	1.58
P₃×P₄	4.83	5.21	4.80	0.45	0.44	0.41	23.19	23.65	23.89	1.21	1.25	1.37
P₃×P₅	4.59	4.50	4.89	0.34	0.35	0.39	20.49	20.64	22.07	1.26	1.29	1.25
P₃×P₆	4.70	4.78	4.60	0.51	0.55	0.43	23.08	23.32	21.85	1.36	1.39	1.32
P₃×P₇	5.33	5.54	5.05	0.43	0.44	0.49	18.99	19.43	21.15	1.34	1.37	1.36
P₃×P₈	5.81	5.89	5.67	0.47	0.39	0.45	19.71	20.23	19.56	1.45	1.49	1.40
P₃×P₉	4.58	4.79	5.23	0.39	0.33	0.39	20.64	21.23	20.43	1.32	1.37	1.40
P₃×P₁₀	4.83	5.34	4.80	0.38	0.34	0.35	22.50	22.90	21.86	1.13	1.17	1.25
P₄×P₅	4.66	4.97	5.14	0.33	0.28	0.35	19.68	19.87	19.82	1.21	1.20	1.48
P₄×P₆	5.71	5.78	5.33	0.43	0.41	0.35	24.88	24.98	22.37	1.39	1.14	1.29
P₄×P₇	5.11	5.23	5.44	0.35	0.35	0.37	23.18	23.78	24.08	1.45	1.47	1.29
P₄×P₈	5.09	5.67	5.15	0.43	0.46	0.39	21.71	21.75	22.74	1.20	1.22	1.33
P₄×P₉	4.53	4.53	5.10	0.51	0.54	0.48	22.99	23.45	22.37	1.50	1.47	1.36
P₄×P₁₀	5.06	4.98	4.79	0.41	0.44	0.47	19.63	19.76	21.54	1.39	1.41	1.42
P₅×P₆	4.98	5.54	4.97	0.39	0.35	0.36	24.76	25.21	22.81	1.16	1.12	1.13
P₅×P₇	5.18	5.86	5.36	0.41	0.38	0.38	23.84	23.98	24.52	1.39	1.42	1.25
P₅×P₈	4.61	4.79	5.23	0.36	0.34	0.37	18.88	19.45	21.43	1.41	1.44	1.41
P₅×P₉	4.58	4.65	4.68	0.48	0.45	0.41	20.53	21.43	19.99	1.72	1.75	1.58
P₅×P₁₀	5.21	5.73	4.92	0.41	0.36	0.42	23.83	24.53	22.63	1.33	1.34	1.54
P₆×P₇	4.40	4.65	4.62	0.53	0.50	0.43	25.03	25.87	23.79	1.88	1.89	1.69
P₆×P₈	4.69	4.73	4.66	0.37	0.37	0.43	24.77	24.83	25.31	1.12	1.15	1.50

P₆×P₉	4.46	4.75	4.59	0.42	0.41	0.39	21.10	21.45	22.96	1.28	1.30	1.21	
P₆×P₁₀	5.32	5.97	5.03	0.42	0.39	0.41	23.10	23.45	22.27	1.18	1.22	1.24	
P₇×P₈	5.26	5.76	5.15	0.38	0.35	0.40	25.21	25.76	24.57	1.11	1.15	1.38	
P₇×P₉	4.58	5.76	5.17	0.36	0.37	0.35	21.21	21.45	23.48	1.12	1.15	1.13	
P₇×P₁₀	4.74	4.87	5.25	0.38	0.32	0.37	18.56	18.78	20.00	1.32	1.35	1.23	
P₈×P₉	5.13	5.65	5.05	0.40	0.37	0.40	20.61	21.21	21.46	1.02	1.05	1.21	
P₈×P₁₀	4.68	4.78	5.17	0.38	0.34	0.37	23.81	23.97	22.51	1.11	1.16	1.08	
P₉×P₁₀	5.13	5.35	5.30	0.37	0.35	0.33	19.58	20.23	19.61	1.31	1.42	1.53	
Mean	4.98	5.15	5.06	0.40	0.38	0.39	21.97	22.27	22.12	1.33	1.40	1.37	
C.V.	8.36	7.16	9.45	7.70	8.07	12.77	7.16	7.63	9.25	10.67	11.05	15.83	
S.E.±M	0.24	0.21	0.19	0.17	0.02	0.02	0.90	0.98	0.83	0.08	0.09	0.08	
C.D. 5%	0.67	0.60	0.54	0.05	0.05	0.05	2.54	2.75	2.32	0.23	0.25	0.24	
Range	Lowest	4.40	4.50	4.54	0.33	0.28	0.31	16.41	16.74	18.93	0.97	0.50	1.08
	highest	5.80	5.97	5.67	0.52	0.55	0.49	25.20	25.87	25.31	1.88	1.89	1.69

...

Genotypes	Non - reducing sugar			Total sugar			Total fruit yields per plant (kg)		
	Y ₁	Y ₂	Pooled	Y ₁	Y ₂	Pooled	Y ₁	Y ₂	Pooled
P₁	2.68	2.76	2.57	3.51	4.08		2.67	2.86	2.46
P₂	2.67	2.68	2.75	3.98	4.12	3.61	1.94	2.32	2.35
P₃	2.49	2.17	2.36	3.45	3.2	4.33	2.80	2.97	2.78
P₄	2.77	2.23	2.71	3.79	3.73	3.51	3.01	3.45	2.81
P₅	2.54	2.64	2.37	4.07	4.39	3.80	2.22	2.54	2.99
P₆	2.46	2.55	2.50	3.70	3.88	3.85	2.25	2.36	3.24
P₇	2.43	2.54	2.34	4.08	4.08	3.80	2.83	2.83	2.64
P₈	2.88	2.94	2.80	4.05	4.59	3.81	1.88	1.90	2.28
P₉	2.78	2.16	2.56	4.16	3.56	4.06	2.84	2.89	3.25
P₁₀	2.60	2.75	2.54	4.21	4.51	3.83	2.62	2.75	3.09
P₁×P₂	2.44	2.25	2.60	3.68	3.59	3.88	2.32	2.56	2.59

$P_1 \times P_3$	2.38	2.56	2.31	4.12	4.40	3.85	2.15	2.22	2.35
$P_1 \times P_4$	2.64	2.65	2.59	3.98	4.28	4.19	3.18	3.89	2.70
$P_1 \times P_5$	2.69	2.19	2.67	3.93	3.73	4.10	2.56	2.56	3.22
$P_1 \times P_6$	2.29	2.57	2.23	3.75	4.13	3.73	1.19	1.78	1.87
$P_1 \times P_7$	2.59	2.22	2.58	3.70	3.97	3.91	2.86	2.35	2.32
$P_1 \times P_8$	2.37	2.28	2.29	3.92	3.98	3.94	1.96	2.67	2.15
$P_1 \times P_9$	2.33	2.84	2.30	4.98	3.98	4.47	2.77	2.87	2.71
$P_1 \times P_{10}$	3.15	2.84	2.99	3.40	4.68	3.69	2.52	2.76	2.69
$P_2 \times P_3$	2.32	2.54	2.50	4.38	3.77	4.24	1.86	1.87	2.09
$P_2 \times P_4$	3.14	2.78	2.83	4.13	4.12	3.95	3.94	4.12	2.90
$P_2 \times P_5$	2.91	2.86	2.84	3.70	4.14	3.91	2.74	2.97	3.43
$P_2 \times P_6$	2.17	2.45	2.51	3.70	3.93	3.92	2.95	3.45	2.96
$P_2 \times P_7$	2.22	2.32	2.33	3.39	3.53	3.65	2.62	2.34	3.03
$P_2 \times P_8$	2.82	2.21	2.56	4.02	3.41	3.77	2.26	2.23	2.29
$P_2 \times P_9$	2.24	2.28	2.22	4.08	4.15	3.74	2.69	2.87	2.46
$P_2 \times P_{10}$	2.37	2.24	2.32	3.67	3.57	3.91	2.54	2.76	2.70
$P_3 \times P_4$	2.48	2.50	2.32	3.69	3.75	3.48	1.86	2.43	2.41
$P_3 \times P_5$	2.32	2.38	2.40	3.58	3.67	3.66	2.87	2.97	2.65
$P_3 \times P_6$	2.96	2.49	2.67	4.32	3.88	3.99	2.81	2.80	2.89
$P_3 \times P_7$	2.15	2.23	2.31	3.49	3.60	3.68	1.76	1.97	2.28
$P_3 \times P_8$	2.74	2.36	2.48	4.19	3.85	3.89	2.93	3.54	2.44
$P_3 \times P_9$	2.36	2.76	2.36	3.68	4.13	3.76	2.45	2.75	2.99
$P_3 \times P_{10}$	3.14	2.65	2.95	4.27	3.82	4.20	2.18	2.61	2.46
$P_4 \times P_5$	2.35	2.64	2.28	3.56	3.84	3.64	2.77	2.62	3.10
$P_4 \times P_6$	2.85	2.93	2.74	4.24	4.07	4.04	3.42	3.82	3.02
$P_4 \times P_7$	2.24	2.55	2.58	3.69	4.02	3.88	2.79	2.85	3.30
$P_4 \times P_8$	2.37	2.37	2.46	3.57	3.59	3.79	3.56	3.93	3.20
$P_4 \times P_9$	2.93	2.26	2.65	4.43	3.73	4.01	2.90	3.33	3.41
$P_4 \times P_{10}$	2.26	2.21	2.26	3.65	3.62	3.69	3.02	3.76	3.17
$P_5 \times P_6$	2.30	2.56	2.47	3.46	3.68	3.92	2.52	2.54	2.52
$P_5 \times P_7$	2.96	2.62	2.75	4.35	4.04	4.01	1.67	1.63	2.10
$P_5 \times P_8$	2.51	2.52	2.56	3.92	3.96	3.98	2.50	2.53	2.06

$P_5 \times P_9$	2.45	2.36	2.48	4.17	4.11	4.06	1.88	1.93	2.20	
$P_5 \times P_{10}$	2.39	2.55	2.37	3.72	3.89	3.91	3.59	4.23	2.75	
$P_6 \times P_7$	2.61	2.92	2.58	4.74	4.81	4.31	2.66	2.72	2.51	
$P_6 \times P_8$	2.36	2.55	2.64	3.48	3.70	4.14	1.99	2.45	2.35	
$P_6 \times P_9$	2.28	2.25	2.41	3.56	3.55	3.63	3.67	3.53	3.06	
$P_6 \times P_{10}$	2.21	2.25	2.22	3.39	3.54	3.47	1.88	2.45	2.70	
$P_7 \times P_8$	2.38	2.40	2.46	3.56	3.55	3.82	2.88	3.56	2.85	
$P_7 \times P_9$	2.35	2.32	2.37	3.47	3.47	3.51	2.84	2.32	3.20	
$P_7 \times P_{10}$	2.40	2.73	2.36	3.73	4.08	3.60	2.83	2.67	2.57	
$P_8 \times P_9$	2.50	2.31	2.72	3.52	3.30	4.05	2.57	2.86	2.2	
$P_8 \times P_{10}$	2.48	2.35	2.39	3.59	3.51	3.44	3.41	3.63	3.13	
$P_9 \times P_{10}$	2.42	2.48	2.29	3.73	3.83	3.64	3.20	3.56	3.04	
Mean	2.52	2.49	2.50	3.86	3.89	3.87	2.61	2.82	2.71	
C.V.	10.67	9.70	11.2394	7.64	10.80	11.07	10.75	11.31	20.34	
S.E. $\pm M$	0.15	0.14	0.1150	0.17	0.24	0.17	0.16	0.18	0.22	
C.D. 5%	0.97	0.39	0.3203	0.47	0.68	0.48	0.45	0.52	0.62	
Range	Lowest	2.14	2.16	2.22	3.39	3.28	3.44	1.18	1.63	1.87
	highest	3.14	2.94	2.99	4.98	4.81	4.47	3.94	4.23	3.43

...

References;

1. Anonymous. Horticulture Data Base, National Horticulture Board, Gurgaon, Ministry of Agriculture, India 2020.
2. Narolia RK, Reddy RVSK, Padma M. Correlation, path coefficient and genetic divergence analysis of growth, yield and quality of tomato (*Lycopersicon esculentum* Mill.) *Indian J. Crop Bio.* 2012; 20(1):65-69.
3. Mohamed SM, Ali EE, Mohamed TY. Study of Heritability and Genetic Variability among Different Plant and Fruit Characters of Tomato (*Solanum lycopersicum* L.). *Int. J Sci. Tech. Res.* 2012; 1(2):55-58.
4. Prakash, O., Choyal, P., Godara, A., & Choudhary, S. Mean performance of tomato (*Solanum lycopersicum* L.) genotypes for yield, yield parameters and quality traits. *J. Pharma. Inno.* 2019; 8,763-765.
5. Anuradha, B., Saidaiah, P., Ravinder Reddy, K., & Harikishan Sudini, G. A. Mean performance of 40 genotypes in tomato (*Solanum lycopersicum* L.). *Int. J. Chem. Stud.* 2021; 9(1), 279-283.
6. Joshi A, Kohli UK. Genetic divergence for qualitative and quantitative traits in tomato (*Lycopersicum esculentum* Mill.). *Indian J. Agri. Sci.* 2003; 73(2):110-113.
7. Jogi P, Shukla N, Mehta N, Sahu M. Genetic divergence for fruit traits in tomato (*Lycopersicum esculentum* Mill.). *Oris.J. Hort.*, 2008; 36(2):149-151.
8. Kerketta, A., Bahadur, V., & Rajesh, J. Performance of different tomato genotypes (*Solanum lycopersicum* L.) for growth, yield and quality traits under Allahabad condition. *J. Pharma. Phyto.* 2018.;7(6), 1766-1769.
9. Martí, R., Roselló, S., & Cebolla-Cornejo, J. Tomato as a source of carotenoids and polyphenols targeted to cancer prevention. *Cancer.*, 2016;8(6), 58.

Comment [r9]: Where is the conclusion section?

Formatted: Tab stops: 1.03", Left

Comment [r10]: Do not use anonymous references

UNDER PEER REVIEW

