

Original Research Article

“Effect of Irrigation Scheduling and Foliar Organic Nutrition on yield and economics of Summer Groundnut (*Arachishypogaea L.*)”

ABSTRACT

A field experiment was conducted during *Summer*season of 2022 at Crop Research Farm (CRF), Department of Agronomy, SHUATS, Prayagraj (UP) on soil with sandy loam in texture to investigate the effect of Irrigation Scheduling and Foliar Organic Nutrition on growth and yield of Zaid Groundnut. The treatments consist of three Irrigation Scheduling viz., I₁: 3 Irrigations (25,45,70 DAS), I₂: 2 Irrigations (25,45 DAS), I₃: 2 Irrigations (25,70 DAS) and three Foliar Organic Nutrition comprising of F₁ – Panchagavya at 3%, F₂ – Jeevamrutha at 3%, F₃ – Panchamrutha at 3% whose effect is observed on Groundnut (var. K-6). The experiment was laid out in Randomized Block Design with Ten treatments replicated thrice. The treatment with application of 3 Irrigations (25,45,70 DAS) + panchagavya-3% recorded significantly higher number of pods per plant (20.33), number of kernels per pod (2.47), seed index (40.08 g), pod yield (2.85 t/ha), haulm yield (4.4 t/ha) gross returns (128344.5 INR), net returns (74,798.30 INR) and B:C ratio (2.02) compared to other treatment combinations.

Keywords: *Groundnut, Irrigation Scheduling, Panchagavya, Jeevamrutha, Panchamrutha Yield, Economics.*

Introduction

A common leguminous crop grown across the tropics and subtropics is groundnut (*Arachishypogaea L.*). It is valued for its high-oil edible seeds, making it the third-most significant source of vegetable protein in the world and the fourth-most significant source of edible oil. Groundnut is not only an important oilseed crop of India but also an important agricultural export commodity. India ranks first in Groundnut acreage and is the second largest producer of Groundnut in the world with 67 lakh tonnes after China with a productivity of 1422 kg per hectare (FAOSTAT, 2019). Although groundnut is grown in one or more of the

seasons (kharif, rabi, and summer), the kharif crop accounts for roughly 80% of the acreage and production (June–October). A significant oilseed crop, groundnut will provide about 37% of the nation's total oilseed production in 2020–21. Groundnut acreage in the nation varies with time, and in the last 20 years, it has decreased from 83 lakh ha to 47 lakh ha as farmers switch to other lucrative crops in response to the green revolution, which mostly focused on cereal crops.

Irrigation plays a major role in crop growth and development. Irrigation also affects the availability of plant nutrients, with proper irrigation to the crop at right time and right intervals maximum growth and yield can be achieved. Both excess and shortage of water can affect crop growth and yield adversely. Being a day neutral crop, ample irrigations are required by groundnut in summer and missing any irrigation at any one of the critical growth stages results into considerable reduction in pod yield. High soil moisture results in pod rot, low yield and poor seed quality and also water stress reduces photosynthesis mainly due to reduction in photosynthetic area. **Madhuri Devi T., et al., (2019)**. Hence proper research is required to find proper irrigation scheduling.

Excessive use of **Chemical** fertilizers in agriculture after green revolution led to soil, water, environmental pollution along with health hazards. Organic liquid manures have variety of advantages over synthetic chemicals and supply all the essential plant nutrients in balanced manner naturally without deteriorating the soil health. Sustainable farming using organic inputs is essential to reduce the effects of synthetic fertilisers on soil health, the environment, and the preservation of ecological biodiversity. Hence this research is involved to find proper irrigation scheduling along with foliar application of liquid organic manures in summer ground nut.

Materials and Methods

A field experiment was conducted during *Zaid* season of 2022 at Crop Research Farm (CRF), Department of Agronomy, SHUATS, Prayagraj (UP). The soil of the experimental plot is sandy loam in texture, nearly neutral in soil reaction (pH 7.4), medium in organic carbon (0.48%), medium in available Nitrogen (278.93 kg/ha), low in available Phosphorous (19.03 kg/ha) and medium in available Potash (238.1 kg/ha). The treatments consist of three Irrigation Scheduling viz., I₁: 3 Irrigations (25,45,70 DAS), I₂: 2 Irrigations (25,45 DAS), I₃: 2 Irrigations (25,70 DAS) and three Foliar Organic Nutrition F₁ – Panchagavya at 3%, F₂ –

Jeevamrutha at 3%, F3 – Panchamrutha at 3% whose effect is observed on Groundnut (var. K-6). The experiment was laid out in Randomized Block Design with ten treatments replicated thrice. The experiment comprising ten treatment possible combination of above factor, viz., T₁: 3 Irrigations (25,45,70 DAS) + panchagavya-3% T₂: 3 Irrigations (25,45,70 DAS) + Jeevamrutha-3%, T₃: 3 Irrigations (25,45,70 DAS) + Panchamrutha-3%, T₄: 2 Irrigations (25,45 DAS) + panchagavya-3%, T₅: 2 Irrigations (25,45 DAS) + Jeevamrutha-3%, T₆: 2 Irrigations (25,45 DAS) + Panchamrutha-3%, T₇: 2 Irrigations (25,70 DAS) + panchagavya-3%, T₈: 2 Irrigations (25,70 DAS) + Jeevamrutha-3%, T₉: 2 Irrigations (25,70 DAS) + Panchamrutha-3%, T₁₀: Control. Observations regarding growth and yield attributes was recorded during the field experiment.

Comment [M1]: How the control was treated?

Result and Discussion

Yield attributes

According to the yield characteristics data that was collected and analysed at harvest, maximum number of pods per plant (20.33), maximum number of kernels per pod (2.47) and higher seed index (40.08g) was recorded in treatment with the application of 3 Irrigations (25,45,70 DAS) + panchagavya-3%.

The improved performance of yield attributes may be due to the availability of adequate soil moisture at the root zone, which in turn feeds the crop with all of the nutrients needed during all phases of growth and therefore increased the values of yield attributes. These findings are closely in line with those of **BN Solanke et al., (2021)**, **Bibhu Santosh Behera et al., (2015)**.

Yield

After evaluated the data recorded post harvesting of crop show that significantly higher pod yield (2.47 t/ha), higher haulm yield (4.4 t/ha) and harvest index (39.31%) was recorded in treatment with the application of 3 Irrigations (25,45,70 DAS) + panchagavya-3%.

The quantity of IAA and GA present in panchagavya spray as well as the simplicity with which nutrients may be applied to plants by foliar spray may have stimulated the plant system, increasing the production of growth regulators in the cell system. Consequently, this

Comment [M2]: expansion

Comment [M3]: expansion

encouraged the necessary growth and development in plants, improving yield. These results are in close conformity with the findings of **Ravi Kumar et al., (2012)**, **Vikash et al., (2022)**.

Economics

The economic return of Groundnut was analyzed after harvesting the crop based on market pricing, the result indicated a growing trend in with the increasing yield trend across treatment.

The maximum Gross returns (INR **128344.5** /ha), Net returns (INR **74,798.30** /ha) and Benefit cost ratio (2.40) was evaluated in treatment with the application of 3 Irrigations (25,45,70 DAS) + panchagavya-3%.

Conclusion

Based on the above experimental findings, it is concluded that application of 3 Irrigations (25,45,70 DAS) + panchagavya-3% accomplished better growth parameters, yield attributes, higher seed yield, higher gross returns and net returns in Groundnut crop under eastern Uttar Pradesh conditions.

Future Scope

The conclusion drawn are based on one season experimentation only which requires further confirmation for recommendation.

References

- Bibhu Santosh Behera, Mohit Das, Anama Charan Behera & Rudra Ashish Behera "Weather Based Irrigation Scheduling in Summer Groundnut in Odisha Condition" *International Journal of Agricultural Science and Research (IJASR)* ISSN(P): 2250-0057; ISSN(E): 2321-0087; Vol. 5, Issue 5, Oct 2015, 247-260
- BN Solanke, RH Shinde, NS Lolamwad and AS Bhosale Yield and economics of summer groundnut (*Arachis hypogaea L.*) as influenced by different irrigation regimes and land configurations. ISSN (E): 2277-7695 ISSN (P): 2349-8242 NAAS Rating: 5.23 TPI 2021; SP-10(11): 2857-2860
- D.C. Lokhande, N.E. Jayewar and A.G. Mundhe Summer Groundnut (*Arachis hypogaea L.*) Productivity Influenced by Irrigation Scheduling: A Climatological Approach.

International Journal of Current Microbiology and Applied Sciences ISSN: 2319-7706 Special Issue-6 pp. 87-91.

J Pradhan, A Baliarsingh, S Pasupalak, AKB Mohapatra. Effect of different irrigation scheduling & mulching on productivity of groundnut (*Arachishypogaea L.*) of east & south eastern coastal plain of Odisha. *Pharma Innovation* 2018;7(10):689-691

Kumawat, R.N., Mahajan, S.S. and Mertia, R.S. 2009. Growth and development of groundnut under foliar application of panchgavya and leaf extracts of endemic plant. *Indian Journal of Agronomy*, 3: 324-331.

Madhuri Devi T., et al., (2019) Effects of Irrigation Regime and Phosphorus Level on Yield and Yield Attributes of Summer Groundnut (*Arachis hypogea L.*). *International Journal of Agriculture Sciences*, ISSN: 0975-3710 & E-ISSN: 0975-9107, Volume 11, Issue 12, pp.- 8673-8676.

Naresha, R., Laxminarayana, P., Suneetha Devi, K. B. And Sailaja, V., Effect of Irrigation Scheduling and Phosphogypsum Levels on Yield Attributes, Yield and Available Nutrients in Soil after Harvest of Rabi Groundnut, *Int. J. Pure App. Biosci.*6(2): 1300-1308 (2018). doi: <http://dx.doi.org/10.18782/2320-7051.6361>.

PB Kotadiya, DS Hirpara, LC Vekariya, HN Kanjiya. Response of summer groundnut (*Arachishypogaea L.*) to irrigation level and anti-transpirant on quality, soil parameters and economics. *Pharma Innovation* 2021;10(10):315-317.

Ravi Kumar H.S. &Janakiraman, N &Sheshadri, T & Gowda, J &Vijaymahantesh, And. (2012). Integrated Organic Nutrient Supply Systems on Growth and Yield of Groundnut (*Arachishypogaea L.*). *Environment and Ecology*. 30. 118-121.

Sunil Kumar Meena, Rajendra Kumar Yadav, Baldev Ram, Vinod Kumar Yadav, SL Yadav, MK Sharma, Anil Meena. Effect of irrigation schedules and soil amendments on growth and nutrient content & uptake of soybean under Vertisols of South-Eastern Rajasthan. *Pharma Innovation* 2021;10(12):1298-1302.

Vikash, Matam& Singh, V. & George, Shruti&Vishkarma, S.P.. (2022). Performance of Different Organic Liquid Manures on Growth, Yield and Economics of Field Pea

(Pisumsativum L.). *International Journal of Plant & Soil Science*. 44-48.
10.9734/ijpss/2022/v34i1931087.

UNDER PEER REVIEW




Table 1. Yield attributes of Groundnut as influenced by Irrigation Scheduling and Liquid Organic Manures.

Treatment	Yield attributes		
	No. of pods/plant	No. of kernels/pod	Seed Index (g)
3 Irrigations (25,45,70 DAS) + panchagavya-3%	20.33	2.47	40.08
3 Irrigations (25,45,70 DAS) + Jeevamrutha-3%	19.67	2.27	38.96
3 Irrigations (25,45,70 DAS) + Panchamrutha-3%	19.53	2.07	38.61
2 Irrigations (25,45 DAS) + panchagavya-3%	18.87	2.00	37.47
2 Irrigations (25,45 DAS) + Jeevamrutha-3%	18.40	1.87	36.77
2 Irrigations (25,45 DAS) + Panchamrutha-3%	18.27	1.73	36.39
2 Irrigations (25,70 DAS) + panchagavya-3%	18.13	1.60	35.73
2 Irrigations (25,70 DAS) + Jeevamrutha-3%	17.53	1.60	35.64
2 Irrigations (25,70 DAS) + Panchamrutha-3%	17.47	1.60	35.17
Control	15.47	1.40	32.43
F test	S	S	S
SEm(±)	0.06	0.05	0.16
CD(p=0.05)	0.19	0.14	0.49

Comment [M4]: dimension missing

Comment [M5]: for area or number of plant??

Table 2. Yield of Groundnut as influenced by Irrigation Scheduling and Liquid Organic Manures.

Treatment	Pod Yield (t/ha)	Haulm Yield (t/ha)	Harvest Index (%)
3 Irrigations (25,45,70 DAS) + panchagavya-3%	2.85	4.40	39.31
3 Irrigations (25,45,70 DAS) + Jeevamrutha-3%	2.76	4.28	39.24
3 Irrigations (25,45,70 DAS) + Panchamrutha-3%	2.55	4.26	37.42
2 Irrigations (25,45 DAS) + panchagavya-3%	2.36	4.22	35.84
2 Irrigations (25,45 DAS) + Jeevamrutha-3%	2.27	4.17	35.21
2 Irrigations (25,45 DAS) + Panchamrutha-3%	2.20	4.13	34.74
2 Irrigations (25,70 DAS) + panchagavya-3%	2.19	3.88	36.02
2 Irrigations (25,70 DAS) + Jeevamrutha-3%	2.12	3.76	36.08
2 Irrigations (25,70 DAS) + Panchamrutha-3%	2.09	3.58	36.84
Control	1.99	3.37	37.11
F test	S	S	S
SEm(±)	0.01	0.01	0.13

CD(p=0.05)

0.04

0.02

0.38

Table 3. Economics of Groundnut at harvest as influenced by Irrigation Scheduling and Liquid Organic Manures.

S. No	Treatments	Economics			
		Cost of Cultivation	Gross returns	Net Returns	B:C ratio
1.	3 Irrigations (25,45,70 DAS) + panchagavya-3%	53,546.2	161808.4	108,262.20	2.02
2.	3 Irrigations (25,45,70 DAS) + Jeevamrutha-3%	54,446.2	124378.5	69,932.30	1.82
3.	3 Irrigations (25,45,70 DAS) + Panchamrutha-3%	54,746.2	114721.5	59,975.30	1.58
4.	2 Irrigations (25,45 DAS) + panchagavya-3%	53,046.2	106059	53,012.80	1.47
5.	2 Irrigations (25,45 DAS) + Jeevamrutha-3%	53,946.2	102012	48,065.80	1.33
6.	2 Irrigations (25,45 DAS) + Panchamrutha-3%	54,246.2	98869.5	44,623.30	1.25
7.	2 Irrigations (25,70 DAS) + Panchagavya-3%	53,046.2	98424	45,377.80	1.29
8.	2 Irrigations (25,70 DAS) + Jeevamrutha-3%	53,946.2	95470.5	41,524.30	1.18
9.	2 Irrigations (25,70 DAS) + Panchamrutha-3%	54,246.2	94042.5	39,796.30	1.14
10.	Control	51,546.2	89391	37,844.80	1.14

*Data was not subjected to statistical analysis.