

Original Research Article

Effect of nanoscale zinc oxide particles on macronutrient concentration of Groundnut (*Arachis hypogaea* L.)

Abstract

In the present investigation, size dependent effects of nanoscale zinc oxide particulates (n-ZnO) on the macronutrient concentration of groundnut leaf, stem and kernel have been analysed. ZnO-nanoparticulates that were used in the study were prepared by modified oxalate decomposition method and the ZnO-nanoparticulates (mean size of 20, 25 and 30 nm) were characterized using techniques such as transmission electron microscopy (TEM), Fourier transform infrared spectroscopy (FT-IR), Dynamic light scattering (DLS) and X-ray diffraction analysis (XRD). Different concentrations (150, 200 and 400 ppm) of ZnO-nanoparticulates were applied (foliar spray) to reveal their effects on groundnut crop in comparison to bulk ZnSO₄. These results indicate that zinc nanoparticles significantly influenced the macronutrient (N, P, K) concentration of groundnut depending on their size and concentration.

Key words: n-ZnO, size, concentration, macronutrients.

1. INTRODUCTION

Nanoscale materials (size less than 100 nm in at least one dimension) exhibits unique and novel properties compared to their bulk counter parts (Prasad et al., 2012). However, the application of nanoscale materials in agriculture as nutrients is relatively new and the record of consequent effects on crops is scant. Further, it is clear from the theory that nanoscale materials possess size dependent characteristics and reactivity and also are distinct to each other.

Nanotechnology plays a vital role in improving soil health, nutrient management, weed management, pest and disease control, through the new scientific approaches to increase production and productivity of crops.

The present study examines the interactions between Zn and other nutrients in soil, behaviour in plant growth. It stresses the need for identification of the factor responsible for any Zn response to the addition of another nutrient compound. Of the many interactions of Zn with other nutrients, the most widespread and important to crop production are those with N and P fertilizers on soils with limiting supplies of both Zn and N or P. Similar interactions of Zn with other essential nutrients will also be important in soils with low fertility. It helps to introduce new techniques through enabling slow and controlled release of nutrients from fertilizers, efficient and targeted delivery of fertilizers coupled with enabling resistance, effective processing, storage and packing. Nanoparticles have smaller particle sizes, higher specific surface area and an increased proportion of reactive surface atoms as compared to bulk particles [12]. Zinc nanoparticles are being used in various agricultural experiments by the researchers to understand its effect on growth, germination, and various other properties [8, 11, 6, and 9] and reported encouraging results.

2. MATERIALS AND METHODS:

ZnO nanoparticles of mean size of 20, 25, 30 nm diameter were used in the study. Nanocrystalline zinc oxide has been prepared by using the oxalate decomposition technique. Zinc oxalate was prepared by mixing equimolar (0.2 M) solutions of zinc acetate and oxalic acid. The resultant precipitate was collected and rinsed extensively with double deionized water (DI-water) and dried in air. The oxalate was then ground and decomposed in air by placing it in a pre-heated furnace for 45 minutes at 500°C. The characterization of the samples was done by Dynamic Light Scattering analysis, Transmission Electron Microscopy. The TEM samples were prepared by drop casting the suspensions on carbon coated Cu grids.

The experiment was conducted at College farm, S. V. Agricultural College, Acharya N.G. Ranga Agricultural University, Tirupati during *Kharif*, 2018-19. The experiment was laid out in sandy clay loam textured soil in a randomized block design (RBD) with three replications and with the plot size of 4m× 4m. The initial soil parameters were pH 6.42 (neutral); EC = 0.132 dSm⁻¹; organic carbon = 0.50% (low); available nitrogen = 188.16 kg ha⁻¹ (low); available P₂O₅ =

14.66 kg ha⁻¹; available K₂O= 564.4 kg ha⁻¹ (high); available zinc = 16.6 ppm; and total zinc content of 21.3 ppm. Laboratory analysis is done by following standard procedures given by Jackson, [4] and piper, [7].

Field experiment was carried out in *kharif* 2018 with twelve treatments and three replications. The treatments were *viz.*, control *i.e.*, no application (T₁), Recommended Dose of Fertilizer RDF (T₂), RDF + Zinc sulphate @ 2000 ppm at 25 and 45 DAS (T₃), RDF + Nanoscale zinc oxide (20 nm) @ 400 ppm (T₄), , RDF + Nanoscale zinc oxide (20 nm) @ 200 ppm (T₅), RDF + Nanoscale zinc oxide (20 nm) @ 150 ppm (T₆), RDF + Nanoscale zinc oxide (25 nm) @ 400 ppm (T₇), RDF + Nanoscale zinc oxide (25 nm) @ 200 ppm (T₈), RDF + Nanoscale zinc oxide (25 nm) @ 150 ppm (T₉), RDF + Nanoscale zinc oxide (30 nm) @ 400 ppm (T₁₀), RDF + Nanoscale zinc oxide (30 nm) @ 200 ppm (T₁₁) and RDF + Nanoscale zinc oxide (30 nm) @ 150 ppm (T₁₂).

3. RESULTS AND DISCUSSION

The data on post-harvest concentration of macronutrients (N, P and K) in leaf, stem and kernel at harvest as influenced by the application of nano ZnO and bulk ZnSO₄ are presented in the Table 1.

3.1 Concentration of macronutrients in leaf, stem, kernel at harvest

3.1.1 Nitrogen content (%)

At harvest, the concentration of nitrogen in groundnut leaves, stem was numerically higher in when compared to control and bulk ZnSO₄ @ 2000 ppm but they are not significantly different. Highest leaf N content (0.84 %) was observed in treatment of 100 % RDF (T₂). Whereas, highest stem N content (0.70 %) was observed in treatment (T₇) n-ZnO of size 25nm @ 400 ppm over other treatments. Statistically significant high kernel N content (0.49 %) was observed in T₁₀ treatment n-ZnO of size 30 nm @ 400 ppm which is 45 % more than control and 49 % more than bulk ZnSO₄ @ 2000 ppm.

3.1.2 Phosphorous content (%)

Phosphorous content in groundnut leaves, stem and kernel was significantly higher when compared to control and bulk ZnSO₄ @ 2000 ppm. Highest leaf P content (0.23 %) was observed in treatment n- ZnO of size 30 nm @ 400 ppm (T₁₀) which is 73 % more than control, 43 % more than bulk ZnSO₄ @ 2000 ppm and it is on par with T₁₁ (0.22) n- ZnO of size 30 nm @ 200 ppm. The next best treatments were T₉ (0.17%) and T₁₂ (0.14%).

Highest stem P content (0.27 %) was observed in treatment T₁₀ (n-ZnO of size 30 nm @ 400 ppm) which is 70 % more than control and 66.6 % more than bulk ZnSO₄ @ 2000 ppm. Highest P content in kernel (0.16 %) was observed in T₁₂ treatment (n- ZnO of size 30 nm @ 150 ppm) which is 62.5 % more than control and 44 % more than bulk ZnSO₄ @ 2000 ppm.

3.1.3 Potassium content (%)

The concentration of potassium in leaves, stem and kernels of groundnut at harvest was significantly higher in when compared to control and bulk ZnSO₄ @ 2000 ppm. Highest leaf K content (1.08 %) was observed in treatment n- ZnO of size 20 nm @ 400 ppm (T₄) which is 20 % more than control and 23 % more than bulk ZnSO₄ @ 2000 ppm. Highest stem K content (1.12 %) was observed in treatment T₇ (n- ZnO of size 25 nm @ 400 ppm) which is 15 % more than control and 20.5 % more than bulk ZnSO₄ @ 2000 ppm. Highest kernel K content (0.7 %) was observed in both T₈ and T₁₁ treatments n-ZnO of size 25 nm @ 200 ppm and n- ZnO of size 30 nm @ 200 ppm respectively, which is 20 % more than control and 10 % more than bulk ZnSO₄ @ 2000 ppm. The next best treatments were T₃ (0.63), T₄ (0.63), T₇ (0.63) and T₁₂ (0.63) and these results were in good agreement with Afshar *et al* [1], Singh *et al.* [10], EI-Metwally *et al.* [3].

Optimal levels of zinc improve the uptake of phosphorus and potassium. Zinc plays a key role in chlorophyll which increases greenness that led to increased uptake of nutrients. The increase in total N, K and Zn uptake could be attributed to the synergistic effect between N and Zn and due to the positive interaction of K and Zn, respectively. The present findings support the results of Ashoka *et al.* [2], Morshedi and Farahbakhsh [5]. The mobility of the nanoparticles is known to be very high which ensures the phloem transport and ensures the nutrient to reach all parts of the plant thereby affecting the enzyme reactions, increased dry-matter production which led to increased nutrient uptake. This may be the reason for higher zinc content in grain and

lower zinc content in dry-matter at harvest with RDF along with nanoscale nutrients in combination than bulk form of nutrient.

Table 1. Size dependent effects of nanoscale ZnO particles on the concentration of macro nutrients in leaf, stem and kernel at harvest

Treatment	N concentration %			P concentration %			K concentration %		
	Leaf	stem	kernel	Leaf	stem	kernel	Leaf	stem	Kernel
T ₁ : control	0.14 ^c	0.17 ^b	0.27 ^{de}	0.06 ^e	0.08 ^e	0.06 ^e	0.86 ^{cd}	0.95 ^{cde}	0.56 ^b
T ₂ : RDF	0.84 ^a	0.60 ^{ab}	0.44 ^{ab}	0.1 ^{cde}	0.13 ^{bcd}	0.08 ^{de}	0.84 ^d	0.97 ^{bcde}	0.60 ^b
T ₃ : RDF + ZnSO ₄ @ 2000 ppm	0.39 ^{abc}	0.26 ^{ab}	0.25 ^e	0.13 ^{bcd}	0.09 ^{de}	0.09 ^{cd}	0.83 ^d	0.89 ^{de}	0.63 ^{ab}
T ₄ : RDF + Nano ZnO (20nm) @ 400 ppm	0.28 ^{abc}	0.38 ^{ab}	0.35 ^{bcd}	0.11 ^{cde}	0.12 ^{bcde}	0.09 ^{cd}	1.08 ^a	0.94 ^{cde}	0.63 ^{ab}
T ₅ : RDF + Nano ZnO (20 nm) @ 200 ppm	0.27 ^{abc}	0.23 ^{ab}	0.31 ^{cde}	0.12 ^{cd}	0.17 ^b	0.12 ^{abc}	0.94 ^{bcd}	0.94 ^{cde}	0.60 ^b
T ₆ : RDF + Nano ZnO (20 nm) @ 150 ppm	0.37 ^{abc}	0.25 ^{ab}	0.35 ^{bcd}	0.14 ^{bcd}	0.16 ^{bc}	0.12 ^{abc}	0.85 ^d	0.84 ^e	0.60 ^b
T ₇ : RDF + Nano ZnO (25 nm) @ 400 ppm	0.79 ^{ab}	0.70 ^a	0.40 ^{abc}	0.09 ^{de}	0.1 ^{de}	0.13 ^{abc}	0.98 ^{abc}	1.12 ^a	0.63 ^{ab}
T ₈ : RDF + Nano ZnO (25 nm) @ 200 ppm	0.40 ^{abc}	0.27 ^{ab}	0.35 ^{bcd}	0.12 ^{cd}	0.11 ^{cde}	0.15 ^{ab}	0.99 ^{ab}	1.09 ^{ab}	0.70 ^a
T ₉ : RDF + Nano ZnO (25 nm) @ 150 ppm	0.29 ^{abc}	0.25 ^{ab}	0.40 ^{abc}	0.17 ^b	0.14 ^{bcd}	0.13 ^{abc}	1.03 ^{ab}	0.95 ^{cde}	0.60 ^b
T ₁₀ : RDF + Nano ZnO (30 nm) @ 400 ppm	0.17 ^{bc}	0.23 ^{ab}	0.49 ^a	0.23 ^a	0.27 ^a	0.11 ^{bcd}	0.92 ^{bcd}	0.98 ^{bcd}	0.56 ^b
T ₁₁ : RDF + Nano ZnO (30 nm) @ 200 ppm	0.29 ^{abc}	0.26 ^{ab}	0.37 ^{bc}	0.22 ^a	0.11 ^{cde}	0.11 ^{bcd}	0.95 ^{bcd}	1.03 ^{abc}	0.70 ^a
T ₁₂ : RDF + Nano ZnO (30 nm) @ 150 ppm	0.29 ^{abc}	0.34 ^{ab}	0.41 ^{abc}	0.14 ^{bc}	0.11 ^{cde}	0.16 ^a	1.03 ^{ab}	1.10 ^{ab}	0.63 ^{ab}
SE(m)	0.18	0.14	0.030	0.013	0.016	0.012	0.039	0.04	0.024
CD	NS	NS	0.09	0.039	0.047	0.036	0.110	0.110	0.069

*The mean values were separated by Duncan's Multiple Range Test (DMRT)

4. Conclusions

While Zn interacts with other nutrients in many ways, few, other than those involving correction of deficiencies of both Zn and another nutrient, appear to be important in crop production. Where interactions do occur, they sometimes result, not from the nutrient to which they are attributed, but from other factors associated with the addition of the nutrient compound. The results of *kharif* season shows that, N, P and K concentrations in leaves, shoot and kernels

varied significantly with the foliar application of different sizes and concentrations nanoscale ZnO particles.

References:

1. Afshar I, Akbar R, Minoos S. Comparison of the effects of spraying different amounts of nano ZnO and bulk ZnO on wheat. *International Journal of Plant, Animal and Environmental Sciences*.2014; 4:688-693.
2. Ashoka P, Mudalagiriappa BK, Desai. Effect of micronutrients with or without organic manures on yield of baby corn-chickpea sequence. *Karnataka Journal of Agricultural Sciences*. 2008; 21(4): 485-487.
3. El-Metwally IM, Doaa MR, Abo-Basha ME, El-Aziz A. Response of peanut plants to different foliar applications of nano- iron, manganese and zinc under sandy soil conditions, *Middle East Journal of Applied Sciences*.2018; 8(2): 474-482
4. Jackson ML. *Soil chemical analysis*. Prentice Hall of India Pvt. Ltd. New Delhi. 1973; 134-204.
5. Morshedi A, Farahbakhsh H. Effects of potassium and zinc on grain protein contents and yield of two wheat genotypes under soil and water salinity and alkalinity stresses. *Plant Ecophysiology*. 2010; 2: 67-72.
6. Naseeruddin R, Sumathi V, Prasad TNVKV, Sudhakar P, Chandrika V, Ravindra Reddy B. Unprecedented synergistic effects of nanoscale nutrients on growth, productivity of sweet sorghum [*Sorghum bicolor* (L.) Moench], and nutrient biofortification. *Journal of Agriculture and Food Chemistry*. 2018; 66: 1075-1084.
7. Piper, CS. *Soil and plant analysis*. 1950; Hans publishers, Bombay. India
8. Prasad TNVKV, Sudhakar P, Sreenivasulu Y, Latha P, Munaswamy V, Raja Reddy K, Sreeprasad, TS, Sajanlal PR and Pradeep T. Effect of nanoscale zinc oxide particles on the germination, growth and yield of peanut. *Journal of Plant Nutrition*. 2012; 35: 905-927.
9. Ramapuram N, Sumathi V, Prasad TNVKV, Sudhakar P, Chandrika V, Reddy, BR. Unprecedented synergistic effects of nanoscale nutrients on growth, productivity of sweet sorghum [*Sorghum bicolor* (L.) Moench] and nutrient biofortification. *Journal of Agricultural and Food Chemistry*. 2018; DOI: 10.1021/acs.jafc.7b04467.
10. Singh MD, Jayadeva HM, Chirag G, Mohan MH. Effects of nano zinc oxide particles on seedling growth of maize (*Zea mays* L.) in germinating paper test. *International Journal of Microbiology Research*. 2017; 9(5): 897-898.
11. Subbaiah LV, Prasad TNVKV, Krishna TG, Sudhakar P, Reddy BR, Pradeep T. Novel effects of nanoparticulate delivery of zinc on growth, productivity, and zinc

biofortification in maize (*Zea mays* L.). *Journal of Agricultural and Food Chemistry*. 2016; DOI:10.1021/acs.jafc.6b00838J.

12. Wigginton NS, Haus KL, Hochella MF. Aquatic environmental nanoparticles. *Journal of Environmental Monitoring*. 2007; 9: 1306 -1316.

UNDER PEER REVIEW