

Effect of addition of carbon through residues in Rice-Rice cropping system under long term fertilizer experiment on soil carbon stock and biological properties in an *Inceptisol* of Bhubaneswar, Odisha

ABSTRACT

Aims: To measure the effect of addition of carbon through residues in Rice-Rice cropping system in the long-term fertiliser experiment on soil carbon stock and biological properties in an *Inceptisol* of Bhubaneswar, Odisha.

Study Design: It was carried out with six treatments during the eighth crop cycle viz, 100% NPK, 100% NPK + FYM, 100% N, 100% NP, 100% NPK +Lime, control (no manuring) with 4 replications in a randomized block design. Both in the kharif and the rabi seasons, rice stubble with undisturbed roots was meticulously collected, processed, and tested for total carbon after rice harvest. The usual approach was followed when collecting and analysing soil samples. Between 1223.5 and 2571.5 kg/ha of carbon and 2998.9 to 6330.85 kg/ha of total stubble were absorbed into the soil.

Results: After kharif and rabi, the surface carbon stock of the soil ranged from 7.41 to 12.50 Mg/ha and 7.14 to 11.76 Mg/ha, respectively. After kharif, the SOC of surface soil ranged between 3.48 and 6.51 g/kg and 3.35 and 6.13 g/kg. In 100% NPK+ FYM, the highest amounts of stubble, stubble carbon, and MBC were found. No manuring enhanced the BD (1.42 Mg/m³) but the addition of FYM and stubble decreased the BD (1.28 Mg/m³). MBC varied from 45.89 to 132.41 g of carbon per g of soil. The importance of stubble addition in enhancing SOC was demonstrated by the significantly positive association between SOC and quantity of stubble addition ($r = 0.85$), carbon addition by stubble ($r = 0.94$), and carbon stock ($r = 0.91^{**}$). Similar to this, the MBC-SOC connection ($r = 0.83^{**}$), which supported the contribution of SOC to collective formation.

Conclusion: The strong positive link between SOC and MBC suggests that adding carbon helps to improve soil health, and rice straw is an excellent source of carbon. It needs to be suppressed in the soil.

1. INTRODUCTION

Rice-rice cropping system is a popular system prevailing in Odisha occupying about 16% of total rice area in Odisha (Agricultural statistics Odisha, 2013). By default at the time of harvest farmers cut paddy plant leaving (20-30) cm above ground stubble which is incorporated in soil act as good source of carbon addition. The amount of soil organic carbon is affected by manuring and fertiliser. Under inorganic fertilisation, SOC concentration rises, especially when using inorganic (N) fertiliser (Majumdar et al., 2008;

Reid 2008; Tong et al., 2009; Batle- Bayer et al. 2010). The soil's agricultural leftovers are improved by inorganic N fertilisers, which raises SOC. To counteract the potential rise in C from crop leftovers, N fertiliser has been demonstrated to improve the breakdown of crop residues and soil carbon (Nayak et al, 2009). Soil carbon stock is a function of soil organic carbon, bulk density, and plough horizon depth. Organic inputs and rice residues (rice stubble and root) are the main sources of converted soil carbon in paddy field ecosystems (Li Z.P. et al., 2010).

To combat the deterioration of soil structure, various organic additions, such as manure (farmyard manure, green manure), compost, and crop residues (especially agricultural residues), are typically recommended. When used, these additions are dispersed in various SOC pools (Majumder et al., 2008). According to Bhattacharyya et al. (2000), SOC stock may be the only metric that can be used to determine where to focus efforts on restoring soil health and reducing climate change. Direct sources of soil carbon include the development and decomposition of plant roots, as well as the movement of carbon-rich chemicals from the roots to soil microorganisms (Ontl and Schulte, 2012). For agriculture to be long-term sustainable, soil organic carbon must be maintained because diminishing levels typically result in lower crop productivity (Allison, 1973). Applications of manure containing NPK enhance the SOC content or produce significantly higher SOC levels (Swarup et al., 2000). Microbial biomass has been found to be a sensitive indicator of changes in soil biology brought on by management (Carter, 1986; Doran, 1987; Powlson et al., 1987). When employed to assess soil organic matter dynamics, Sparling (1992) found that the soil organic C ratio and microbial biomass C were more sensitive indices than total soil organic C. The ratio of microbial biomass carbon to total organic carbon in the control and fertilised treatments, according to Rochette and Gregorich (1998), was around 1.5% and 3.5%, respectively. The higher proportion of total organic C as microbial biomass C in manured soil suggests that the C levels in soil receiving manure will rise if the microbial biomass C is utilised as an early indicator of altering soil organic matter contents.

Hu and Cao (2007) conducted a field experiment, where microorganism growth in soil carbon, microbial respiration and soil enzyme activity were estimated from six different manure treatments. The experimental results showed that soil microbial biomass C, microbial respiration rate and enzyme activity in compost system were significantly higher than in the chemical fertilizer and control treatment. Dubey *et al.*, (2009) reported that microbial biomass carbon was more in wheat straw-treated treatments followed by FYM and green manure-treated treatments respectively. The highest biomass carbon in wheat straw-treated treatments may be attributed to the fact that the total as well as individual microbial population was the highest in wheat straw-treated treatment followed by FYM and green manure-treated treatment. Microbial biomass C was higher at surface than sub-surface soil. According to a study by Mandal

et al. (2007), a stable application of NPK + FYM resulted in the highest levels of microbial biomass carbon (MBC), mineralizable nitrogen (MBN), dehydrogenase and phosphate activities in soil, and the lowest values at the control. As a result of the use of nitrogen and phosphorus-containing inorganic fertilisers, the MBC and MBN levels rose. Yet, when organic manure was put with inorganic fertilisers, these metrics increased more. Application of organic manure has also dramatically boosted the activity of the enzymes soil dehydrogenase, alkaline phosphatases, B-glucosidase, and urease (Liu Enke et al, 2010). Bhubaneswar soil is lateritic in origin, light textured, acidic and contains low carbon (Nayak *et al.*, 2002). The hot and sub humid climatic condition with high monsoon rainfall causes the annual depletion of carbon. There is a scope to capture atmospheric carbon by different manuring practices through stubble in rice-rice cropping system. In Bhubaneswar soil under rice-rice cropping system, stubble is incorporated into soil. But information on how much of C is added in different type of manuring through stubble is not available. Our objective of study is quantification of incorporated stubble in soil by different manurial treatments, C added through stubble in different treatments, SOC and its effect on soil biological properties.

2. MATERIALS AND METHOD

The long-term fertilizer experiment at Central Research Station, OUAT, Bhubaneswar Odisha at (20°16' - 20°17' N latitude and 85°49'E longitude) was selected for this study. The experiment was started in 1994 on Inceptisol (710 g kg⁻¹ sand, 120 g kg⁻¹ silt and 170 g kg⁻¹ clay). The initial pH (1:2) was 5.8, electrical conductivity 0.12 dS m⁻¹, SOC: 4.3 g kg⁻¹, CEC: 3.75 c mol (P⁺) kg⁻¹, Alkaline KMnO₄ available N: 187 kg ha⁻¹, NaHCO₃ extractable P: 19.4 kg ha⁻¹, NH₄OAC extractable K: 93.4 kg ha⁻¹ content of soil. The field experiment was laid out in randomized block design with four replication with treatments viz., T1(100% NPK), T2 (100%NPK +FYM), T3 (100% N) , T4 (100% NP), T5 (100%NPK +Lime), T6 (Control- no manure and fertilizer). 100% recommended dose of fertilizer 80 kg N, 40 kg P₂O₅, 60 kg K₂O was applied each year per season. Nitrogen as (Urea) was applied in three splits i.e., 25% basal, 50% at maximum tillering (MT) stage and 25% at panicle initiation (PI) stage. Total phosphorus as basal as DAP and potassium as MOP in two splits 50% as basal and 50% at panicle initiation stage was applied. Manure in the form of FYM @ 5 t ha⁻¹ and lime@ 1 t ha⁻¹ was applied per cropping season. Manure with chemical composition and C: N ratio of FYM is given in Chart no.1

Chart 1. Chemical composition and C: N ratio of FYM used during rabi season.

Properties	Content
Carbon (%)	20.8

N (%)	0.79
P ₂ O ₅ (%)	0.50
K ₂ O (%)	0.80
C: N value	26.28

Soil Sampling

Pre-rabi soil samples were taken in December 2013 and post-rabi soil samples were taken in April 2014. SOC (MBC) was measured using composite surface soil samples from each treatment that were collected at a depth of 0 to 15 cm, air dried, and then put through a 2 mm filter. For the purpose of estimating bulk density, soil cores from 0 to 15 cm of soil depth were obtained.

Collection and processing of stubble

1 square meter metallic frame of 10 cm height was randomly installed in the treatment plot, pressed to (1-2) cm, water was added to make the soil wet, soft and loose. Next day morning stubble was uprooted carefully so that loss of rootlets will be negligible. Root was washed with water, treatment wise stubble was made bunches and tagged, dried for a week under sun then oven dried. Stubble (both shoot and root) were chopped and used for carbon estimation in muffle furnace.

Soil Analysis

The bulk density of soil samples was determined by applying the core method, which Black (1965) described. The wet oxidation method devised by Walkley and Black was utilised to assess the soil's organic carbon, and the Vance et al. (1987) reported fumigation and extraction procedure was employed to estimate the carbon in the soil's microbial biomass.

Plant Analysis

For determining the organic content of root & shoot of recognisable water activity were weighed in a dry silica crucible. Heat was given slowly by raising the temperature setting in 3 steps (100, 400 & 550 °C). The final temperature setting was at 550 °C maintained for 8 hrs. Then crucible containing ash was cooled & weighed. Percentage ash and organic matter were calculated by difference between crucibles before & after combustion as follows:

$$\text{Ash (\%)} = \frac{W_3 - W_1}{W_2 - W_1} \times 100$$

And organic matter (%) = 100 – ash %

(Where W_1 = The weight of empty crucible, W_2 = The weight of dry crucible containing root & shoot, W_3 = The weight of dry crucible containing root & shoot after following ignition).

Statistical analysis

According to the conventional statistical technique established by Gomez and Gomez (1976), data about soil qualities for various treatments were subjected to analysis of variance, and the link between sets of variables was determined by performing linear regressions.

3. RESULTS AND DISCUSSION

Addition of quantity of rice stubble (20 cm) in soil

A comparison of stubble addition by different treatments stated that 100% NPK + FYM treatment helped in adding 111.10% more over control followed by 63.96 %, 51.39 %, 30.16 %, 21.55 % in 100% NPK + Lime, 100% NPK, 100%NP and 100% N treatment respectively (Table 1). Similarly, comparison of quantity of addition by different treatment over RDF it was found that in 100% NPK +FYM 39.43 % and in 100% NPK + Lime 8.29 % which was positive and in other treatments carbon is added in a drastic reduced quantity of -14.02 %,-19.70 %, -33.94 % in100%NP ,100% N and control treatment. Always in control treatment addition of stubble was the minimum irrespective of season and year. The quantity of stubble in differential treatment was maximum in 100% NPK + FYM followed by 100% NPK + Lime, 100% NPK, 100% NP, 100 %N, control. This trend of quantity of stubble added was due to the differential biomass production in their respective treatments.

Table 1. Quantity of rice stubble (20 cm) added (kg/ha) in different manurial treatments.

Treatment	<i>Kharif</i> 2012	<i>Rabi</i> 2012-13	<i>Kharif</i> 2013	<i>Rabi</i> 2013-14	Avg/yr	Increase over control (%)	Increase over RDF (%)
100% NPK	2172.5	2452.5	2045.3	2410.3	4540.3	51.39	-
100% NPK + FYM	3086.5	3355.2	2872.5	3347.5	6330.85	111.10	39.43
100% N	1768.4	2002.5	1542.5	1977.5	3645.45	21.55	-19.70
100%NP	1911.2	2097.5	1782.5	2015.6	3903.4	30.16	-14.02

100% NPK + Lime	2522.6	2596.4	2302.5	2412.5	4917	63.96	8.29
Control	1489.1	1597.5	1380.6	1530.6	2998.9	-	-33.94
SE(m)±	191.93	98.05	28.73	88.97			
CD(0.05)	578.45	295.52	86.58	268.15			

Addition of carbon in soil through rice stubble

A comparison of carbon addition by different treatments stated that 100% NPK +FYM treatment helped in adding 110.17% more over control followed by 65.50 %, 50.42%, 38.12%, 28.93% in 100% NPK + Lime, 100% NPK, 100%NP and 100% N treatment respectively (Table 2). Similarly comparison of carbon addition by different treatment over RDF it was found that in 100% NPK +FYM 39.71% and in 100% NPK + Lime 10.02% which was positive and in other treatments carbon is added in a drastic reduced quantity of -8.17%, -14.28%,-33.52% in 100%NP ,100% N and control treatment. The quantity of carbon added through rice stubble including root was maximum in 100% NPK + FYM followed by 100% NPK + Lime, 100% NPK, 100% NP, 100%N, control. This trend of quantity of 'C' added was due to the multiple quantity of stubble incorporated in the respective treatments (Holeplass *et al.*, 2004). Due to differential biomass production, variable quantity of stubbles and carbon were added in their respective treatments.

Table 2. Effect of long-term manuring on addition of carbon in soil through rice stubble (kg/ha)

Treatment	Kharif 2012	Rabi 2012-13	Kharif 2013	Rabi 2013-14	Avg/yr	Increase over control (%)	Increase over RDF (%)
100% NPK	866	981	858	976	1840.5	50.42	-
100% NPK + FYM	1233	1342	1229	1339	2571.5	110.17	39.71
100% N	706	881	697	871	1577.5	28.93	-14.28
100%NP	762	939	753	926	1690	38.12	-8.17
100% NPK +Lime	1008	1024	1001	1017	2025	65.50	10.02
Control	592	639	584	632	1223.5	-	-33.52

SE(m)±	28.08	57.91	25.32	28.37			
CD(0.05)	84.66	174.57	76.33	85.52			

Carbon stock

Organic carbon stock was calculated as per the formula $[C_{\text{stock}} = \text{SOC} (\%) \times \text{BD} (\text{Mg m}^{-3} \times \text{soil depth (m)} \times 100)]$. Data showed that it varied from 7.41 to 12.50 Mg ha^{-1} and 7.14 to 11.76 Mg ha^{-1} of (0-15) cm soil depth post-harvest soils of *kharif* and *rabi* season respectively (Table 3). The minimum SOC stock was observed in control where as maximum was found out in 100% NPK+ FYM treatment followed by 100% NPK + Lime. The SOC stock value was in order of control < 100% N < 100% NPK < 100% NP < 100% NPK + Lime < 100% NPK+ FYM. After continuous 7 cropping cycles a good amount of organic matter in the form of in situ stubble and external FYM were added in the treatment resulting in significantly higher quantity of soil organic carbon stock than any other treatment. This type of findings corroborated using the results of (Holeplass et al., 2004 and Jiang et al., 2006).

Table 3. Effect of long term manuring on Carbon stock (Mg ha^{-1}) of soils after harvest of *Kharif* and *Rabi* crops.

Treatment	(0-15)cm	
	<i>Kharif</i>	<i>Rabi</i>
100% NPK	10.57	9.37
100% NPK+ FYM	12.50	11.76
100% N	10.11	9.14
100%NP	10.25	9.30
100% NPK+ Lime	10.92	9.80
Control	7.41	7.14
SE(m)±	0.44	0.43
CD(0.05)	1.31	1.30

Soil Organic Carbon

SOC value varied from 3.48 to 6.51(g/kg) and 3.35 to 6.13 (g/kg) of (0-15) cm soil during *kharif* and *rabi* respectively. Similarly, in (15-30) cm of depth SOC value for above season varied from 1.65 to 4.65 g/kg and 1.38 to 4.32 g/kg respectively. SOC value of surface soil was found to be higher than subsurface soil in respective treatments irrespective of seasons. The minimum SOC was observed in control where as maximum SOC was found out in 100% NPK+ FYM treatment followed by 100% NPK+ Lime treatment. The SOC value was in the order of control < 100% N < 100% NP< 100% NPK < 100% NPK + Lime < 100% NPK+ FYM.SOC value was found maximum in NPK +FYM treated plot followed by NPK + Lime , 100% NPK, 100% NP, 100% N control in decreasing order (Table 4). The significantly greater SOC in the NPK+ FYM treatment was caused by both the addition of more stubble and FYM. In their research on agricultural soils, they reported identical outcomes that have been amply documented by (Rudrappa et al., 2006). The amount of organic carbon in soil was thought to depend on the farming system's net input of organic residues (Gregorich et al., 1996). The findings made it abundantly evident that the carbon store was higher in *kharif* than in *rabi* due to more SOC, a cooler environment, and more stubble addition.

Table 4. Effect of long-term manuring on SOC (g/kg) of soils after harvest of *Kharif* and *Rabi* crops.

Treatment	(0-15)cm	
	<i>Kharif</i>	<i>Rabi</i>
100% NPK	5.34	4.73
100% NPK+ FYM	6.51	6.13
100% N	4.92	4.45
100%NP	5.10	4.63
100% NPK+ Lime	5.6	5.03
Control	3.48	3.35
SE(m)±	0.20	0.23
CD(0.05)	0.60	0.70
Initial	4.3	

Bulk Density (BD)

The BD varied from 1.28 to 1.42 Mg m⁻³ in (0-15) cm, less than the initial BD (1.55 Mg m⁻³) and 1.32 to 1.49 Mg m⁻³ in (15-30) cm of soil (Table 5). Addition of stubble, FYM and inorganic fertilizer helped in reducing the BD. Imbalanced or no fertilization increase the BD due to comparatively less stubble addition and compaction. The BD value of subsurface soil is higher than the surface soil irrespective of treatments. (K.M. Hati et al.,2006)

Table 5. Effect of long-term manuring on BD (Mg m^{-3}) of soils after harvest of *Kharif* crops

Treatment	(0-15) cm
100% NPK	1.32
100% NPK+ FYM	1.28
100% N	1.37
100%NP	1.34
100% NPK+ Lime	1.30
Control	1.42
SE(m) \pm	0.04
CD(0.05)	0.11
Initial	1.55

Microbial biomass carbon (MBC)

The data revealed that the value of MBC varied from 111.21 to 204.93 $\mu\text{g of C g}^{-1}$ and 45.89 to 132.41 $\mu\text{g of C g}^{-1}$ (0-15) cm soil in *kharif* and *rabi* respectively (Table 6). Similarly, in (15-30) cm depth the value of MBC for above seasons varied from 44.32 to 93.79 $\mu\text{g of C g}^{-1}$ and 32.18 to 81.59 $\mu\text{g of C g}^{-1}$ respectively. Regardless of the season, the MBC value of surface soil was found to be higher than subsurface soil in the corresponding treatments. The control had the lowest MBC value, whereas 100% NPK+ FYM therapy had the highest MBC value, followed by 100% NPK + Lime. The MBC value fell into the following categories: 100% N, 100% NP, 100% NPK, 100% NPK + Lime, and 100% NPK+ FYM. The agro-nutrient ecosystem's cycling depends greatly on the microbial biomass. The amount of microbial biomass in a soil is significantly influenced by soil management practises. In the current experiment, a number of long-term manuring practises had a considerable impact on soil MBC (Table 6). The MBC was low in the control and unbalanced fertiliser treatments, but it drastically rose after applying manure and the proper quantity of NPK. Similar findings have been reported by a number of other researchers (Goyal et al., 1992; Chakrabarti et al., 2000; Kaur et al., 2005). The rapidly metabolizable carbon and nitrogen in organic manure, as well as growing root biomass and root exudates as a result of improved crop development, are the main contributors to the biomass rise.

Table 6. Effect of long-term manuring on MBC ($\mu\text{g of C g}^{-1}$) of soils after harvest of *Kharif* and *Rabi* crops.

Treatment	(0-15) cm	
	<i>Kharif</i>	<i>Rabi</i>
100% NPK	166.86	86.76

100% NPK+ FYM	204.93	132.41
100% N	126.65	55.39
100%NP	156.49	71.88
100% NPK+ Lime	184.51	100.66
Control	111.21	45.89
SE(m)±	0.81	0.54
CD(0.05)	2.46	1.68

Correlation between SOC with addition of stubble, carbon added, carbon stock.

Results on Correlation study of SOC with C enhanced by stubble, C stock of soil (Fig. 1, 2, 3, 4 ,5,6) was found to be significantly high. Correlation between SOC (*Kharif* 2013) with addition of stubble (*Rabi* 2012-13), SOC (*Rabi* 2013-14) with addition of stubble (*Kharif* 2013) was found to be ($r = 0.80^{**}$), ($r = 0.85^{**}$) respectively. This showed contribution of stubble in SOC was more in *Rabi* 2013-14 than that of *Rabi* 2012-13 and between SOC (*Kharif* 2013) with carbon added through stubble (*Rabi* 2012-13) and SOC (*Rabi* 2013-14) with carbon added through stubble (*Kharif* 2013-14) found to be ($r = 0.94^{**}$) and ($r = 0.85^{**}$) respectively. A significant interactions across SOC (*kharif*) with carbon stock in *Kharif* and SOC (*rabi*) with carbon stock in *rabi* were ($r = 0.88^{**}$) and ($r = 0.91^{**}$) respectively. A substantial relationship between SOC and MBC ($r = 0.83^{**}$) (Fig.6) were found indicating role of stubble incorporation in elevated SOC, MBC.




Fig.1. Correlation between SOC (*Kharif* 2013) and addition of stubble (*Rabi* 2012-13)




Fig.2 Correlation between SOC (*Rabi* 2013-14) and addition of stubble (*Kharif* 2013)




Fig.3. Correlation between SOC (*Rabi* 2013-14) with carbon addition through stubble (*Kharif* 2013)




Fig.4 Correlation between SOC and carbon stock in *Kharif*




Fig.5 Correlation between SOC and carbon stock in *Rabi*




Fig.6 Correlation between Soil organic carbon and Microbial biomass carbon

4. CONCLUSION

In a long-term manuring experimental field under rice-rice cropping system at Bhubaneswar soil seasonally added stubble enhanced the carbon content in stubble, carbon stock in soil and soil organic carbon. The microbial population and activity of microorganisms in soil were enhanced which was indicated by higher MBC where more stubble was added. Increased soil carbon stock, SOC, and MBC were found in 100% NPK+ FYM, which was proceeded in decreasing order by 100% NPK + Lime, 100% NPK, 100% NP, 100% N, and control.

From the entire study following conclusions can be drawn:

1. Balance fertilization is good, balanced fertilizer with lime treatment in acid soil is better but integration of organic with inorganic is the best way of manuring in a light textured, low fertile acid soil at Bhubaneswar in rice-rice cropping system.
2. Recommend dose of fertilizer was better over imbalanced or no manuring where less stubble was produced and incorporated.
3. For the soil, rice straw is a good source of carbon. It has a positive effect on the condition of the soil, improving rice quality and yield. Soil has to be amended with stubble.

REFERENCES

- Allison FE. Soil Organic Matter and its Role in Crop Production. *Soil Science*, Elsevier, Amsterdam. 1973. p.3
- Bhattacharya T, Pal DK, Mandal C and Velayutham M. Organic carbon stock in Indian Soils and their geographical distribution. *Curr Scien.* 2006;79(5):655-660.
- Carter MR. Microbial biomass carbon as an index for tillage induced changes in soil biological properties. *Soil Til Res.* 1986;7:29-40.
- Doran J and Smith MS. 1987. Organic Matter Management and utilization of soil and fertilizer nutrients in Folett RF et al.,eds. Soil fertility and organic matter as critical component of systems. ASA Spec. Publ.19 ASA & SSSA Madison, WI.
- Dubey YP, Bedi P and Datt N. 2009. Microbial properties under rice-wheat cropping sequence in an acid *alfisol*. *Journal of Indian Society of Soil Science*, **57**(3):373-377.
- Goyal SM, Mishra M, Hooda IS and Singh R. 1992. Organic matter microbial biomass relationships in field experiments under tropical conditions: effect

- of inorganic fertilization and organic amendments. *Soil Biology and Biochemistry* **24**:1081-1084.
- Holeplass H, Singh BR and Lal R .2004. Carbon sequestration in soil aggregates under different crop rotation and nitrogen fertilization in an *Inceptisols* in south eastern Norway. *Nutr Cycl Agroecosyst.* **70**:167-177.
- Hu C and Cao Z.2007. Size and activity of soil microbial biomass and soil enzyme activity in long term field experiments, *World Journal of Agricultural Science.***1**:63-70.
- Jiang D, Hengsdijk H, Dai TB, de Boer W, Jiang Q and Cao WX. 2006. Long term effects of manure and inorganic fertilizers on yield and soil fertility for a winter –maize system in Jiangsu, China. *Pedosphere.* **16**(1):25.32.
- Kaur K, Kapoor KK and Gupta AP.2005. Impact of organic manures with and without mineral fertilizers on soil chemical and biological properties under tropical conditions. *Journal of Plant Nutrition and Soil Science* **168**:117-122.
- Li ZP,Liu M, Wu XC, Han FX, Zhang TL .2010. Effects of long-term technical fertilization and organic amendments on dynamics of soil organic C and total N in paddy soil derived from barren land in subtropical China. *Soil& Tillage Reseach.***106**:268-274
- Liu Enke, Yan C, Mei X, He W, Bing SH., Ding L, Liu Q, Liu S and Fan T (2010) *J.Geoderma.***158**:173-180.
- Majumdar B, Mandal B, Bandyopadhyaya PK, Gangopadhaya A, Mani PK, Kundu AL and Mazumdar D. 2008. Organic amendments influence soil organic carbon pools and rice-wheat productivity. *Soil Science Society of America Journal.***72**:775-785.

- Mandal, A;Patra, A.K.;Singh, D;Swarup, A and Masto, R.E. (2007) Effect of long-term application of manure and fertilizer on biological and biochemical activities in soil during crop development stages.*J.Bioresource Technology*. **98**: 112-117.
- Nayak P, Patel D, Ramakrishnan B, Mishra AK and Samantaray RN. 2009. Long term application effects of chemical fertilizers and compost on soil carbon under intensive rice cultivation. *Nutrient Cycling in Agroecosystems*.**83**:259-269.
- Nayak RK, Sahu GC and Nanda SSK.2002. Characterisation and classification of soils of Central rice research station, O.U.A.T, Bhubaneswar. *Journal of Ageo Pedology* **7**(2):27-31.
- Ontl TA & Schulte IA.2012. "C" storage. *Nature education knowledge*.**3**(10):35.
- Rochette P and Gregorich EG. 1998. Dynamics of soil microbial biomass carbon C, soluble organic C and CO₂ evolution after 3 years of manure application. *Canadian Journal of Soil Science Society*.**78**(2):283-292.
- Rudrappa L, Purakayastha TJ, Dhyan S and Bhadrarary S. 2005. Long term manuring and fertilization effects on soil carbon pools in a Typic Haplustept of semi arid sub- tropical India. *Soil & Tillage Research*. **88**:180-192.
- Sparling GP.1992. Ratio of microbial biomass to soil organic carbon as a sensitive indicator of changes in soil organic matter. *Australian Journal of Soil Research*.**30**:195-207.
- Swarup A, Manna MC and Singh GB. 2000. Impact of land use management practices on organic dynamics in soils of India in Global Climate change and Tropical Ecosystem (Lal et al., eds). *Advances in Soil Sci.(USA)*.PP.261-281.

Vance ED, Brookers PC and Jenkison DS.1987. An extraction method for measuring soil microbial biomass carbon. *Soil Biology and Biochemistry*.**19**:703-706

UNDER PEER REVIEW