

ORIGINAL RESEARCH ARTICLE

TAXONOMIC SIGNIFICANCE OF FOLIAR AND STEM ANATOMY IN THE DELIMITATION OF *PIPTADENIASTRUM AFRICANUM* (HOOK.F.) BRENAN AND *CATHORMION ALTISSIMUM* (HOOK.F.) HUTCH. & DANDY

Authors Contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript

ABSTRACT

The current monotypic status of the genus *Piptadeniastrum* Brenan is reasonably challenged as *Cathormion altissimum* is claimed to be closely related to *Piptadeniastrum africanum* – the only species in the genus – by some indigenous people of Bayelsa State. This work is a systematic examination of this claim by studying some aspects of the foliar and stem anatomy of *P. africanum* and *C. altissimum*. Leaf peels and stem sections were made and studied according to standard procedures. Anatomical characteristics considered include stomata distribution, stomatal Index, type of stomata, the shape of leaf epidermal cells and anticlinal walls, the outline of the stem, and the arrangement of vascular bundles. Results showed that stomatal distribution in both plants was hypostomatic. Stomatal index and stomata type were 21.5% and mainly anomocytic respectively, for *P. africanum* and 14.4% and mainly paracytic, respectively, for *C. altissimum*. Epidermal cell shape is irregular with undulating anticlinal walls in *P. africanum*, while epidermal cell shape is rectangular to polygonal with curved or straight anticlinal walls in *C. altissimum*. The outline of the stem is angular with four pronounced ridges and furrows in *P. africanum* while stem outline is wavy with very mild protrusions in *C. altissimum*. The differences in anatomical characteristics may be justifying the placement of these plants in different genera. *Piptadeniastrum* Brenan is monotypic.

Keywords: *Piptadeniastrum africanum*, *Cathormion altissimum*, anatomical character, anticlinal walls, paracytic, hypostomatic, stomatal index

INTRODUCTION

The species *P. africanum* is a forest tree of the rain forest, published and widely accepted as the only species in the genus *Piptadeniastrum* Brenan, a genus that belongs to the family

Fabaceae Lindsl, Subfamily Mimosoideae [1,2,3]. *P. africanum* (Hook.f.) Brenan is endemic to tropical Africa and occurs in the Niger Delta, especially on riverbanks in the riverine areas of the rainforest [4,5,6]. This forest tree is of great interest due to its wide range of ethnobotanical uses – medicinal, domestic, social, and agricultural applications. Thus, studies have concentrated on its chemical constituents and their usefulness to man. Consequently, there are comparatively few taxonomic studies that are necessary to clear the doubts that surround its monotypic taxonomic status. This research gap is evident because names such as *Piptadeniastrum manni* Oliver, with an unresolved status and low confidence level, were placed in the Plant List [7] and later placed with an accepted status and medium confidence level [8], but placed as a species of the genus *Piptadeniastrum* Brenan in TROPICOS [9]. Moreover, reports of varieties of *P. africanum* such as *Piptadeniastrum africanum* var. *africanum* and *Piptadeniastrum africanum* var. *brachysperma* Pellergr. exist in the literature [8].

Similarly, the identification, nomenclature, and taxonomic status of this species (whether it is the only species in the genus *Piptadeniastrum* Brenan) among some indigenous people in Bayelsa State is unclear. While most of the local people confidently affirm that there are two types of plants, they disagree on the forms in which they occur in the wild. A survey in the study area showed that one of the trees closely related to *P. africanum* is *C. altissimum* [10]. Also of the Fabaceae family, *C. altissimum* is of great ethnobotanical importance. It is used as food spice [11], medicine [12, 13], and wood of choice for carving durable, quality kitchen utensils like mortar, pestle, and spatula that are indispensable in a typical Niger Delta family. These carved utensils serve as valuable traditional gifts to new couples during marriage ceremonies.

Observation is a critical element of the scientific method. The indigenous people, who are close to the forest and have depended on it for their sustenance over the years, have significant ethnobotanical knowledge about these trees, and the importance of applying appropriate scientific methods to study their observations cannot be overemphasized. The fact that indigenous knowledge, sometimes called Traditional Ecological Knowledge (TEK), has contributed immensely to discoveries in natural sciences and medically necessary substances in plants cannot be controverted [14,15,16]. Moreover, when the identity of even a single plant is not sure and authentic from broad-based, empirical taxonomic evidence, then extant erroneous phylogeny in the relevant taxa is undeniable. This study, therefore, is aimed at describing some leaf epidermal and stem anatomical characteristics that may be significant, at least in serving as supportive taxonomic evidence, in the description and delimitation of *P.*

africanum and *C. altissimum* as either belonging to different genera (as they are present) or the possibility of belonging to the same genus and thus establishing their degree of relatedness.

MATERIALS AND METHODS

Study area

Bayelsa State, where this study was carried out, comprises eight Local Government Areas (Figure 1). The state is located within latitudes $04^{\circ}15'$ North and $05^{\circ}23'$ south and longitudes $05^{\circ}22'$ West and $06^{\circ}45'$ East. It is bounded by Delta State on the North, Rivers State on the East, and on the west and south by the Atlantic Ocean (Figure 1).




Fig. 1: Bayelsa State, showing the Local Government Areas (Inset: Map of Nigeria showing Study Area)

Sample collection and identification:

Matured samples of the plants (Figure 2) were collected from the study area and identified at the Forest Herbarium Ibadan of the Forestry Research Institute of Nigeria (FRIN). Voucher specimens of the plants – *P. africanum* (FHI 110137) and *C. altissimum* (110884) – were deposited in the herbarium for reference and further studies.




Fig. 2(a-b): (a) *C. altissimum* (b) *P. africanum*

Leaf epidermal studies

Cuttings from identical regions were made from fresh leaves of the plants collected from the field. Peels from the adaxial and abaxial surfaces of the leaves were obtained carefully with the aid of pins and forceps. Samples were stained with alcian blue or 1% safranin, rinsed in distilled water to remove excess stain, and mounted on microscope slides with a drop of glycerol. Epidermal features such as the shape of epidermal cells, type of anticlinal wall, stomatal distribution, stomata type, and stomatal index were studied. Stomatal index (SI) was estimated as reported by Obembe [17], while the terminology for stomata type was taken after Metcalfe and Chalk [18]

Stem anatomical studies


Matured plants from fresh vegetation were used for this study. Cut sections of the stem were fixed at room temperature (23⁰C – 28⁰C) in Formaldehyde-acetic acid- alcohol (1 part of 40% formaldehyde, 1 part of acetic acid, 18 parts of 70% ethanol) for at least 48 hours. Then, samples were washed with several changes of distilled water, dehydrated through 2-hour incubations in graded alcohol solutions (30%, 50%, 70%, and 100%) at room temperature, and embedded in wax. Hand sections were cut with blades. Thin sections were selected and dewaxed by passing the specimens through a series of 3 hours of incubations in graded ethanol: chloroform solutions (1:3, 1:1, and 3:1), with a final 3-hour incubation in absolute ethanol. Samples were stained with alcian blue, counterstained with 1% safranin, and mounted on microscope slides. Photomicrographs were acquired with a digital camera.

RESULTS

Leaf epidermal studies: Results of the leaf epidermal studies of the plants are presented in table 1 and figure 3.

Table 1: Leaf epidermal characteristics of *P. africanum* and *C. altissimum*

Characters/ Species	Stomatal Index (%)	Stomata Type	Stomata distribution	Epidermal cell shape	Anticlinal wall
<i>P. africanum</i>	21.5	Mainly anomocytic	Hypostomatic	Irregular	Undulating
<i>C. altissimum</i>	14.4	Mainly paracytic	Hypostomatic	Rectangular/ polygonal	Curved/ straight



A: Adaxial/Upper epidermis of *P. africanum* B: Adaxial/Upper epidermis of *C. altissimum*

Figure 3: Adaxial and Abaxial leaf epidermis of *P. africanum* and *C. altissimum*

Stem anatomical studies: Result of anatomical studies of the stems of *P. africanum* and *C. altissimum* is shown in Figure 4.




Figure 4: Cross-section of stems of *P. africanum* and *C. altissimum*

DISCUSSION

Stomatal distribution in both the leaves of *P. africanum* and *C. altissimum* is Hypostomatic i.e. stomata are found only on the abaxial surfaces of the leaves. The epidermal cell shape is irregular with undulating anticlinal walls in *P. africanum*, while the epidermal cell shape for *C. altissimum* is rectangular/polygonal with curved/straight anticlinal walls. The stomatal index is 21.5% in *P. africanum* with mainly anomocytic stomata type, while the stomatal Index for *C. altissimum* is 14.4% with mainly paracytic stomata.

The taxonomical value of foliar epidermal characters such as hairs, stomata, and epidermal cells has been well established. Though epidermal cells vary considerably in size, shape, and outline in different plants, they are affected by environmental factors. Consequently, the shape of epidermal cells should be helpful as confirmatory evidence [18]. However, variations in leaf epidermal characters have been used in the identification and classification of several plant groups because they are primarily controlled by genes and are structurally diverse [19,20,21,22]. Nurul-Aini *et al* [23] showed that leaf micromorphological features such as stomata characteristics and the presence of trichomes were taxonomically significant in some selected taxa of *Acanthaceae*. Researchers have found leaf epidermal micromorphological characteristics such as differences in epidermal cell shape and size and

the stomatal index, and the pattern of anticlinal walls of epidermal cells to be of taxonomic significance among 43 species of *Allium* [24]. It was reported that the pattern of the anticlinal walls of epidermal cells was consistent within a species, but there were differences between species. Consequently, micromorphological characteristics of the epidermis of *P. africanum* and *C. altissimum* in this study were of taxonomic significance in their delimitation into different genera.

The stem of *P. africanum* has an angular outline with four pronounced ridges and furrows, while the stem of *C. altissimum* has a wavy outline. Furthermore, although the vascular bundles of both plants are scattered in the ground tissue, they differ in the patterns of grouping and arrangement. In *C. altissimum*, the vascular bundles are arranged in groups of about five, while in *P. africanum*, vascular bundles are concentrated at the ridges or protrusions. These anatomical characteristics may be significant in the delimitation of these plants.

The significance of using stem anatomical features in collaboration with other lines of evidence in taxonomy is well known [25,26,27,28,29]. Cavente et al. [30] worked on six species of *Rhipsalis* and found that features of the stem anatomy discriminate between the species. Furthermore, Kaplan & Symoens [31] used stem anatomy to distinguish between two confusing African taxa (*Potamogeten schweinfurthii* and *Potamogeten richardii*).

CONCLUSION

The results of the studies showed no sufficient degree of relationship that could warrant the placement of *C. altissimum* under the genus *Piptadeniastrum* Brenan. Consequently, the current monotypic status of the genus *Piptadeniastrum* Brenan is maintained.

SIGNIFICANCE STATEMENT

The nomenclatural conflict between *P. africanum* and *C. altissimum*, arising from the similar morphology of the leaves, was systematically resolved using evidence from foliar and stem anatomy. Furthermore, the information from the results of this study has enriched the existing literature on the taxonomy of plants.

CONSENT AND ETHICAL APPROVAL

It is not applicable.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Airy Shaw HK. J.C. Willis: *A dictionary of flowering plants and ferns* 8th ed. Revised. Cambridge: Cambridge University Press. 1985
2. Brusotti G, Tosi S, Tava A, Picco AM, Grisoli P, Cesari I, Caccialanza G. Antimicrobial and phytochemical properties of stem bark extracts from *Piptadeniastrum africanum* (Hook f.) Brenan. *Industrial Crops and Products*. 2013; 43:612-616
3. Gunn CR. Fruits and Seeds of Genera in the Subfamily Mimosoideae (Fabaceae). US Department of Agriculture, Technical Bulletin. 1984; 1681
4. Brenan JPM. *Piptadeniastrum africanum*. *Kew Bull.* 1955; 179.
5. Hutchinson J, Dalziel JM. *Flora of West Tropical Africa*. (2nd ed.). Vol. 1, Part 2. Revised by Keay RWJ; 1958.
6. Nyananyo BL. *Plants from the Niger Delta*. Nigeria; 2006
7. The Plant List. *Piptadeniastrum*. 2010; Accessed 19 May 2013. Available: <https://www.theplantlist.org/>
8. The Plant List. *Piptadeniastrum*. 2013; Accessed 19 May 2013. Available: <https://www.theplantlist.org/>
9. TROPICOS. *Piptadeniastrum manni* Oliv. 2013; Missouri Botanical Garden. Accessed 19 May 2013. Available: <https://www.tropicos.org/Name/13069459>.
10. Youkparigha FO, Nyananyo BL, Oyedeji AA. Morphology and Distribution of *Piptadeniastrum africanum* (Hook.f.) Brenan and *Cathormion altissimum* (Hook.f.) Hutch. & Dandy in Bayelsa State, Nigeria. *International Journal of Advanced Research in Botany*. 2019; 5(2): 8-14.
11. Jolaoso AA, Ajayi JO, Ogunmuyiwa JO, Albert OM. Changes in Functional Properties as a Measure of Biochemical Deterioration of Oso (fermented seeds of *Cathormion altissimum*). *Journal of Emerging Trends in Engineering and Applied Sciences*. 2012; 3(4): 608-613.
12. Burkill HM. *The Useful Plants of West Tropical Africa*. Edn. 2, Vol. 3, Families J-I. Royal Botanic Gardens, Kew, Richmond, Surrey, England. 1985
13. Lemmens RHMJ. *Albizia altissima* Hook.f. Record for PROTA4U. Louppe, D. Oteng-Amoako, A.A. and Brink, M. (Editors). PROTA (Plant Resources for Tropical Africa), Wageningen, Netherlands. 2006. Available: <https://www.prota4u.org/search.asp>.
14. Brush SB. Indigenous knowledge of biological resources and Intellectual Property Rights: The role of Anthropology. *American Anthropologist*. 1993; 95(3): 653-671
15. Reyes-Garcia V. The relevance of traditional knowledge systems for ethnopharmacological research: theoretical and methodological contributions. *Journal of Ethnobiology and Ethnomedicine*. 2010; 6: 32
16. Popp J. How indigenous knowledge advances modern science and technology. *The Conversation*. 2018. Accessed 18 April 2020. Available <https://theconversation.com/how-indigenous-knowledge-advances-modern-science-and-technology-89351>.
17. Obembe OA. Stomata complex in some shrubs and trees. *Global Journal of Biology, Agriculture and Health Science*. 2015; 4(2):164-172

18. Metcalfe CR, Chalk L. Anatomy of the Dicotyledons. Volume 1. Oxford University Press, London. 1950
19. Pant DD, Verma BK. Taxonomy of the genus Ephedra: Significance of stem and leaf epidermis and cuticle. Botanical Journal of the Linnean Society. 1974; 69: 287-308.
20. Wilkinson HP. The plant surface (mainly leaf). In Metcalfe CR, Chalk L. (Eds.). *Anatomy of the Dicotyledons* (2nd ed.). Oxford, England: Oxford Press. 1979.
21. Adegbite AE. Leaf epidermal studies in three Nigerian species of *Aspilia* (Heliantheae-Asteraceae) and two hybrids. Nigerian Journal of Botany. 1995; 8: 25-33.
22. Pandey BP. *Plant Anatomy: Including Embryology and Morphogenesis of Angiosperms* (6th ed.). New Delhi: S. Chand. 2004
23. Nurul-Aini CAC, Noraini T, Latiff A, Amirul-Aiman AJ, Ruzi AR, Idris S. Taxonomic Significance of Leaf Micromorphology in Some Selected Taxa of Acanthaceae (Peninsular Malaysia). AIP Conference Proceedings. 2014; 727-733
24. Chen-Yi L Dun-Yan T. The taxonomic significance of leaf epidermal micromorphological characters in distinguishing 43 **species** of *Allium* l. (Amaryllidaceae) from Central Asia. Pakistan Journal of Botany. 2015; 47(5): 1979-1988.
25. Wiegleb G. The importance of stem anatomical characters for systematics of genus *Potamogeton* L. Flora. 1990; 184: 197-208.
26. Metcalfe CR. On the taxonomic value of the anatomical structure of the vegetative organs of the dicotyledons: An introduction with special reference to the anatomy of the leaf and stem. Proceedings of the Linnean Society of London. 1942; 155(3): 211-223.
27. Mbagwu FN, Inyama CN. *Stem anatomy of three Chrysophyllum Species (Sapotaceae) and their relevance for taxonomy (Systematic Botany)*. International Conference on Plant Physiology and Pathology. June 09-10, Dallas, USA. 2016.
28. Aziagba BO, Okeke CU. Taxonomic significance of stem and petiole anatomy of three varieties of *Vigna unguiculata*. American Journal of Life Sciences Research. 2017; 5(1): 1-5.
29. Goswami S. *Anatomy in relation to plant Taxonomy*. 2018. Available: <https://www.biologydiscussion.com/plant-taxonomy/30463>.
30. Cavente AM, Andreato RHP, Vieira RC. Stem anatomy of *Rhipsalis* (Cactaceae) and its relevance for taxonomy. Plant Systematics and Evolution. 2008; 276 (1): 1-7.
31. Kaplan Z, Symoens J. Taxonomy, distribution and nomenclature of three confused broad-leaved *Potamogeton* species in Africa and surrounding islands. Botanical Journal of the Linnean Society. 2005; 148(3): 329-357.