

Original Research Article

Does Oral Administration of Fresh and Dry Ginger (*ZingiberOfficinale*) Juice have effect on The Lipid Profile of Albino Rats.

ABSTRACT

Aim: This study compares the effect of fresh and dry ginger juice on the lipid profile of albino rats.

Methodology: Fifteen(15) albino rats with average weight of 150g divided into three groups of five (5) rats each were used for the study.5ml of the ginger juice was administered orally for 21 days using the gavage oral tube to the experimental animals. Blood samples were collected via cardiac puncture after a chloroform anesthesia was given to the animals to a heparinized sample bottle. Plasma samples were analyzed for total cholesterol, triglyceride, high density lipoprotein using the colorimeter while and low-density lipoprotein level was calculated.

Results: There was no significant difference in cholesterol, triglyceride, high density lipoprotein and low-density lipoprotein levels ($P>0.05$) when the group 1(control group) was compared with group 2 (dry ginger group) and group 3 (fresh ginger group).

Conclusion: Ginger juice in different preparation state does not alter the lipid profiling in the experimental animals over a period of 21 days.

Key words: *Dry ginger, Fresh ginger, Ginger juice, Lipid profile, Albino rats.*

1.INTRODUCTION

Hyperlipidemias reveals the onset of abnormalities of lipid metabolism and it is secondary to the manifestation and progression of arteriosclerotic state in humans [1, 2].One of the most important risk factors associated with coronary artery disease (CAD) is dyslipidemia. Low HDL-cholesterol (HDL-C), high triglycerides (TG) and high LDL-cholesterol (LDL-C) levels have been associated with increased incidence of CAD [3]. In addition to diet, herbal medicine has been explored as a pharmacologic modality in preventing alterations in lipid metabolism [4,5].

Ginger (*Zingiberofficinale*) a well-known spice plant with a pungent appetizing taste has been in use in traditional oriental medicine [6] to treat a number of medical conditions including headache, cold, arthritis etc.[7]. It is included in the family of Zingiberaceae .This family covers up to 24 genus and around 300 species with the genus Zingiber having about 20 species. It possesses a perennial tuberous or rhizomatous root with an upright annual stalk (pseudo stem about 60-90cm tall) covered with flat sheaths that may be taken off as stalk with about 8-12 distiches dark green leaves present on the stem [8, 9].Ginger (*Zingiberofficinale*) has been used in folk medicine for a long time. Its extract and juice have been shown to exhibit a variety of biological activities.

Ginger is found almost in every home and plays a major role as a spice for food as well as a herbal and dietary supplement with lesser toxic effects but of more therapeutic effect. As a means to achieving certain health goals, individuals resort to taking ginger due to its numerous health benefits including its role in lipid metabolism and as a hypolipidemic agent [10, 11]. Many varieties of ginger are found such as processed, coated or unscrapped, unbleached (natural) and bleached ginger. Aqueous and organic extracts of ginger have illustrated a high number of active components including gingerols, shogaols, alkaloids, tannin, saponin, curcumin phenolic and ketone derivatives [12,13,14,15].

Comment [a1]: In Asian subcontinent

Comment [a2]: Reference should be added .

Comment [a3]: Like candies ,flavored drinks etc

Comment [a4]: Check it one more time , curcumin is found in Haldi (turmeric).

High plasma levels of lipoproteins are strongly associated with coronary artery disease and an independent risk factor for atherosclerosis. Hence investigation of the effect of either fresh or dried ginger juice on lipid profile of albino rats in this study.

Comment [a5]: Reference should be added .such as Prevalence and determinants of dyslipidemia: data from a Saudi University Clinic YT Al-Hassan, EL Fabella, E Estrella, M Aatif The Open Public Health Journal 11 (1)

2.MATERIALS AND METHODS

2.1. Experimental animals- Fifteen (15) albino rats weighing averagely 150g were used for this study. They were purchased from the Animal house of the Department of Human Physiology, University of Port Harcourt. . They were housed in wired cages, acclimatized for two weeks with temperature controlled at 25°C before commencement of treatments They were fed with rat feed (finisher) commercially prepared from the Top feed Company, Eastern Premier Feed Mill Ltd, Aba, Abia State, Nigeria and had access to water (ad libitum) throughout the period. The conditions of the animals were in conformity with standards as outlined by the National Academy of Science [16,17].

Comment [a6]: Comparative

Comment [a7]: It can be , So aim of this study is to find comparative effect of Fresh and Dry juice of ZingiberOfficinaleon The Lipid Profile of Albino Rats.

2.2. Ginger juice- The ginger was purchase from the mile 3 market in Port Harcourt. 600g of fresh rhizome of ginger after cleaning and scrapping the superficial skin was cut into smaller pieces. Using a grinder, the pieces were made into a paste like form which was taken unto a white clean sieve cloth and the liquid squeezed out from it into a clean airtight sterile container yielding about 300mls.The stocked juice as obtained was refrigerated in the refrigerator at 4°C and used for the experiment.

Another set of fresh ginger rhizome were also dried using a hot air oven at a temperature of 140°C for 3 days, the dried ginger was weighed to a 600g weight, then ground to powder using a grinder. This was dissolved in 300mls of distilled water, sieved out from a white clean cloth into a clean container and stored in the refrigerator for administration each week as the experiment lasted.

Comment [a8]: At this temperature might be some active ingredients will lose their activity .

Comment [a9]: 300 ml.

2.3. Experimental Design The 15 albino rats were divided into 3 groups namely Group 1 with 5 rats serving as Control fed only with food and water. Group 2 (Dry Ginger Juice) with rats that were fed with food and water and administered orally 5mls of dry ginger juice while the last is group 3 (fresh ginger juice). This group was fed with food and water then administered 5mls of fresh ginger juice during the experiment which lasted for 21 days.The administration of the juice was done once per day.

2.4. Collection of blood samples- After the treatment period of 21days, the rats were left 24hours in a fasting state, then on the 22nd day, they were anaesthetized using chloroform, sacrificed and blood collected via cardiac puncture into lithium heparinized sample bottles. This was spun using a centrifuge after which the plasma was separated and used for the estimation of total cholesterol, triglycerides, high density lipoprotein using colorimetric methods with the

Randox kit UK. The Low-density lipoprotein concentration was calculated using the formula as described by Fried Wald formula [18].

$$\text{LDL-C} = (\text{TC}) - (\text{HDL-C}) - (\text{TGs}/5)$$

2.5. Statistical Analysis- Results were presented as mean \pm standard deviation while the data obtained were analyzed statistically using Graph pad prism version 8.2.0. Values were considered significant at 95% confidence interval.

3. RESULTS: The effect of fresh and dried ginger juice on plasma total cholesterol (TC), triglyceride (TG). High density lipoprotein (HDL) and low-density lipoprotein (LDL)) levels of albino rats were compared and reflected in the Tables 1-3.

Table 1: Mean \pm SD of Lipid Profile of Control (Group 1) and Dried Ginger Juice (Group 2)

Groups/Parameters	TC mmol/L	TG (mmol/L)	HDL (mmol/L)	LDL (mmol/L)
Group 1 (Control)	2.70 \pm 0.74	0.94 \pm 0.25	0.66 \pm 0.21	1.61 \pm 0.63
Group 2 Dried Ginger juice)	2.68 \pm 1.59	0.92 \pm 0.90	0.64 \pm 0.98	1.62 \pm 0.39
T Valve	0.1377	2.154	0.6730	0.3372
Pvalue	0.893	0.052	0.514	0.745
Remark	NS	NS	NS	NS

Key: NS-Non significant

Table 2: Mean \pm SD of Lipid Profile of Control (Group 1) and Fresh Ginger Juice (Group 3)

Groups/Parameters	TC mmol/L	TG (mmol/L)	HDL (mmol/L)	LDL (mmol/L)
Group 1 (Control)	2.73 \pm 0.74	0.94 \pm 0.25	0.66 \pm 0.21	1.61 \pm 0.63
Group 3 Fresh Ginger juice)	2.69 \pm 1.20	0.93 \pm 0.36	0.67 \pm 0.30	1.60 \pm 0.46
T valve	0.4416	0.0849	0.2414	0.8470
Pvalue	0.667	0.934	0.813	0.102
Remark	NS	NS	NS	NS

Key: NS-Non significant

Table 3: Mean \pm SD of Lipid Profile parameters in Albino rats administered fresh and Dried ginger juice.

Groups/Parameters	TC mmol/L	TG (mmol/L)	HDL (mmol/L)	LDL (mmol/L)
Group 2 (Dried Ginger)	2.68 \pm 1.59	0.92 \pm 0.90	0.64 \pm 0.98	1.62 \pm 0.39
Group 3(Fresh Ginger)	2.69 \pm 1.20	0.93 \pm 0.36	0.67 \pm 0.30	1.60 \pm 0.46
Tvalue	0.1595	1.879	0.1589	3.297
Pvalue	0.854	0.185	0.854	0.072
Remark	NS	NS	NS	NS

Key: NS-Non significant

4.DISCUSSION: The non-significant levels seen when the control was compared with the fresh ginger treated groups is in variance with study carried out by Prasad et al., [19] who reported a

Comment [a10]: Data of all three tables can also be represented via a bar graph.

Comment [a11]: How it can be 2.73 now while earlier it was 2.70 in table -1.

decrease in cholesterol level and increase in HDL level which could be adduced to modulatory influence on cholesterol metabolism and turnover. This observation is also in contrast with the work done by [20] who administered ginger orally to the rats resulting in a reduction in cholesterol and a lesser effect on LDL level though the nature of the ginger preparations are not the same.

The nature of preparing the ginger might have an effect on the results observed in this study as compared to the study of Li et al.[21] where activities of ginger prepared in different modes were compared and significant differences were observed, However, in this study the effect from the dried ginger juice and the fresh ginger juice on the lipid profile of the rats were not significantly different. The variation noted cannot be easily **accounted for**. The differences in preparation of ginger may be contributory factors for this variation. Also, Cheng et al.[22] in their study showed that steamed ginger had the potentials to change chemical profile of ginger thereby enhancing some of its beneficial activities however this kind of observation was not seen in this study considering dried and fresh ginger prepared juice. This study is also in contrast with the **study of** [23] where animals fed with ginger had a significantly increased HDL and reduced cholesterol which might be adduced to the activity of hepatic cholesterol-7- hydroxylase, a rate limiting enzyme in bile acids biosynthesis which stimulates cholesterol conversion to bile acids thereby removing cholesterol from the body [24]. However in a meta analysis carried out by Pourmasoumi et al.[25], ginger was considered a lipid lowering agent as reflected in the reduced levels of Triglycerides (TG) and low density lipoproteins (LDL) with no significant effect on Cholesterol and high density lipoprotein (HDL) in the plasma. A study by Murad et al.[26] revealed that powdered ginger administration was able to significantly reduced LDL and Cholesterol levels though this was carried out in humans. In the study of Fakhri et al [27], ginger supplementation was able to reduce cholesterol, triglycerides and LDL significantly but no significant effect on HDL which is also in contrast with our present observation in this study. Reviews of the more recent trials, suggest that ginger shows considerable hypolipidemic effect in vitro and in some animal studies [28]. Hence it will be worth studying and comparing the effects and potency of the different forms of ginger preparation; fresh, dry, steamed, powdered, juice or extract considering their doses and mode of administration.

5.CONCLUSION. Ginger juice in their different prepared forms had no significant effects on the lipid profile of the **albino rats**.

REFERENCE

1. Vodnala D, Rubenfire M, Brook RD. Secondary causes of dyslipidemia. American Journal of Cardiology. 2012;110(6):823-825.
2. Veerkemp MJ, De Graaf J, Den Heijer M, Blom HJ, Stalenhoef AF. Plasma homocysteine in subjects with familial combined hyperlipidemia. Artherosclerosis. 2003; 166:111-117
3. Weissglas-Volkov D, Pajukanta P. Genetic causes of high and low serum HDL cholesterol. The Journal of Lipid Research, 2010; 51: 2032-2057.

Comment [a12]: Conclusion should be elaborated to some more extent as, what can be the reasons here not finding any significant differences. While in discussion part there were so many studies quoted, which are contradictory to the finding of this manuscript. Thanks

4. Maqbool M, Dar MA, Gani I, Mir SA, Khan M. Herbal Medicines as an Alternative Source of Therapy: A Review. *World Journal of Pharmacy and Pharmaceutical Sciences*. 2012; 8(2): 374-380.
5. Bhandari U, Sharma JN, Zafar R. The protective action of ethanolic ginger (*ZingiberOfficinale*) extract in cholesterol fed rabbits. *Journal of Ethnopharmacology*. 1998; 61:167-171.
6. Ghayur MN, Gilani AH. Ginger lowers blood pressure through blockage of voltage-dependent calcium channels. *Journal of Cardiovascular Pharmacology*. 2005; 45: 74-80.
7. Ebrahimzadeh AX, Malex MA, Jaxadixala Z, Zununi VS, Ostradrahimi A. A systemic review of the anti-obesity and weight lowering effect of ginger and its mechanism of actions. *Phytotherapy Research*. 2018; 32: 577-585.
8. Moghaddasi MS, KashaniHH. Ginger (*Zingiberofficinale*): A review. *Journal of Medicinal Plants Research*. 2012; 6(26): 4255-4258,
9. Grzanna R, Lindmark L, Frondoza CG. Ginger an herbal medicinal product with broad anti-inflammatoryactions. *Journal of Medicine and Food*. 2005; 8:125-132.
10. Leech J. 11 proven Health Benefits of Ginger, *Nutrition*. 2021;1: 1-34.
11. Pillai KK, Bhandari U, Kanojia R. Effect of ethanolic extract of *ZingiberOfficinale* on dyslipidaemia in rats, *Journal of Ethnopharmacology*. 2005; 97: 227-230.
12. Qian-Qian M, Xiao-Yu Xu, Sui-Yu C, Ren-You G, Harold Corke TB, Hua-Bin Li. Bioactive compounds and Bioactivities of ginger (*Zingiberofficinale* Roscoe. *Foods*,2019; 8(6):185-187.
13. Wang J, Ke W, Bao R, Hu X, Chen F. Beneficial effect of ginger *Zingiberofficinale* Roscoe on obesity and metabolic syndrome, A review. *Annals of Academic Science*. 2017; 1398: 83-98.
14. Rahmani AH, Aly SM. Active ingredients of ginger as potential candidates in the prevention and treatment of diseases via modulation of biological activities. *International Journal of Physiology, Pathophysiology & Pharmacology*. 2014; 6(2): 125–136.
15. Shrin Adel PR & Prakash J (2010). Chemical composition and qanti oxidant properties of ginger root. *Journal of Medicinal plants and Reserves*,4(24),2674-2679.
16. ILAR. Guideline for the care and use of laboratory animal.8th edition. National academic press. Washington D.C, 2011.
17. OLAW. Institution Animal care and use committee Guidebook.2nd edition NIH publication; Bethesda. 2002.

18. Eblen-ZajjurA Eblen-ZajjurM. Estimation of low density lipoprotein cholesterol concentration: regression analysis versus Friedwald's formula. *Revista Medica de Chile*. 2001; 129(11):1263-1270.
19. Prasad SS Sushil, K Vaspeyee SK Bhavsar VH. To establish the effect of ginger juice on important parameters of Lipid profile. *International Journal of Pharmacology Sciences and Research*. 2012; 3(4): 352-357.
20. Thomson M, Al-Qattan KK, Al-Sawan SM, Alnaqeeb MA, Khan I, Ali M. The use of ginger (*Zingiberofficinale* Rosc.) as a potential anti-inflammatory and antithrombotic agent. *Prostaglandins Leukotrienes and Essential Fatty Acids*. 2002; 67(6):475-478.
21. Li Y Hong Y Han Y Wang Y Xia L. Chemical characterization and anti-oxidant activities comparison in fresh, dried, stir-frying and carbonized ginger. *Journal of Chromatography*. 2016; 1011: 223-232.
22. Cheng XL Lin Q Peng YB Qil W Li P. Steamed ginger (*Zingiberofficinale*): changed chemical profile and increased anti-cancer potentials. *Food Chemistry*. 2011; 129:1785-1792.
23. Ahmed RS Sharma SB. Biochemical studies on combined effects of garlic (*Allium sativum* Linn) and ginger (*Zingiberofficinale* Rosc) in albino rats. *Indian Journal of Experimental Biology*. 1997; 35(8):841-843.
24. Gälman C Arvidsson I Angelin B Rudling M. Monitoring hepatic cholesterol 7-hydroxylase activity by assay of the stable bile acid intermediate 7-hydroxy-4-cholesten-3-one in peripheral blood. *Journal of Lipid Research*. 2003; 44:859-866.
25. Pourmasoumi M, Hadi A, Rafie N, Najafgholizadeh A, Mohammadi H, Rouhani MH. The effect of ginger supplementation on lipid profile: A systematic review and meta-analysis of clinical trials. *Phytomedicine*. 2018; 1(43): 28-36.
26. Murad S Niaz K Aslam H. Effects of Ginger on LDL-C, Total Cholesterol and Body Weight. *Clinical & Medical Biochemistry*. 2018; 4(2): 1-4.
27. Fakhri Z, Shab-Bidar S, Firoozi S, Djafarian K. The Effects of Ginger Supplementation on Lipid Profile: A Meta-Analysis of Randomized Clinical Trials. *Herbal Medicine Journal*. 2018; 3(3):120-131.
28. Nicoll R, Henein MY. Ginger (*Zingiberofficinale* Roscoe): A hot remedy for cardiovascular disease?, *International Journal of Cardiology*. 2009; 131(3):408-409,

UNDER PEER REVIEW

