

Original Research Article

Impact of Host age, Gestational week and Parity on Susceptibility to Urinary Tract Infections

ABSTRACT

Background: Urinary tract infections are inflammatory responses of the urothelium to non-flora microorganisms. **Aim:** This study aims at identifying the urinary tract pathogens in association with other risk factors in pregnant women residing in the rural community of Okada, Edo State. **Methods:** The study entails a cross sectional study involving pregnant women at various gestational age, a mid stream, non-repetitive urine samples were collected in sterile bottles and cultured on bacteriological media at 37°C overnight, isolates were identified using standard conventional procedures. **Results:** The organisms isolated include Staphylococcus aureus, Escherichia coli, Candida albicans with prevalence of 35.7%, 5.1% and 27.1% respectively, the age range 26-31 have a prevalence of 37.7% with a p-value of 0.006, the distribution of UTI in relation to other risk factors such as parity, haemoglobin genotype and level of education had p-value of 0.179, 0.067 and 0.0001 respectively. **Conclusion:** The relationship between UTI and parity is not statistically significant, but statistically significant with level of education

Comment [MOU1]: to

Comment [MOU2]: this target isn't reported in the title

Comment [MOU3]: where is age in the conclusion

Background to the Study

Urinary tract infections (UTIs) are inflammatory responses of the urothelium to bacterial invasion by non-flora microorganisms. It is one of the most common medical issues associated with pregnancy and may involve the reproductive organs if not well managed (Brook *et al.*, 2001). Despite major progress in the management of urinary tract infections in

pregnancy, they are still linked to negative feto-maternal outcomes (Ezugwuet *al.*, 2021). According to Ali and Abdallah, (2019), urinary tract infection describes microbial colonization or inflammation of the bladder (cystitis), urethra (urethritis), or renal pelvis and kidneys (pyelonephritis) with growth of up to 10^6 organisms per μl of urine in an asymptomatic presentation (Emilie *et al.*, 2011). The presence and growth of these microorganisms in the urinary tract can lead to these infections, irrespective of age and mostly in pregnancy in association with other risk factors can be life threatening (Tamalliet *al.*, 2013).

Culturing of urine specimens is the gold standard for diagnosing urinary pathogen, as well as allowing for quantitative bacteriuria quantification (Ezugwuet *al.*, 2021). According to Nwachukwu *et al* (2018), the common etiologic agents of UTI are *Escherichia coli*, *Klebsiella spp*, *Staphylococcus aureus*, *Pseudomonas spp*, and *Streptococcus*. These organisms can be found in the external genitalia, vagina, rectum, genital tract, and gastrointestinal tract and can infect cells in the urinary tract by attachment and eventual ascension, causing bacteremia, due to the renal cells being compromised. Advanced age, number of parturitions, number of coitus per week, recessive sickle cell anemia, and diabetes, previous history of UTI, immunodeficiency, and urinary tract abnormalities are conditions that can increase the risk of UTI in pregnant women (Giraldo *et al*, 2012).

Pregnancy changes including anatomical, physiological and hormonal increases susceptibility to the development of UTI. These changes can be due to heightened levels of progesterone and estrogen but also due to pressure created by the growing uterus (Taye *et al.*, 2018). About 70% of women during pregnancy develop glycosuria, which supports bacterial growth in the urine (Kant *et al.*, 2017). UTIs are may also be associated with premature delivery, low-birth-weight infants, cesarean delivery, morphological abnormalities, and infant mortality (Nabbugodiet *al.*, 2015). Early diagnosis, administering of adequate treatment using

Comment [MOU4]: What is the role of Ali and Abdallah in urinary tract infection describes microbial colonization

antibiotics has been shown to help reduce these risks by 80% (Nicolle, 2015). In this rural community there is paucity of UTI in relation to other associated risk factors such as gestational age, parity and haemoglobinopathy. This study aims at identifying the urinary tract pathogens in association with other risk factors in pregnant women residing in the rural community of Okada, Edo State, Nigeria.

Materials and methods

The study area

The study was carried out in Igbinedion University Teaching Hospital, Okada in Edo State using the facilities of the Medical Microbiology Laboratory.

Study design

This is a cross-sectional study involving pregnant women visiting the antenatal clinic and general outpatient clinic. The study comprised of 150 women who were pregnant at various gestational ages. A mid-stream non-repetitive urine was collected from each participant following administration of questionnaire after informed written consent. Urine samples were collected for urine microscopy, culture and sensitivity. Furthermore, those currently on antibiotics were excluded from the study.

Ethical Approval

This study was approved by the ethical committee of Igbinedion University Teaching Hospital, Okada Edo state. The purpose of this study was well explained to the pregnant women and strict adherence to confidentiality maintained.

Screening of urine bacteria:

Culture method described by Chesbrough, (2010) was used for the isolation of bacteria.

Comment [MOU5]: You must specify criteria of inclusion / asymptomatic, symptomatic, both?
Urine sample at least after 3 hours without urination

Briefly, mid-stream urine was cultured on Cysteine lactose electrolyte deficient agar (CLED) and Sabouraud dextrose agar. At 37°C, all cultures were incubated aerobically for 18-24hrs.

Identification of Isolates

All bacterial isolates were subjected to taxonomic analysis by the determination of their morphological, physiological and biochemical characteristics as adopted by (Fawole and Oso, 1988; Holt et al., 1994; Cowan and Steel, 2003; Cheesbrough, 2010).

RESULTS

150 urine samples were collected from pregnant women attending Igbinedion University Teaching Hospital (IUTH). The age ranges of the subjects were between 20-47 years as shown in Table 1. The age ranges 20-25, 26-31, 32-37 and 38-43 had 17.1%, 37.7%, 32% and 15.4% prevalence for urinary tract infections respectively and is statistically significant with P- value 0.006. Table 2 shows the distribution of urinary tract infections in pregnant women in relation to gestational age, the week, (1- 12), (13- 25) and (26-40) had 13.6%, 17.9% and 56.8% prevalence rate of infection and are statistically significant with p- value 0.0001. The distribution of UTI in relation to other factors such as marital status, level of education, past history of infection, sexual activity and haemoglobin genotype is shown in Table 3; the level of Education with respect to UTI is statistically significant with p-value of 0.0001. The distribution of UTI in pregnancy in relation to parity is shown in Table 4, parity of 0, 1, 2, 3, and > 4 had UTI prevalence of 31.3%, 21.1%, 15.4%, 22.2% and 43.8% respectively and is not statistically significant with respect to UTI.

Table 5 shows the microbial distribution; *Staphylococcus aureus*, *Candida albicans*, and *Escherichia coli* had occurrence rate of 35.7%, 27.1% and 5.1% respectively.

Table 1: Prevalence of urinary tract infections in pregnant women in relation to Age

| AGE GROUP | FREQUENCY | NUMBER POSITIVE | (%) | P-value |
|-----------|-----------|-----------------|------|---------|
| 20-25 | 41 | 7 | 17.1 | 0.006 |
| 26-31 | 69 | 26 | 37.7 | |
| 32-37 | 25 | 8 | 32 | |
| 38-43 | 13 | 2 | 15.4 | |
| 44-47 | 2 | 0 | 0 | |
| TOTAL | 150 | 43 | | |

Table 2: Distribution of urinary tract infections in pregnant women in relation to gestational period

| GESTATION (WEEK) | FREQUENCY | NUMBER POSITIVE | (%) | p- value |
|---------------------|-----------|-----------------|------|----------|
| 1-12 | 22 | 3 | 13.6 | < 0.0001 |
| 13-25 | 84 | 15 | 17.9 | |
| 26-40 | 44 | 25 | 56.8 | |
| TOTAL | 150 | 43 | 100 | |

Table 3: Distribution of urinary tract infections in relation to other factors

| FACTORS | | N | NOS POS | (%) | p value |
|----------------------|------------|-----|---------|------|---------|
| STATUS | MARRIED | 139 | 35 | 25.2 | |
| | UNMARRIED | 11 | 8 | 72.7 | |
| EDUCATION | PRIMARY | 9 | 8 | 88.9 | |
| | SECONDARY | 72 | 15 | 20.8 | |
| | TERTIARY | 69 | 20 | 28.9 | |
| PAST HISTORY OF UTI | PRESENT | 101 | 33 | 32.7 | |
| | ABSENT | 49 | 10 | 20.4 | |
| SEXUAL ACTIVITY | ACTIVE | 135 | 42 | 31.1 | |
| | NON-ACTIVE | 15 | 1 | 6.7 | |
| HAEMOGLOBIN GENOTYPE | AA | 114 | 38 | 33.3 | |
| | AS | 33 | 5 | 15.2 | |
| | AC | 3 | 0 | 0 | |

STATUS P = 0.126

EDUCATION = 0.0001

PAST HISTORY OF UTI P= 129

SEXUAL ACTIVITY P= 0.067

HB GENTOTYPE P= 0.683

Comment [MOU6]: All married or pregnant women are sexually active

Comment [MOU7]: To report on the Last column

TABLE 4: DISTRIBUTION OF UTI IN **PRENANT** WOMEN IN RELATION TO PARITY

Comment [MOU8]: pregnant

| PARITY | NOS EXAMINED | NOS WITH UTI | (%) | P-value |
|--------------|--------------|--------------|------|---------|
| 0 | 32 | 10 | 31.3 | 0.179 |
| 1 | 57 | 12 | 21.1 | |
| 2 | 39 | 6 | 15.4 | |
| 3 | 36 | 8 | 22.2 | |
| >4 | 16 | 7 | 43.8 | |
| TOTAL | 150 | 43 | | |

UNDER PEER REVIEW

TABLE 5: DISTRIBUTION OF ISOLATES FROM PREGNANT WOMEN

| ISOLATES | NUMBER ISOLATED | (%) |
|----------------------|------------------------|------------|
| Bacillus specie | 11 | 8.5 |
| S. epidermidis | 11 | 8.5 |
| Escherichia coli | 7 | 5.4 |
| Enterobacter specie | 7 | 5.4 |
| S. aureus | 46 | 35.7 |
| Lactobacillus specie | 3 | 2.3 |
| Candida albicans | 35 | 27.1 |
| Klebsiella aerogenes | 9 | 6.9 |
| TOTAL | 129 | |

Comment [MOU9]: report the total percentage on this line (100%)

UNDER PEER REVIEW

DISCUSSION

Infection of the urinary tract (UTI) represents the most common medical complication of pregnancy especially in asymptomatic bacteriuria. Pregnant women are at greater risk of UTIs, particularly because of the physiologic, hormonal and anatomic changes that occur in normal pregnancy.

Comment [MOU10]: bacteriuria

In this research, about 150 urine samples of pregnant women were collected of which the overall prevalence of UTI among pregnant women in this study was 43(28.7%). This is less than the prevalence reported in Saudi Arabia 53.5% (El-Kashif *et al.*, 2019), but within the range obtained in Ambo Central Ethiopia 18.7%, Ismailia, Egypt 29%, Benin city, Nigeria 21% and in Nepal 37.8% respectively (Thakur *et al.*, 2020). These variations in prevalence might be as a result of differences in social habits, religious affiliation, and personal hygiene of participants. However, spirituality has a way of making people play down on personal and environmental hygiene.

The age range in this research with high prevalence was between 26-31 years with 26(37.7%) which agrees with a study carried out in Onitsha, Nigeria (Nwachukwu *et al.*, 2018). This may be as a result of sexual activity which increases the risk of UTI and the women of such age group are mostly sexually active.

Comment [MOU11]: perhaps high rate of sexual activity at this age

With respect to parity; Parity greater than 4 has the highest prevalence with 4(43.8%). however, in a study conducted by in Gorgan, the highest rate of infections was after the third pregnancies (4.73%) (Mobbasher *et al.*, 2001). The results from the studies of Mobbasher *et al.* were consistent with the results from the study in America (Gibbs *et al.*, 1994).

Comment [MOU12]: reformulate

The gestational period of the participants showed that those in their 2nd trimester have the highest prevalence with 25(56.8%). This is consistent with the result reported by Okonko, 2009. This difference may be as a result of either change in urinary stasis and vesicoureteral

reflux or decrease in urinary progesterones and oestrogens in the various trimester of pregnancy.

Based on level of education, the highest prevalence was among those who attained primary level of education with 8(88.9%), while those with tertiary education had 28.9%, difference could be attributed to awareness and hygiene. Also based on marital status of participants, the unmarried status had a distributions of 8(72.7%), while the married status had 35 (25.2%) with UTI. This could be due to the level of socioeconomic tendencies of the participants.

Among isolated pathogens, *Staphylococcus aureus* was the most predominant bacteria with 46(35.7%). This is in contrast to other results gotten in other studies conducted in Babol city, *E. coli* was indicated to be the cause of 83% of UTIs in pregnant women (Yaghoobiet *al.*, 2006) and Moreover, in studies conducted by Masinde in Tanzania, Al- Haddad in Yemen, Hamdan in Sudan and Totsika in Australlia (Totsikaet *al.*, 2012), *E. coli* was the main cause of UTIs among women of those areas and the prevalence in those regions were 47.2%, 41.5%, 42.4%, respectively.

5.1 Conclusion

This study concludes that relationship between UTI and parity is not statistically significant. However, education status and gestational growth are statistically significant in relation to UTI among pregnant women, regardless of other associated risk factors such as haemoglobin genotype, past history of UTI and marital status.

References

- Ali, M., & Abdallah, M. S. (2019). Prevalence of Urinary Tract Infection among Pregnant Women in Kano, Northern Nigeria. *Archives of Reproductive Medicine and Sexual Health*, 2(1), 23-29.

- Cheesbrough, M. (2010). District Laboratory practice in tropical countries (Part II). Cambridge University; pp.19-11.
- Cowan , S.T.and Steel, K. J. (2003). Cowan and Steels Manual for the Identification of Medical Bacteria. (5thEdn). Cambridge University Press,Cambridge, London, p.1-210.
- El-Kashif, M. M. L. (2019). Urinary tract infection among pregnant women and its associated risk factors: a cross-sectional study. *Biomedical Pharmacology Journal*,12(4), 2003–2010.
- Ezugwu,I. E., Afunwa, R. E., Onyia, F. A., Chukwunwejim,, C. R., & Eze, A. E. (2021). Prevalence of Urinary Tract Infections and Associated Risk Factors among Pregnant Women in Enugu Metropolis, *Nigerian Journal of Biosciences and Medicines*,9(10)
- Fawole, M.O. and Oso, B.A. (1988). Laboratory manual for microbiology. Spectrum Book Ltd,Ibadan.5th edition. **1**:22-45.
- Gibbs, R. S., McGregor, J. A., Mead, P. B., Eschenbach, D. A., Hager, W. D., & Sweet, R. L. (1994). A survey of practices in infectious diseases by obstetrician-gynecologists. *Obstetrics and Gynecology*,83(4):631–636.
- Giraldó, P. C., Araújo, E. D., Junior, J. E., Amaral, G. D., Passos, R. L., & Gonçalves, A. K. (2012). The Prevalence of Urogenital Infections in Pregnant Women Experiencing Preterm and Full-Term Labor. *Infectious Diseases Obstetrics Gynecology*,1–4. doi: 10.1155/2012/878241
- Holt, J.G., Krieg, N.R., Sneath, P.H.A., Stanley, J.T. and Williams, S.T. (1994). Bergey's Manual of Determinative Bacteriology, 3rd edition. *Williams and Wilkins, Baltimore*.**9**:787.
- Mobbasheri, E. T., Ghaemi, E., Moujloo, M., & Vakili, M. A. (2001). Prevalence of bacteriuria during pregnancy in Gorgan, Iran. *Gorgan Medical Journal*,**9**:42–47.

- Nabbugodi, W. F., Gichuhi, J. W., & Mugo, N. W. (2015). Prevalence of Urinary Tract Infection, Microbial Etiology, and Antibiotic Sensitivity Pattern among Antenatal Women Presenting with Lower Abdominal Pains at Kenyatta National Hospital, Nairobi, Kenya. *The Open Access Journal of Science and Technology*, 3, 1-6.
- Nwachukwu, E., Onyebuchi, O., & Michael, O. (2018). Prevalence of urinary tract infections in pregnant women in Onitsha, Nigeria. *Journal of Bacteriology Mycology*, 6(5), 284-285. DOI: 10.15406/jbmoa.2018.06.00219
- Okonko, I. O., Ijandipe, L. A., Ilusanya, O. A., Donbraye-Emmanuel, O. B., Ejembi, J., Udeze, A. O., Egun, O. C., Fowotade, A., & Nkang, A. O. (2009). Incidence of Urinary Tract Infection (UTI) among Pregnant Women in Ibadan, South-Western Nigeria. *African Journal of Biotechnology*, 8(23), 6649-6657.
- Tamalli, M., Sangar, B., & Alghazal, A. M. (2013). Urinary tract infection in pregnancy at Al-khoms, Libya. *International Journal of Medicine and Medical Sciences*, 3(5), 455-459.
- Taye, S., Getachew, M., Desalegn, Z., Biratu, A., & Mubashir, K. (2018). Bacterial profile, antibiotic susceptibility pattern and associated factors among pregnant women with urinary tract infection in Goba and Sinana Woredas, bale zone, Southeast Ethiopia. *BMC Research Notes*, 11(1), 799.
- Thakur, S. & Nagpal, K. L. (2020). Urinary tract infection in pregnant women at Kathmandu, Nepal. *Journal of Medical Care Research Review*, 3(9), 454-458.
- Totsika, M., Moriel, D. G., Idris, A., Rogers, B. A., Wurpel, D. J., & Phan, M. D. (2012). Uropathogenic Escherichia coli mediated urinary tract infection. *Current Drug Targets*, 13(11):1386-1399.
- Yaghobi, R. H., Roshan, AZ. PZ., Hajiahmadi, M., & Nasirim-Amiri, F. (2006). The role of hygienic behavior in incidence of urinary tract infection among pregnant women referred to health center, Babol, 2003- 2005]. *Journal of Babol University Medical Science*, 8(2):56-62.