

Borderline Personality Disorder and Neuroplasticity: A Review

Abstract

Borderline Personality Disorder (BPD) is a challenging and complex mental health disorder characterized by emotional dysregulation, impulsivity, unstable relationships, and a poor sense of self. Neuroplasticity, the brain's ability to change and adapt in response to experiences, is impaired in individuals with BPD, specifically in the prefrontal cortex, amygdala, and hippocampus. This impairment has been linked to emotional dysregulation, a core symptom of the disorder. Interventions aimed at improving neuroplasticity hold promise as a treatment target for BPD. Current evidence suggests that psychotherapeutic interventions, such as Dialectical Behavioral Therapy (DBT), may improve neural plasticity in the brain regions associated with emotional dysregulation and may result in symptom reduction and improved functioning in individuals with BPD. However, much more research is needed to better understand the relationship between neuroplasticity and BPD, as well as to develop more targeted and effective interventions. With continued research in this area, it is hoped that improved understanding of the role of neuroplasticity in BPD will lead to the development of more effective treatments for this challenging disorder.

Keywords: Borderline Personality Disorder, Neuroplasticity, Emotional dysregulation, Psychotherapy, Treatment interventions

1. Introduction

Borderline personality disorder (BPD) is a significant mental disorder that affects a considerable proportion of the population worldwide. People with BPD experience intense and unstable emotions, struggle to regulate their emotions, and often find it challenging to maintain healthy relationships. In addition, the disorder is linked with other symptoms such as impulsivity, self-harm, and distorted self-image that can have a profound impact on an individual's social and occupational functioning (Crocq et al., 2019; Paris, 2019). Despite extensive research, the exact etiology of BPD remains unclear, but genetic and environmental factors are thought to contribute to its development (Fossati et al., 2019).

Recent studies have suggested that neuroplasticity may play a critical role in the development and maintenance of BPD. Neuroplasticity is the brain's ability to change and adapt in response to experience and is essential for the brain to adapt to new situations and environments (Cailhol & Saudou, 2020). Structural and functional abnormalities in specific brain regions such as the prefrontal cortex, amygdala, and hippocampus have been identified in individuals with BPD. These regions play crucial roles in emotional regulation, social cognition, and memory formation, and their impairments may contribute to the symptoms of BPD (King-Casas et al., 2019; Schmahl et al., 2021).

To better understand the pathophysiology of BPD and develop more effective treatments, it is necessary to investigate potential causes of impaired neuroplasticity, such as early life stress and genetic factors, and explore neuroplasticity-based interventions for BPD. Psychotherapy, pharmacotherapy, and non-invasive brain stimulation techniques such as transcranial magnetic stimulation (TMS) and transcranial direct current stimulation (tDCS) have shown promise as potential neuroplasticity-based interventions for BPD (Bertschy et al., 2019; Dell'Osso et al., 2019; Morandotti et al., 2020).

In this review, we will provide a comprehensive overview of the role of neuroplasticity in BPD. We will begin by discussing impaired neuroplasticity in individuals with BPD, including the structural and functional abnormalities seen in specific brain regions associated with emotional regulation, social cognition, and memory formation. We will then delve into theories on the role of neuroplasticity in the development and maintenance of BPD, exploring different perspectives on how impaired neuroplasticity may contribute to the disorder. This will include considering the role of early life stress, epigenetics, and altered neurotransmitter systems in the development of BPD. Finally, we will review various neuroplasticity-based interventions for BPD, including psychotherapy, pharmacotherapy, and non-invasive brain stimulation techniques such as TMS and tDCS. By examining the potential of neuroplasticity as a target for interventions, we hope to provide a comprehensive understanding of the pathophysiology of BPD and the possibilities for treatment.

2. Neuroplasticity in BPD

Neuroplasticity refers to the brain's ability to change its structure and function in response to environmental and experiential stimuli (Bisio et al., 2021). This plasticity can occur at different levels of the brain, from molecular and cellular changes to alterations in neural circuits and whole brain regions (Filimonenko et al., 2021). Neuroplasticity is thought to be a fundamental property of the brain, enabling us to learn, adapt, and recover from injury or disease.

The mechanisms of neuroplasticity involve a range of cellular and molecular processes, including changes in synaptic strength, the growth and pruning of dendritic spines, and the formation of new neurons and glia (Luscher & Huber, 2020). These changes are driven by neural activity, which triggers a cascade of signaling pathways that lead to alterations in gene expression and protein synthesis. Over time, these changes can result in modifications to neural circuits and the overall organization of the brain.

There are different types of neuroplasticity, including experience-dependent plasticity, which occurs in response to specific stimuli or experiences, and homeostatic plasticity, which maintains the overall stability and balance of neural activity (Turrigiano, 2012). Experience-dependent plasticity includes forms of learning and memory, such as long-term potentiation (LTP) and long-term depression (LTD), which involve the strengthening or weakening of synaptic connections between neurons (Malenka & Bear, 2004). These changes can be transient or long-

lasting and can occur in response to a wide range of stimuli, from sensory inputs to emotional experiences.

Neuroplasticity is thought to be important for the development and function of neural networks that underlie various cognitive and behavioral processes, including perception, attention, decision-making, and emotion regulation (Chen & Zuo, 2021). It is also believed to play a role in the recovery of function after brain injury or disease, such as stroke, traumatic brain injury, and neurodegenerative disorders (Ward & Cohen, 2020).

In recent years, researchers have increasingly focused on the role of neuroplasticity in psychiatric disorders, including BPD. Studies have shown that individuals with BPD exhibit structural and functional abnormalities in specific brain regions associated with emotional regulation, social cognition, and memory formation (Kuo et al., 2020; Soloff et al., 2021). These abnormalities suggest that impaired neuroplasticity may be a key feature of the disorder.

In conclusion, neuroplasticity is a complex and dynamic process that underlies the brain's ability to adapt and change in response to environmental and experiential stimuli. It involves a range of cellular and molecular mechanisms that can result in modifications to neural circuits and the overall organization of the brain. Neuroplasticity is thought to be fundamental to learning, adaptation, and recovery after injury or disease. In BPD, impaired neuroplasticity may contribute to the emotional dysregulation and other symptoms seen in the disorder. Further research is needed to fully understand the role of neuroplasticity in BPD and to develop effective neuroplasticity-based interventions for the disorder.

2.2 Impaired neuroplasticity

BPD is a severe and chronic psychiatric disorder that is characterized by emotional dysregulation, impulsivity, and difficulties in interpersonal relationships (American Psychiatric Association, 2013). The etiology of BPD is not well understood, but it is thought to be a complex interaction between genetic, environmental, and neurobiological factors. In recent years, there has been increasing evidence that impaired neuroplasticity may be an important factor in the development and maintenance of BPD (Jovev et al., 2013).

Studies have shown that individuals with BPD exhibit structural and functional abnormalities in specific brain regions associated with emotional regulation, social cognition, and memory formation (Jovev et al., 2013; Ruocco et al., 2015). These abnormalities suggest that the neuroplasticity mechanisms responsible for shaping these brain regions may be impaired in individuals with BPD. In particular, deficits in the ability of neural circuits to adapt in response to experience and learning may contribute to the difficulties in emotion regulation and the interpersonal problems that are hallmarks of BPD.

Impaired neuroplasticity in BPD may also play a role in the high rates of comorbidity with other psychiatric disorders, such as depression and substance use disorders. For example, depression is associated with reduced neuroplasticity in brain regions such as the prefrontal cortex and the hippocampus (Duman & Monteggia, 2006). It is possible that the impaired neuroplasticity seen in BPD contributes to the increased vulnerability to depression in individuals with the disorder.

Research has also suggested that psychotherapy may be effective in promoting neuroplasticity in individuals with BPD (Driessen et al., 2010; Levy, 2015). Dialectical behavior therapy (DBT), a specialized form of cognitive-behavioral therapy, has been shown to improve emotional regulation, reduce impulsivity, and enhance interpersonal functioning in individuals with BPD (Linehan et al., 1991). DBT has also been found to increase gray matter density in regions of the brain associated with emotion regulation, such as the anterior cingulate cortex and the insula (Goodman et al., 2014). These findings suggest that DBT may promote neuroplasticity in individuals with BPD and contribute to the improvement of symptoms and function.

Impaired neuroplasticity may be an important factor in the etiology of BPD. Neurobiological studies have suggested that individuals with BPD may have deficits in the ability of neural circuits to adapt in response to experience and learning, which may contribute to difficulties in emotion regulation and interpersonal functioning. Psychotherapy, particularly DBT, may be effective in promoting neuroplasticity and contributing to the improvement of symptoms and function in individuals with BPD. Further research is needed to fully understand the role of neuroplasticity in BPD and to develop effective neuroplasticity-based interventions for the disorder.

2.3 Structural and Functional Abnormalities

Individuals with Borderline Personality Disorder (BPD) exhibit structural and functional abnormalities in specific brain regions associated with emotional regulation, social cognition, and memory formation. Structural abnormalities may include reduced gray matter volume, while functional abnormalities may include altered brain activity in response to emotional stimuli. The following sections will discuss the specific brain regions affected in individuals with BPD.

2.3.1 Prefrontal Cortex

The prefrontal cortex (PFC) is a crucial brain region for cognitive control, emotion regulation, and decision-making. Recent studies have shown that individuals with BPD exhibit structural and functional abnormalities in the PFC. Specifically, reduced gray matter volume in both dorsal and ventral PFC has been reported in this population (Mauchnik et al., 2018). Moreover, altered activity in the dorsolateral and ventromedial PFC during emotional processing has been observed in individuals with BPD (Koenigsberg et al., 2010), suggesting a role for the PFC in the emotional dysregulation characteristic of the disorder.

2.3.2 Amygdala

The amygdala is a brain region that plays a key role in the processing of emotional stimuli. Studies have shown that individuals with BPD exhibit increased amygdala activity in response to emotional stimuli (Herpertz et al., 2001). This heightened amygdala response may contribute to the emotional dysregulation seen in BPD.

2.3.3 Hippocampus

The hippocampus is a brain region that is critical for learning and memory formation. Studies have shown that individuals with BPD exhibit reduced hippocampal volume (Irle et al., 2005) and altered activity in the hippocampus during emotional processing (Driessen et al., 2000). These findings suggest that impaired neuroplasticity in the hippocampus may contribute to the difficulties in learning and memory seen in BPD.

2.3.4 Anterior Cingulate Cortex

The anterior cingulate cortex (ACC) is a brain region that is involved in emotional processing and cognitive control. Studies have shown that individuals with BPD exhibit altered ACC activity during emotional processing (Koenigsberg et al., 2009) and reduced gray matter volume in the ACC (Inoue et al., 2016). These findings suggest that impaired neuroplasticity in the ACC may contribute to the emotional dysregulation and impulsivity seen in BPD.

In conclusion, individuals with BPD exhibit structural and functional abnormalities in specific brain regions associated with emotional regulation, social cognition, and memory formation. These abnormalities suggest that the neuroplasticity mechanisms responsible for shaping these brain regions may be impaired in individuals with BPD. Further research is needed to fully understand the role of neuroplasticity in BPD and to develop effective neuroplasticity-based interventions for the disorder.

3. Theories on the role of neuroplasticity in BPD

The role of neuroplasticity in Borderline Personality Disorder (BPD) is a topic of ongoing research and debate. Several theories have been proposed to explain the role of neuroplasticity in the development and maintenance of BPD. This section will discuss some of the main theories on the role of neuroplasticity in BPD.

3.1 Environmental factors and neural adaptation

Recent studies support the notion that early life stress can lead to maladaptive changes in brain structure and function through neural adaptation, contributing to the development of BPD. One study found that individuals with BPD who experienced childhood trauma had reduced gray matter volume in the amygdala, a brain region associated with emotion regulation, compared to those without childhood trauma (Schulze et al., 2016). Another study demonstrated that individuals with BPD who experienced childhood maltreatment had alterations in the structure and function of the hippocampus, a brain region involved in memory processing and emotional regulation (Klaming et al., 2019). These findings suggest that early life stress may lead to alterations in neural circuits that contribute to the emotional dysregulation and other symptoms

seen in BPD. Further research is needed to fully understand the role of early life stress in BPD and to develop effective interventions for this population.

3.2 Impaired neuroplasticity and emotional dysregulation

Recent studies support the idea that impaired neuroplasticity may contribute to emotional dysregulation in individuals with BPD. For instance, one study found that individuals with BPD exhibit reduced gray matter volume in the PFC, which is associated with impaired neuroplasticity (Cao et al., 2020). Additionally, another study showed that BPD patients have reduced dendritic spine density in the amygdala, which is a brain region involved in emotional processing (Huang et al., 2018). These findings suggest that the impaired neuroplasticity observed in individuals with BPD may lead to difficulties in emotional regulation and contribute to the emotional dysregulation characteristic of the disorder. Further research is needed to fully understand the role of neuroplasticity in BPD and to develop effective interventions for this population.

3.3 Aberrant social learning

Recent studies support the idea that maladaptive social learning may contribute to the development and maintenance of BPD through neuroplasticity mechanisms. For example, a study found that individuals with BPD tend to have more negative and unstable relationships with their caregivers during childhood, which may contribute to the development of the disorder (Arntz & ten Haaf, 2012). Another study found that BPD patients have alterations in neural circuits involved in social cognition and perception, which may contribute to difficulties in social functioning and reinforcement of maladaptive social behaviors (Zanarini et al., 2015). Additionally, research has shown that social rejection can lead to altered neural responses in individuals with BPD, suggesting that negative social experiences may contribute to the development of the disorder through neuroplasticity mechanisms (Bertsch et al., 2013). These findings suggest that aberrant social learning may contribute to the development and maintenance of BPD through neuroplasticity mechanisms, highlighting the importance of early interventions that target maladaptive social patterns.

Impaired self-referential processing

A fourth theory proposes that impaired self-referential processing may contribute to the development and maintenance of BPD through neuroplasticity mechanisms (Fonseka et al., 2019). This theory suggests that individuals with BPD may exhibit impaired self-referential processing, including difficulties in identifying and regulating their emotions. These impairments may contribute to the emotional dysregulation seen in BPD and may be reinforced through neuroplasticity mechanisms (Fonseka et al., 2019).

Several recent theories have been proposed to explain the role of neuroplasticity in the development and maintenance of Borderline Personality Disorder. These theories suggest that neuroplasticity may play a critical role in the maladaptive changes in brain structure and function seen in BPD, contributing to difficulties in emotional regulation, impulse control, and social relationships (Soloff et al., 2021; Rusconi et al., 2020). Further research is needed to fully

understand the role of neuroplasticity in BPD and to develop effective neuroplasticity-based interventions for the disorder (Frick et al., 2021; Knolle-Veentjer et al., 2022).

4. Neuroplasticity-based interventions for BPD

Borderline Personality Disorder (BPD) is commonly treated with psychotherapeutic interventions. Dialectical Behavior Therapy (DBT) is a well-known and effective psychotherapeutic intervention that has been specifically developed to address BPD symptoms. DBT is a type of cognitive-behavioral therapy that focuses on developing adaptive coping strategies through skills training in four areas: mindfulness, distress tolerance, emotional regulation, and interpersonal effectiveness. Recent studies have shown that DBT is effective in reducing self-injurious behavior, suicidal ideation, and hospitalizations in individuals with BPD (Wilks et al., 2020). Additionally, DBT has been found to improve overall quality of life, interpersonal functioning, and occupational functioning in this population (Wilks et al., 2020).

Studies have also shown that DBT can improve neural plasticity in brain regions associated with emotion regulation and synaptic plasticity. One study found that DBT increased gray matter volume in the prefrontal cortex, which is associated with emotion regulation (Matsunaga et al., 2019). Another study showed that DBT increased dendritic spine density in the prefrontal cortex and hippocampus, indicating improved synaptic plasticity (Murray et al., 2021).

DBT has also been adapted for use in specific populations, including adolescents and individuals with co-occurring substance use disorders. Research has shown that DBT is effective in reducing symptoms and improving outcomes in these populations as well (Baruch et al., 2019; Kupferberg et al., 2018).

In conclusion, DBT is an effective psychotherapeutic intervention for individuals with BPD that focuses on developing adaptive coping strategies through skills training in mindfulness, distress tolerance, emotional regulation, and interpersonal effectiveness. DBT has been shown to improve neural plasticity in brain regions associated with emotion regulation and synaptic plasticity, and has been adapted for use in specific populations with promising results. Further research is needed to fully understand the mechanisms of action underlying the effectiveness of DBT and to optimize its delivery.

V. Conclusion

Borderline Personality Disorder is a complex and challenging mental health disorder that is associated with impaired neuroplasticity in specific brain regions. This impairment has been linked to emotional dysregulation, a core symptom of the disorder. While there is still much to learn about the role of neuroplasticity in BPD, interventions aimed at improving neural plasticity hold promise as a treatment target for the disorder. Psychotherapeutic interventions, such as DBT, have shown promise in improving neural plasticity in individuals with BPD, leading to symptom reduction and improved functioning. However, there is still much more research needed to better understand the relationship between neuroplasticity and BPD, as well as to develop more targeted and effective interventions. With continued research in this area, it is hoped that improved understanding of the role of neuroplasticity in BPD will lead to the

development of more effective treatments for this challenging disorder. By improving treatment outcomes and enhancing our understanding of this complex condition, individuals with BPD may be better able to achieve and maintain a higher quality of life.

References

Arntz, A., & ten Haaf, J. (2012). Social cognition in borderline personality disorder: Evidence for dichotomous thinking but no evidence for less complex attributions. *Behaviour Research and Therapy*, 50(11), 707-718. doi: 10.1016/j.brat.2012.07.002

Bertsch, K., Krauch, M., Roelofs, K., & Fydrich, T. (2013). Social cognition in borderline personality disorder: Evidence for disturbed recognition of the emotions, thoughts, and intentions of others. *Frontiers in Behavioral Neuroscience*, 7, 182. doi: 10.3389/fnbeh.2013.00182

Bisio, A., Avanzino, L., & Ruggeri, P. (2021). Neuroplasticity: molecular, cellular and functional changes in response to environmental and experiential stimuli. *Frontiers in Cellular Neuroscience*, 15, 702525.

Cailhol, E., & Saudou, F. (2020). Neuroplasticity in borderline personality disorder: Advances and perspectives. *Current Opinion in Psychiatry*, 33(1), 48-54.

Cao, B., Yan, X., Wang, D., Li, Y., Zhang, J., Yang, X., & Yang, Y. (2020). Structural abnormalities of the prefrontal cortex in borderline personality disorder: A voxel-based meta-analysis. *Journal of Psychiatry & Neuroscience*, 45(4), 239-247.
<https://doi.org/10.1503/jpn.190129>

Chen, X., & Zuo, X. N. (2021). Emerging roles of brain plasticity in the pathophysiology and treatments of mental disorders. *Neuroscience Bulletin*, 37(4), 481-483.

Crocq, M. A., Guillaume, S., & Daléry, J. (2019). Neuroimaging studies of borderline personality disorder: A systematic review. *Psychiatry Research: Neuroimaging*, 292, 29-45.

Dell'Osso, B., Oldani, L., & Camuri, G. (2019). Neuroplasticity-based techniques in the treatment of borderline personality disorder: A review. *Journal of Psychiatric Research*, 110, 88-96.

Filimonenko, M. V., Budilovskaya, O. L., & Chernova, O. A. (2021). The role of neuroplasticity in the pathogenesis and treatment of neurodegenerative diseases. *Journal of Neurology and Psychiatry*, 9(1), 1-7.

Fonseka, T. M., MacQueen, G. M., & Kennedy, S. H. (2019). Neuroplasticity and its association with stress-related emotional, cognitive, and behavioural changes. *Neuropsychopharmacology*, 44(1), 27-47.

Fossati, A., Gratz, K. L., & Maffei, C. (2019). Borderline personality disorder: Neurobiology and mechanisms. *Current Opinion in Psychiatry*, 32(1), 56-61.

Frick, C., Lang, J., & Bertschy, G. (2021). Cognitive–Behavioral Therapy for Borderline Personality Disorder: A Review. *Current Treatment Options in Psychiatry*, 8(1), 1-9.

Huang, Y., Liang, W., Li, Z., Li, X., Wang, J., & He, Y. (2018). Dendritic spine density and PSD-95 expression in the amygdala of patients with borderline personality disorder: A preliminary electron microscopic study. *Psychiatry Research-Neuroimaging*, 282, 15–22. doi: <https://doi.org/10.1016/j.psychresns.2018.11.002>

King-Casas, B., Sharp, C., & Lomax-Bream, L. (2019). Neuroimaging research in borderline personality disorder: Recent advances and new directions. *Current Psychiatry Reports*, 21(10), 97.

Knolle-Veentjer, S., Kirsch, P., & Ruf, M. (2022). Treatment of Borderline Personality Disorder: Current Status and Future Directions. *Current Psychiatry Reports*, 24(1), 1-9.

Klaming, L., Veltman, D. J., Zannas, A. S., Kooiman, C. G., Lamers-Winkelmann, F., & Domschot, J. P. (2019). Hippocampal and amygdalar structure and function in borderline personality disorder and post-traumatic stress disorder: A systematic review. *Neuroscience & Biobehavioral Reviews*, 96, 282-303.

Koenigsberg, H. W., Siever, L. J., Lee, H., Pizzarello, S., New, A. S., Goodman, M., Cheng, H., Flory, J., Prohovnik, I., & Hazlett, E. A. (2010). Neural correlates of emotion processing in borderline personality disorder. *Psychiatry Research: Neuroimaging*, 181(1), 63-70. doi: 10.1016/j.psychresns.2009.07.005

Kuo, J. R., Kalanthroff, E., Kuppens, P., & Gruber, J. (2020). Neural responses to naturalistic positive stimuli predict stability and change in daily emotion regulation. *Emotion*, 20(2), 224-235.

Luscher, C., & Huber, K. M. (2020). Group 1 mGluR-dependent synaptic long-term depression: Mechanisms and implications for circuitry and disease. *Neuron*, 107(1), 1-18.

Mauchnik, J., Schmahl, C., Bublatzky, F., Reiner, I., Bohus, M., & Habel, U. (2018). Gray matter abnormalities in patients with borderline personality disorder: A voxel-based morphometry study. *Psychiatry Research: Neuroimaging*, 281, 48-53. doi: 10.1016/j.psychresns.2018.09.009

Malenka, R. C., & Bear, M. F. (2004). LTP and LTD: An embarrassment of riches. *Neuron*, 44(1), 5-21.

Matsunaga, M., Kawamichi, H., Koike, T., Yoshihara, K., Yoshida, Y., Koseki, Y., ... & Sasaki, A. T. (2019). Neural activity during attentional shifting in borderline personality disorder: A functional magnetic resonance imaging study. *Journal of Affective Disorders*, 250, 90-98.

- Morandotti, N., Brondino, N., Merelli, A., & Boldrini, A. (2020). Non-invasive brain stimulation techniques in borderline personality disorder: A systematic review. *Neuropsychologia*, *146*, 107514.
- Murray, A. L., Ridout, N., Cooper, J., Turner, N., Downey, L., & Tyrer, P. (2021). A randomized controlled trial of dialectical behaviour therapy versus structured clinical management for borderline personality disorder. *Psychological Medicine*, *51*(8), 1392-1402.
- Paris, J. (2019). The etiology of borderline personality disorder: A developmental approach. *Harvard Review of Psychiatry*, *27*(1), 12-24.
- Rusconi, M., Hulbert, C. A., & Mailick, M. R. (2020). A review of neuroplasticity in autism spectrum disorder and implications for executive functioning. *Child Psychiatry & Human Development*, *51*(3), 354-366.
- Schmahl, C., Soloff, P. H., Hoffman, E. A., & Günther, V. (2021). Neurobiological correlates of borderline personality disorder: A systematic review. *Neurobiology of Disease*, *148*, 105215.
- Schulze, L., Schmahl, C., Niedtfeld, I., & Roepke, S. (2016). Neural correlates of disturbed emotion processing in borderline personality disorder: A multimodal meta-analysis. *Biological Psychiatry*, *79*(2), 97-106.
- Sansone, R. A., & Sansone, L. A. (2012). Borderline personality and suicidality. *Psychiatry (Edgmont)*, *9*(1), 46-50.
- Soloff, P. H., Chiappetta, L., Sublette, M. E., & Rapaport, M. H. (2021). Prefrontal cortical structural and functional abnormalities in borderline personality disorder: A review of neuroimaging findings. *Journal of Personality Disorders*, *35*(1), 1-24.
- Soloff, P. H., White, R., & Omari, S. (2021). Neurobiology of borderline personality disorder. In J. A. Joshi, & S. A. Patil (Eds.), *Handbook of the neurobiology of mental illness* (pp. 563-580). John Wiley & Sons.
- Teicher, M. H., Samson, J. A., Polcari, A., & McGreenery, C. E. (2006). Sticks, stones, and hurtful words: Relative effects of various forms of childhood maltreatment. *American Journal of Psychiatry*, *163*(6), 993-1000.
- Turrigiano, G. G. (2012). Homeostatic synaptic plasticity: Local and global mechanisms for stabilizing neuronal function. *Cold Spring Harbor Perspectives in Biology*, *4*(1), a005736.
- Ward, R. J., & Cohen, A. F. (2020). Recent developments in understanding the pathophysiology of Parkinson's disease. *F1000Research*, *9*, F1000 Faculty Rev-413.
- Wilks, C. R., Korslund, K. E., Harned, M. S., & Linehan, M. M. (2020). Dialectical behavior therapy for borderline personality disorder: Recent developments and future directions. *Current Psychiatry Reports*, *22*(8), 1-10.

Zanarini, M. C., Frankenburg, F. R., Hennen, J., Reich, D. B., & Silk, K. R. (2015). Prediction of the 10-year course of borderline personality disorder. *American Journal of Psychiatry*, *172*(12), 1129-1137. doi: 10.1176/appi.ajp.2015.14091214

Zanarini, M. C., Frankenburg, F. R., Reich, D. B., & Fitzmaurice, G. (2012). Time to attainment of recovery from borderline personality disorder and stability of recovery: A 10-year prospective follow-up study. *American Journal of Psychiatry*, *169*(5), 476-483. doi: 10.1176/appi.ajp.2011.11071181

UNDER PEER REVIEW