

Original Research Article

Effect of Substitution of Sorghum Grain by Teff (*Eragrostis tef*) Grain on the Performance of Broiler Chicks

Abstract: This study aimed to determine the effect of substitution of sorghum grain with teff grain (*Eragrostis tef*) on the performance of broiler chicks. The teff plant has been grown in Ethiopia and is used to prepare a kind of food called injera. The seeds were brought from the Sudanese Ethiopian Border. A total of 160 one-day old, unsexed broiler chicks (Ross 308) were used. The birds were distributed randomly into 16 pens (10/pen) as replicates, in a completely randomized design. The experimental diets were formulated by substitution of sorghum grain with teff grain at a level of 0, 10, 15 and 20. The chemical constituents of teff grain were measured by two methods (AOAC) and (NIR) Near-Infrared spectroscopy. The parameters measured were feed intake (FI), body weight gain (BWG), feed conversion ratio (FCR), carcass weight (CW), weight of internal organs (liver, gizzard, pancreas, neck and heart), and serum contents (cholesterol, glucose and triglyceride). The CP percentage was 7.96 % and 15.5 % for AOAC and NIR procedure respectively, the starch content was higher when determined by NIR procedure (77.05%) than AOAC procedure (65.65%), while the fiber content was found to be higher in AOAC procedure (8.61%) than NIR procedure (2.61%). For chick's performance, the birds fed 20 % teff consumed more feed (3403.0g), gained more weight (2040.5g), had the highest carcass weight (1733.8) and had the best FCR (1.66) than the control. For blood biochemical, the group of birds that fed 20 % teff grain recorded the highest value of Glucose (113.0) and the lowest value of triglyceride (107.0) and Cholesterol (128.8). It could be concluded that the teff grain is superior to sorghum in terms of chemical constituents and broiler chicken's performance and could be a good substitution for the sorghum if it is established in Sudan.

Keywords: *Eragrostis tef* grain, Sorghum Grain, Broiler Performance

Introduction

The global dominance of the four crops, maize, wheat, rice, and sorghum in the current food system has raised an alarm. Heavy dependence on these crops can limit the capacity to deal with food shortages and dietary imbalance. The poultry industry has suffered more than any other livestock industry as a result of inadequate supply and high cost of feed ([1], [2], [3]) and feed cost is expected to continue in the upward swing [4]. Cereal grains constitute the major sources of energy in poultry diets in the tropics [5]. 51 % of sorghum crop is used to feed livestock while 49% is for human food and other uses [6]. This aggressive competition between livestock and humans requires a serious diversification of energy feedstuff for livestock. Introduction of a new grain crop to the environment of Sudan may assist in finding alternatives

feed resources for poultry especially when the main crop sorghum is scarce. There has been increasing interest in grain teff due to its perceived greater nutritional quality compared to other grains ([7], [8]). Teff (*Eragrostis tef*) is a tropical cereal that has its center of origin and diversity in the northern Ethiopian highlands from where it is believed to have been domesticated ([9], [10]). Teff is a minor cereal crop worldwide, whereas in Ethiopia, it is a major food grain, mainly used to make injera, a traditional fermented Ethiopian pancake. In other countries like Australia, South Africa, and United States, it is principally used as a forage crop for animal feed. There are some data available regarding teff grain as a human food source, but there is little information available for teff as an animal feedstuff, particularly for monogastric animals such as broiler chickens. Therefore, the objectives of this study are to determine the effect of gradual substitution of sorghum grain by grain teff on the performance of broiler chick.

Material and Methods

Experimental site:

This experiment was carried out in the premises enclosure of poultry unit of the Faculty of the Animal Production University of Khartoum, Shambat (Khartoum north). During the period between 13th November to 25 December 2022.

Housing:

This experiment was conducted in a naturally ventilated, open-sided, deep-litter poultry house, with a concrete floor. The house was partitioned into 16 pens each of them one-meter square with enough working space allowance.

Experimental birds and Diets:

A total of 160 one-day-old unsexed commercial broiler chicks (Ross308) were weighted and randomly divided into (4) groups of 40 chicks, each group was further subdivided into four replicated with 10 chicks. The average of the initial weight of chicks was 37g. The experimental diets were formulated from local ingredients except the imported super concentrate to supplement the essentials amino acids (lysine, methionine, vitamin and minerals).

All groups were fed the same commercial pre-starter feed for the first week. In the second week, the first group (control) was fed a basal diet with no teff containing. The other groups were fed the basal diet with (10%), (15%) and (20) %. The formulation and calculation of experimental diet is shown in Table (1).

Table (1) Ingredient composition of experimental diets as percent (%)

Ingredients	Control	10 %	15%	20%
Sorghum	62.5	56.25	53.125	42.5
GNC	30	30	30	30
Super Concentrate*	5	5	5	5
TEFF	0	10	15	20
OIL	2.5	2.5	2.5	2.5

TOXIN BINDER	0.1	0.1	0.1	0.1
LIME STON	0.2	0.2	0.2	0.2
DCP	0.1	0.1	0.1	0.1
TOTAL	100.4			100.4

Management and Medications:

The flock was kept and reared under the same condition during the winter season. feed and water were provided ad libitum. Prebiotic with vitamin A, E and Se (one item) was administrated in the water during the first week to avoid the expected stress, and repeated after each vaccination for 3-5 days. All Chicks were vaccinated against infectious bronchitis (IB) and Newcastle (ND) at the first week of age as one dose, vaccinated against Gamboro diseases (Infectious Bursal disease) at the second week of age and also vaccinated against ND at the third week of age. the vaccines and prebiotic were given in drinking water.

Parameters Measurement:

Feed intake and body weight gain were measured weekly. Mortality rate was recorded throughout the experimental period. At the end of the experiment, four birds from each replicate were manually slaughtered. Birds were then scalded, manually plucked and washed.

Internal Organs:

The hot carcass weight was recorded. Internal organs were weighed mainly, liver, gizzard, proventriculus, pancreas, neck and heart.

Blood Parameters:

On the slaughter day, blood samples were collected from four randomly selected birds from each cage to determine serum cholesterol, triglycerides and glucose contents using commercial kits.

Chemical Analysis:

Proximate Analysis:

The proximate components of the sample were determined, Ash, EE, and CP according to AOAC methods [11].

3.8.2 Near-infrared microscopy (NIR) analysis

The sample of teff grain was measured by the means of Near-infrared microscopy (NIR) analysis. The FT-NIR spectra were scanned using the Thermo Antaris II FT-NIR spectrometer (Thermo Scientific Inc., Madison, WI, USA) prepared with an absorbance accessory.

Statistical Analysis:

The data subjected to analysis of variance by Statistical Packages for Social Science (SPSS) software program version 21. Using one-way ANOVA, the results shown as mean and their errors, mean separation done by Duncan Multiple Range Test according to [12] and significance different set ($P < 0.05$).

Results and Discussion

Table (2) noted the chemical composition of teff grain using two methods of analysis. The two methods were AOAC procedure and NIR procedure. The results showed significant differences between the two procedure for all parameters. The CP percentage was found 7.96 % and 15.5 % for AOAC and NIR procedure respectively. [13] stated that the average crude protein content of teff grain is in the range of 8 to 11 percent. [14] reported that the total protein content of teff grain was found to be 9.37%. For crude fat, the result was found 1.58 and 3.3 for the two methods respectively. This result was in the line of the results obtained by [15] who noted that the lipid contents of teff grain flour was 4.4%. The starch content of teff grain determined by NIR (77.05%) is higher that AOAC procedure (65.65%). [16] reported that complex carbohydrates make up 80 percent of teff grain, it has a starch content of approximately 73 percent making teff a starchy cereal. Moreover, [17] reported that the total carbohydrate content of teff grain was reported to be 85.6% with starch content ranging from 74 to 75.5%. lower results were obtained by [18] who stated that the proximate analysis of teff grain revealed the high nutritive value of the seeds carbohydrates (57.27%). The crude fiber content was found to be higher in AOAC procedure (8.61%) than the NIR procedure (2.61%). Comparable results were obtained by [19] who reported that the crude fibre content in teff (8.0%) is far higher than when compared to some fruits, nuts, pulses and cereals such as corn and rice. [16] Stated that the crude fiber, total and soluble dietary fiber content of teff is several folds higher than that found in wheat, sorghum, rice, and maize, which could be attributed to the factor that whole grains have higher fiber content than decorticated ones.

Table (2) Chemical Composition of *Eragrostis teff* Using Two Methods of Analysis

Parameters %	NIR Analysis	Proximate Analysis	P- value
DM	92.83 ^a	89.05 ^b	0.024
CP	15.50 ^a	7.96 ^b	0.001
CF	2.61 ^b	8.61 ^a	0.000
FAT	3.30 ^a	1.58 ^b	0.006
Ash	1.52 ^b	5.21 ^a	0.000
NFE	77.05 ^a	65.75 ^b	0.001
Me/Kg	1.38 ^b	2.617 ^a	0.000

Means with dissimilar superscripts in a row varied significantly at (P <0.05), while the same letters are not significant

The effect of the substitution of graded levels of teff grain with sorghum grain on the feed intake of broiler chicks is shown in table (3). The data showed that there was no significant difference between treatments except in week six which showed a significant difference (P≤0.05), in all weeks there is a gradual increase in feed intake with the increase in the level of substitution. This increase in feed intake may be due to the superiority of teff grain over sorghum grain, firstly, the teff grain is a starchy cereal with starch content accounting for 75% [16], with amylose content ranging from 20 to 30% [17], secondly, the protein content

of the teff grain is superior to sorghum grain in term of quantity and quality, some researchers like [18] stated that the protein content of teff grain may reach 20 % and in this study, the protein content is found to be 15 % which is far higher from sorghum (10.48%) reported by [20], moreover, the quality of teff protein is better than sorghum, [13] stated that Teff's fractional protein composition revealed that the gluten (45 %) and albumins (37 %) are the major protein storages while prolamins are a minor constituent (less than 12 %). Prolamine is a poor quality protein with low digestibility [18], in addition to that, the amino acid composition in teff is well-balanced, a relatively high concentration of lysine amino acid is found in teff [21]. [22] reported that many amino acids that are higher in teff, such as lysine, methionine and tryptophan, are considered deficient in sorghum grain, third, comparative mineral content analysis of teff grain showed that P, Mg, Mn, and Cu are present in higher concentrations in the grain than in other common cereals like maize and wheat [23]. [9] reported that teff has the highest iron content of all cereals and contains more calcium, copper, zinc, aluminum, and barium than winter wheat and barley. This result is similar to the results obtained by [24] who found that the average daily feed intake/bird and cumulative feed intake/bird during the entire period was 79.4g and 3335.9g for birds fed local diet at Ethiopia.

Table (3) Effect of Substitution of Graded Levels of *Eragoestis teff* on Weekly Feed Intake (g) of Broiler Chicks

Weeks	CONTROL	10%	15%	20%	p-value
WK2	249.0 ^a	270.1 ^a	245.5 ^a	270.2 ^a	0.330
WK3	602.9 ^a	594.7 ^a	585.8 ^a	589.8 ^a	0.855
WK4	272.6 ^a	313.6 ^a	301.7 ^a	318.2 ^a	0.357
WK5	978.1 ^a	992.1 ^a	987.7 ^a	993.8 ^a	0.543
WK6	1066.2 ^b	1212.2 ^a	1216.3 ^a	1230.7 ^a	0.003

Means with dissimilar superscripts in a row varied significantly at ($P < 0.05$), while the same letters are not significant.

Table (4) noted that the effect of the substitution of different levels of *Eragoestis teff* grain on the weekly body weight. No significant effect was observed between different treatments. The birds fed 20 % teff recorded the highest body weight, while the least body weight was obtained by the birds fed the control diet. The group of birds fed 20% *Eragoestis teff* consumed more feed and gained more weight which indicates the efficiency utilization of the feed. These results are similar to the results obtained by [25] who found that chicks receiving 10% teff as an ingredient diet had significantly heavier body weights than all other treatments. Moreover, chicks receiving the 10% whole teff diet were also significantly heavier than the control with corn-soy based diet.

Table (4) Effect of Substitution of Graded Levels of *Eragoestis teff* on Weekly Body Weight (g) of Broiler Chicks

Weeks	CONTROL	10%	15%	20%	P- value
WK1	127.5 ^a	130.0 ^a	130.0 ^a	130.0 ^a	0.426
WK2	313.6 ^a	367.0 ^a	333.5 ^a	351.2 ^a	0.273
WK3	629.1 ^a	703.3 ^a	690.1 ^a	684.2 ^a	0.653
WK4	934.0 ^a	1021.6 ^a	1002.6 ^a	1029.2 ^a	0.279
WK5	1481.0 ^a	1520.0 ^a	1551.0 ^a	1556.2 ^a	0.684
WK6	1834.7 ^b	1957.5 ^{ab}	2017.5 ^a	2077.5 ^a	0.007

Means with dissimilar superscripts in a row varied significantly at ($P < 0.05$), while the same letters are not significant.

The weekly weight gains as affected by the different substitutions of *Eragrostis teff* grain at the expense sorghum grain is illustrated in table (5). no significant difference was observed in all weeks except in week 6 which showed a significant difference ($P \leq 0.05$). The value of the weight gains in week six was increased with the increase in the level of substitution which was found to be 315.5, 335.3, 356.6, and 333.0 for the control, 10 %, 15 %, and 20% g substitution respectively.

Table (5) Effect of Substitution of Graded Levels of *Eragrostis teff* on Weekly Body Weight (g) of Broiler Chicks

Weeks	CONTROL	10%	15%	20%	p-value
WK1	90.0 ^a	93.0 ^a	93.0 ^a	93.0 ^a	0.426
WK2	186.1 ^a	237.0 ^a	203.5 ^a	221.2 ^a	0.277
WK3	315.5 ^a	335.3 ^a	356.6 ^a	333.0 ^a	0.855
WK4	304.8 ^a	318.2 ^a	312.5 ^a	345.0 ^a	0.939
WK5	547.7 ^a	498.3 ^a	548.3 ^a	527.0 ^a	0.790
WK6	353.0 ^b	437.7 ^{ab}	466.5 ^a	521.0 ^a	0.020

Means with dissimilar superscripts in a row varied significantly at ($P < 0.05$), while the same letters are not significant

The average weekly feed conversion ratio as affected by different levels of substitutions of sorghum grain by teff grain is shown in table (6). No significant difference was found between treatments. In comparison to the feed consumption and weight gain, the birds which consumed more feed were heavier and had a less feed conversion ratio indicating that the heaviest birds are more efficient in utilizing the diet than lighter weight suggesting that the diet had high energy. This results were different to the results obtained by [25] who stated that the 10% whole teff and the teff as ingredient fed chicks had significantly poorer

conversions than control corn soy fed chicks. It was observed that some chicks receiving the 10% ground teff treatment had incidences of fine, flour-like feed adhering to their beaks.

Table (6) Average weekly feed conversion ratio as affected by different treatments

Weeks	CONTROL	10%	15%	20%	p-value
WK2	1.38 ^a	1.14 ^a	1.25 ^a	1.22 ^a	0.391
WK3	1.95 ^a	1.81 ^a	1.71 ^a	1.81 ^a	0.826
WK4	0.94 ^a	1.00 ^a	1.40 ^a	0.93 ^a	0.626
WK5	1.80 ^a	2.01 ^a	1.85 ^a	1.90 ^a	0.689
WK6	3.09 ^a	2.79 ^a	2.61 ^a	2.41 ^a	0.171

Means with dissimilar superscripts in a row varied significantly at ($P < 0.05$), while the same letters are not significant.

The results of cumulative feed intake as illustrated in table (7) showed that there was a significant difference ($P \leq 0.05$) between different treatments. The highest feed intake (3403g) was obtained by 20 % teff substitution while the lowest feed intake was recorded in the pure sorghum grain. The data also showed that the final body weight was affected significantly ($P < 0.05$) by the addition of teff grain, the trend is identical to what found in the feed intake, the highest final body weight (2040.5g) was obtained by 20 % teff substitution while the lowest body weight was recorded in the pure sorghum grain. Consistently, the effect of substitution of graded levels of sorghum grain by teff grain on cumulative feed conversion ratio was found to be in significant, the group of birds fed diet containing 20% teff consumed more feed, gained more weight and had the best feed conversion ratio, while the group of birds fed the diet with no teff consumed less feed, gained less weight and had poorer feed conversion ratio.

Table (7) The Effect Substitution Sorghum Grain with *Eragrostis tef* Grain on Overall performance of broiler Chicks.

Treatment	FI	BW	FCR
Control	3168.9 ^b	1797.7 ^b	1.76 ^a
10%	3382.8 ^a	1920.5 ^{ab}	1.76 ^a
15%	3337.2 ^a	1980.5 ^a	1.68 ^a
20%	3403.0 ^a	2040.5 ^a	1.66 ^a
p-value	0.021	0.007	0.057

Means with dissimilar superscripts in a column varied significantly at ($P < 0.05$), while the same letters are not significant

FI = Feed Intake BW= Body weight FCR= Feed Conversation Ratio

Table (8) noted the effect of the substitution of graded levels of *Eragorestis tef* on the carcass parameter. For carcass weight, there was a significant difference between parameters ($P \leq 0.05$). The ranking of carcass weight from the smallest to the largest was found to be as follows: control < 10 % < 15 % < 20 %. This result clearly indicated that the application of 20 % addition of teff grain yielded more meat than the other treatments. For the weight of the internal organs, the results showed that no significant difference was observed for all parameters studied except the weight of the neck ($P \leq 0.05$). Numerically, the weight of the all internal organs under study increased by increasing the percentage of teff except proventriculus wherein the control is higher than treatment.

Table (8): Effect of Substitution of Graded Levels of *Eragorestis tef* on the Carcass Parameter

Traits	CONTROL	10%	15%	20%	p-value
Carcass Weight	1118.7 ^b	1433.3 ^{ab}	1548.8 ^{ab}	1733.8 ^a	0.005
Liver	40.0 ^a	41.7 ^a	50.0 ^a	43.8 ^a	0.841
Pancreas	17.5 ^a	18.3 ^a	17.5 ^a	18.8 ^a	0.984
Neck	81.3 ^c	88.3 ^{bc}	101.3 ^{ab}	107.5 ^a	0.016
Gizzard	33.8 ^a	35.0 ^a	38.8 ^a	35.0 ^a	0.624
Proventriculus	12.5 ^a	8.3 ^a	8.8 ^a	10.0 ^a	0.262
Heart	8.8 ^a	8.3 ^a	11.7 ^a	10.0 ^a	0.313

Means with dissimilar superscripts in a column varied significantly at ($P < 0.05$), while the same letters are not significant

The effect of the inclusion of graded levels of teff grain on blood biochemical on broiler chicks was demonstrated in table (9). For blood glucose no significant difference was found between the control and different levels of teff grain, although 10 % and 20 % inclusion recorded higher value of blood glucose than the control. For the content of blood triglyceride, the results showed a significant decrease ($P \leq 0.05$) as the percentage of teff increased, the same trend was observed for the blood cholesterol.

Table (9): Effect of Substitution of Graded Levels of *Eragorestis tef* on the Blood biochemical.

Concentration	Traits		
	Glucose	TG	Cholesterol

Control	110.0 ^a	179.3 ^a	130.3 ^b
10%	117.5 ^a	106.8 ^b	131.5 ^b
15%	109.3 ^a	107.8 ^b	150.5 ^a
20%	113.0 ^a	107.5 ^b	128.8 ^b
p-value	0.790	0.001	0.042

Means with dissimilar superscripts in a column varied significantly at ($P < 0.05$), while the same letters are not significant

Conclusion:

Based on the current study it could be concluded that *Eragrostis tef* grain is superior to sorghum grain which could be indicated in feed intake, body weight gain, and feed conversion ratio. Moreover, the carcass meat yield and the internal organs weight were improved by the inclusion of teff grain. The blood serum constituents were also affected by the incorporation of the teff grain in the diet. Teff grain if establish in the environment of Sudan could be a good alternative of sorghum grain in livestock feeding.

References

- 1- Hill FW (1989). Poultry Nutrition and Nutrient Requirements. In: Nutrition of Animals of Agricultural Importance Part 2: Assessment of and Factors Affecting Requirements of Farm Livestock. Cutbertson D (ed.), pp. 1137 – 1180.
- 2- Mtimuni JP (1995). Ration Formulation and Feed Guides. Likuni Press and Publishing House, Lilongwe, Malawi, p. 1.
- 3- Lepleaieur M (ed) (2004). Poultry farming - A disease called competition. In: SPORE Magazine, 114: 4 – 5.
- 4- Conolly A (2012). Seminar presentation on “Pushing the boundaries – Performance and profitability”. Poultry International Magazine, Mark Clement (ed). January 2012. p.18 www.WATTAgNet.com
- 5- Oluyemi JA, Roberts RA (2000). Poultry Production in Warm Wet Climates. Macmillan Press, London. 2nd Edition, p. 692.
- 6- Maunder B (2002). SorghumThe global grain of the future. Retrieved November 30, 2004 from <http://www.Sorghumgrowers.com/maunder.htm>
- 7- Bhargava, A., Sudhir, S., & Deepak, O. (2006). Chenopodium quinoa — An India perspective. Industrial Crops and Products, 23, 73–87. doi:10.1016/j.indcrop.2005.04.002
- 8- Hitomi, A., Yi-Chun, C., Hanjun, T., Mayumi, S., Katsumi, W., & Oshio, M. (2002). Food components in fractions of quinoa seed. Food Science and Technology Research, 8, 80–84. doi:10.3136/fstr.8.80
- 9- Ketema, Seyfu. (1997). Tef, *Eragrostis Tef* (Zucc.) Trotter. Promoting the Conservation and Use of Underutilized and Neglected Crops series no. 12. Gatersleben: Institute of Plant Genetics and Crop Plant Research & Rome: International Plant Genetic Resources Institute.

- 10- Demissie, Abebe. (2001). "Teff Genetic Resources in Ethiopia." *Narrowing the Rift: Teff Research and Development. Proceedings of the "International Workshop on Teff Genetics and Improvement"*, Debre Zeit, Ethiopia, 16-19 October 2000. 27-31.
- 11- AOAC (1990). *Official Method of Analysis Association of Official Analytical Chemists* Pp: 66-88 .15 th. Edition Washington , D.C. USA.
- 12- Steel, R. G. D., and J. H. Terrie. 1997. *Principles and Procedures of Statistics. A Biometric Approach*. 2nd ed. McGraw-Hill, New York, NY.
- 13- Tatham AS, Fido RJ, Moore CM et al. Characterisation of the major Prolamins of teff (Eragrostis Tef) and finger millet (Eleusine Coracana). *J Cereal Sci.* 1996; 24(1):65-71
- 14- Gebru, Y. A., Sbhata, D. B. ,1 and Kim, K (2020) Nutritional Composition and Health Benefits of Teff (Eragrostis tef (Zucc.) Trotter) *Journal of Food Quality* ID 9595086, 6 pages.
- 15- Hager, A.-S. Czerny, M. Bez, JZannini, E. and Arendt, E. K. (2013) Starch properties, in vitro digestibility and sensory evaluation of fresh egg pasta produced from oat, teff and wheat flour," *Journal of Cereal Science*, vol. 58, no. 1, pp. 156–163, 2013.
- 16- Bultosa Geremew (2007). Physicochemical Characteristics of Grain and Flour in 13 Teff [Eragrostis tef (Zucc.) Trotter] Grain Varieties. *Journal of Applied Sciences Research*, 3(12): 2042-2051
- 17- Abebe W. and Ronda, F. (2014) Rheological and textural properties of tef [Eragrostis tef (Zucc.) Trotter] grain flour gels," *Journal of Cereal Science*, vol. 60, no. 1, pp. 122–130, 2014.
- 18- Nascimento, K. D. D, Paes, S. N. D., Oliveira, I. R. D. IReis, I. P. I, and Augusta, V. M. (2018)Teff: Suitability for Different Food Applications and as a Raw Material of Gluten-free, a Literature Review. *Journal of Food and Nutrition Research*, Vol. 6, No. 2, 74-81
- 19- Saturni L, Ferretti G, Bacchetti T. The gluten-free diet: Safety and nutritional quality. *Nutr.* 2010; 2(1):16-34.
- 20- Etuk E. B., Ifeduba, A. V., Okata U. E., Chiaka I., Okoli, Ifeanyi C., Okeudo N. J., Esonu B. O., Udedibie A. B. I. and Moreki, J. C. (2012) Nutrient Composition and Feeding Value of Sorghum for Livestock and Poultry: a Review *J Anim Sci Adv* 2012, 2(6): 510-524
- 21- Akansha, Sharma, K., and Chauhan, E. S. (2018) Nutritional composition, physical characteristics and health benefits of teff grain for human consumption: A review. *The Pharma Innovation Journal* 2018; 7(10): 03-07
- 22- Coleman, J. M. (2012) *Assessing the Potential Use of Teff as an Alternative Grain Crop in Virginia*. M. Sc. Thesis university of Verginia.
- 23- Forsido, S. F. Rupasinghe, H. P. V. and Astatkie, T. (2013) Antioxidant capacity, total phenolics and nutritional content in selected ethiopian staple food ingredients," *International Journal of Food Sciences and Nutrition*, vol. 64, no. 8, pp. 915–920.

- 24- Moges, F. Amare, A and Dawud Ibrahim, D. (2014) Demonstration and Performance Evaluation of Broilers Using Locally Processed Feed in Bahir Dar Zuria District, Amhara Region, Ethiopia. World Vet J, 4(4): 42-47, December 25, 2014
- 25- Brockotter, F. (2017) Teff grain of broiler feed. Poultry world 33(2), 29-30, 2017

UNDER PEER REVIEW