

**FRESHWATER PRAWNS DIVERSITY (CRUSTACEA: DECAPODA)
IN THE RIVER KOSHI, NORTH BIHAR**

ABSTRACT

Introduction: The river Koshi is one of the major tributary of river Ganga, which originated from the northern slopes of the Himalayas and flows southwards through north Bihar with draining of the Ganga basin near Bhagalpur Bihar.

Objectives: Diversity and abundance of freshwater prawns were studied from January 2020 to December 2022 in the Koshi river of district Katihar north Bihar, India.

Methodology: The prawn was collected from the bank of Koshi river with the support of fishermen at Kursela Katihar district, north Bihar and preserved in 5% formalin for further study.

Results: Koshi river is also known as Saptakoshi for its seven upper tributaries, that transports a large amount of nutrient and perhaps encourages species prosperity both culturally and individually. The efficiency of water is significant and therefore the freshwater prawns are divers. Prawn is a very important foodstuff in developing countries, due to high protein content and its other nutrition values. In present study we identified & reports commercially and physically important total four species of prawn belonging to *Macrobrachium* genus in the river Koshi around Kursela, Katihar district north Bihar. They are namely *Macrobrachium gangeticum*, *M. gandaki*, *M. tiwari* and *M. lamarrei*. The composition of prawn species among which *M. lamarrei* and *M. tiwari* were dominant species throughout the year. Whereas the catch of *M. gangeticum* and *M. gandaki* were restricted during June to September.

Conclusion: The current investigation important to the presence of commercially important and environmentally sustainable inhabitants available in the river Koshi, Kursela Katihar, North Bihar. The suggests to needs appropriate management and conservation strategies to prevent the further loss of populations from this site. The details of data collected are in the flesh in this communication.

Keywords: Shrimp, Taxonomy, *Macrobrachium species*, Fishery, Biology.

1. INTRODUCTION

The Koshi is one of the major perennial river in the peninsular India, originated from Nepal and flows southwards through north Bihar with draining of the Ganga basin near Bhagalpur, district Bihar. Its facts, river Koshi having unstable nature attributed to the heavy silt carried

during the flood (monsoon) season and also transports a large amount of nutrient, perhaps encourages species prosperity both culturally and individually. Freshwater prawns play a crucial role in sustaining the food web by recycling nutrients and energy from decay matter as well as structuring and functioning of the ecosystem and also play an important role as scavengers [1,2]. Freshwater prawns are present in all biogeographically regions with the exception of Antarctica, however, the Oriental region reported as harbours and the majority of the prawn species [3,4]. As a matter of facts, large and medium sized freshwater prawns, form economically important group as most of them are harvested from the wild or culture for food as well as aquarium trade [5]. Overall diversity and abundance of freshwater prawns is rapidly declining or under risk category mainly due to competition on from invasive species, change in climate and commercial developmental projects [6,7], quality of water and anthropogenic pollution can be determined by presence of species and their abundance [8,9]. Whereas, [10] reported four species of freshwater prawns in the river Ganga up to Kanpur its availability year around during (1956). However, two major *Macrobrachium* species *M. gangeticum* and *M. malcolmsonii* was reported in limited months and there is significant reduction of the natural stock at middle stretch of the river Ganga documented by [6,11].


It facts, North Bihar is donated with number of perennial rivers with draining and ultimately part of Ganges river system. The Koshi river is one of them and also known as 'Sorrow of Bihar' that flow in north east region of Bihar as the annual floods effect of fertile agricultural lands thereby disturbing the rural economy and also abundant fishery resources. Since freshwater fauna especially prawn resources are at risk of being carried away from their respective habitats [12,13]. The size group, sex-ratio, arrangement, spreading, larval biology and incidence of freshwater prawn in the river Ganga has been studied with recorded by [14,15,16,30]. The information about Koshi river prawns data is not available, therefore, it is necessary to have a reliable data based on surveys and identification of prawns from this natural habitats.

Thus the aim of the present study is to documented the diversity and abundance of *Macrobrachium species* in the lower stretch of the river Koshi near Kursela Katihar, south Bihar. These records are prerequisite for further biological studies related to hatchery seed production and cultural aspects.

2. MATERIALS AND METHOD

2.1 Study area: The Koshi region is sedimentary zones appearances like fan and covering 180 km long and 150 km wide northeast Bihar and eastern Mithila to the Ganges,. It shows evidence of twelve major lateral channels shifting exceeding 120 km during the past 250

years: (www.mapsofindia.com). The rivers, which flowed near Purnea in the 18th century, presently it flows west of Saharsa, & South of Katihar district of North Bihar (Map-1). In the Koshi region, Kursela, Purnea, Araria, Forbesganj, Supaul, Madhepura and Khagaria are the main flood affected cities in North Bihar.



Map-1. Showing alluvial flood affected areas of Koshi river (Source: www.mapsofindia.com)

2.2. Step for the collection of the species:

In the present study freshwater prawns specimen were collected from the Koshi river of near between the road Bridge and the Rly Bridge on Koshi river at Kursela Katihar district, north Bihar. They were collected from the bank with the support of fishermen. The each species of prawn individual was preserved in 5 percent neutralized formalin for further identification. Further studies on the biology, classification and allied aspects of the collected prawn specimens as live and preserved were brought at Shellfishes Biodiversity Lab, at Bhola Paswan Shastri Agricultural College (BAU, Sabour), Purnea, Bihar, India.

2.3. Identification and measurement : The species was identified in consultation with related literature previously published by [13,17,18,19,20]. The total length of individual specimen of all the group were recorded from the tip of the rostrum to the tip of the telson., tip of the chalet leges to tip of the telson were recorded to nearest millimetre. The appendix masculine was clearly visible only in the male specimens of 50 mm size and above which formed the base of identification of sex in the commercial catches. The measurements of these species were compared with the *Macrobrachium species*. Morphometric measurements collection are given in (Table -1 & plate a,b,c&d).

3. RESULTS AND DISCUSSION

Populace constraints are essential in order to understand the well-being of any fishery to determine proper management measures. Structural identification of the prawn specimen demonstrated the presence of four *species* of commercial important from the river Koshi near Kursela, Katihar north Bihar. The identification of prawn belonging to genus *Macrobrachium* (Bate) is mainly based on the proportion of different segments of the second chelipeds in addition to the characterization of rostrum [14]. All the specimens have been identified up to the species level & described and body measured in millimetre are predicted in the (Table-1 & Plates a,b,c&d).

Table: 1. A comparative morphological study of *Macrobrachium species*

S. N.	Characters	<i>M. gangeticum</i>	<i>M. gandaki</i>	<i>M. tiwari</i>	<i>M. lamarreii</i>
1	Rostrum extended	Shorter & not beyond antennal scale	Short & not reaching as far as antennal peduncle	Elongated reaching upto or beyond antennal scale	Rostrum long sword shaped reaching past antennal scale
2	Basal crest deviation	Basal crest highly elevated to forma keel	Basal crest moderately elevated	Basal crest moderately elevated	Basal crest moderately elevated
3	Distal end of rostrum	Distal end directed forwards	Distal end directed forwards	Distal end directed forwards	Distal end directed forwards
4	Rostral formula	9-11/4-6	8-10/2-3	9-10/2-0	8-9/10-12
5	Second pereopods	Largest, equal & larger than body length	Largest, & unequal	Larger, equal & shorter to body length	Smaller, equal & larger than other pereopods
6	Maximum size	200 mm	75 mm	85 mm	80 mm

3.1. *Macrobrachium gangeticum*: This is the largest representative of freshwater prawn in the Koshi region. The male attain maximum growth of 200 mm and female attains maximum 180 mm in the Koshi river, whereas, [15,16,29,20] reported regarding *M. gangeticum* male attain maximum growth of 215 mm and female attains maximum 200 mm in the middle stretch of the river Ganga near round Patna, Bihar. These are called ‘Gangetic river prawn’ it also known as *M. birmanicum choprai* (Tiwari) it is yellowish brown, it has an elevated dorsal rostral crest. Rostral formula of the species was found with 9-11 dorsal teeth and 4-6 ventral. The rostrum of this species was also found elongated short extended & reached upto the length of the antennal peduncle scale, which is highly convex, and slightly elevated upturned. The dorsal rostral teeth on elevated crest usually found closely set up with each

other, whereas, ventral teeth were separated from each other. The second pair of pereopod (chelate lag) is longer than body also having equal size and carpus as long as merus. The prawns are distributed in the Ganga and Brahmaputra river system detail presented in the (Table: 1 & Plate- a,b,c&d).

3.2. *Macrobrachium gandaki*: This is the third largest medium sized prawns groups in this region. The male and female attains a maximum growth of 75 and 65 mm. Whereas, [30] reported a new species of *Macrobrachium gandaki* (Bate, 1868) in middle stretch of the river Ganga India, sized ranging from 60-70 mm in total length (tip of the rostrum to tip of the telson) They are locally known as ‘Gandak river prawn’ possesses un-equal size of pair of 2nd chelate legs. Rostrum moderately short, elevated, not reaching up to antennal peduncle. The distal end of rostral tip directed forward upper margin moderately convex, with 8-10 teeth placed at distal half of the rostrum. Small setae were present the teeth of both dorsal & ventral margin. The chelipeds are either the left of right in male unequal, one big, strong and another smaller, body colour is translucent blue. Shape of Telson were found triangular pointed at the tip, slender with a dorsal spines, antenna segmented. It is distributed middle stretch of the river Ganga, Gandak and Koshi. *Macrobrachium gandaki* having black band on the abdominal dorsal part were characteristics to the present species.



Plate-a. *Macrobrachium gangeticum*



Plate-b. *Macrobrachium gandaki*



Plate-c. *Macrobrachium tiwari***Plate-d. *Macrobrachium lamarrei***

[21] reported two species of prawn, whereas, [22] reported 4 species of prawn namely *Macrobrachium gangeticum*, *M. malcolmsonii*, *M. lamarrei* and *M. lamarrei lamarrei* in the middle stretch of the river Ganga around Patna, Bihar. The matter of facts, *M. gangeticum* is aquaculture importance and developed its hatchery technology (outdoor & indoor) by the different researcher like [23, 24,] its need to explore this technology from hatchery to pond condition (lab to land programme). It is facts, *M. gangeticum* reported third largest freshwater prawn in India and apparently less in natural system in the Gangetic river [6] however this species needs further observation particularly on aspects like conservation and culture in pond condition, detail presented in the (Table: 1 & Plate-a,b,c&d).

3.3. *Macrobrachium tiwari*: This is comes under second largest prawn group in the Koshi region. This species are capture to be marketable importance in different state of India. [25] reported also having aquaculture potential, but unutilized and produced in hatchery system. These are relatively medium / small prawns, maximum size of male is 65 mm and female 85mm which are looking in pale grey colour. The rostrum of this species is elongated reaching up to beyond the length of the antennal peduncle. It may be curved or slightly curved upwards. The 2nd pair of chelate legs shorter that the body length. Antenna having dot marking looks like segmented. Uropod consist of accessing spine on the telson. Black band on lateral side of the rostrum from base to tip. Entire elongated pair of cheliped legs in which carpus is distinctly longer than merus. However, the percentage composition in fish landing is comparatively less than *M. lamarrei*, detail presented in the (Table: 1 & Plate-a,b,c&d).

3.4. *Macrobrachium lamarrei*: Length of the rostrum of this decapodis long sword shaped, tip sharper and unturned, reaching past the antennal scale. Distal end of rostrum is slender and turned upwards. the most important features of this species is the wide irregular gap between first two and third dorsal teeth. The second pair of pereopods is much shorter than the total length of the body, merus but carpus is longer than merus. Body were bulky and broader shape. Telson possesses two pairs of dorsal spines and uropod does not have accessory spine (Plate-d). [19], reported regarding this species is found in the Indo-west Pacific: India, Pakistan, Bangladesh. In India, it occur mostly in north-east coast and contributed local fishery. The bulk quantity of prawn available during monsoon at the landing centre. Aquaculture has been attempted but no record for production. It fetches good price in local market. It is often difficult to distinguish Decapodes Crustacean the juveniles of *M. lamarrei* from those of *M. rosenbergii*, *M. malcolmsonni* and *M. gangeticum*, detail presented in the (Table: 1 & Plate-a, b, c &d).

3.5 Abundance of prawn species: The species *Macrobrachium gangeticum*, *M. gandaki*, *M. tiwari*, and *M. lamarrei* occurred at the sampling sites. The *Macrobrachium lamarrei* and *M. tiwari* were found dominant species in year around, whereas in case of *M. gandaki* and *M. gangeticum* were observed restricted during May to October and June to September during the investigation period in the Koshi river near Kursela, Katihar. There was a slight seasonal variations observed in the species availability. The number of individuals were found more and subsequently decreases as *Macrobrachium lamarrei*, *M. Tiwari*, *M. gandaki* and *M. gangeticum* respectively during the investigation period. It may due to water temperature, depth and abundance of food resources present in these site like algae and planktons. Therefore, there is significant correlation between its abundance and food resources as indicated by [19,22,25] documented commercially important availability and distribution of 12 species of freshwater prawn fauna occurring in India. Whereas [26] reported, regarding period wise sex population varied from a maximum in winter whereas minimum in summer season of *Macrobrachium assamense peninsuliric* in the Khohi river Uttarakhand, India. [31] studied taxonomic and identification of freshwater prawn *Macrobrachium assamense peninsuliric* in Garhwal region, this prawn characterized by small rostrum with 5 to 10 dorsal teeth and 0 to 3 ventral. [22], recorded four species of freshwater prawns in the middle stretch of the river Ganga near Patna belonging to genera *Macrobrachium* such as *Macrobrachium gangeticum*, *M. malcolmsonii*, *M. lameraii*, *M. lameraii lameraii*, however, *M. gangeticum* only available in the rainy season. In current survey, low rate of wealth of species was studied in sampling site warning loss of its suitability for nourishing life due to devastation of ecological habitats, increased level of pollution and over exploitation of the species. [8] reported 5 species of freshwater shrimp consisting of 2 families, they also determine diversity and the abiotic conditions of freshwater shrimps habitat in Indonesia and [32] reported relationships between water quality constraints and the development rate of white shrimp (*Litopenaeus vannamei*) in concentrated ponds. [27], reported the impact of climate change on prawn post larvae fishing in coastal Bangladesh. [28] observed similar reasons for declined diversity and low abundance in one of the studied habitats.

4.0. CONCLUSION

This study will serve as starting point evidence regarding freshwater prawn *Macrobrachium* diversity and assortment in the Koshi region of North Bihar. Variety and richness of prawn species is affected by the human caused factors that has considerable impact. The report recommends that site requires proper management and develop hatchery technology for its conservation and to prevent the further loss of populations from this region.

ACKNOWLEDGEMENT

The First Author indebted to Honor'ble Vice Chancellor and Director research, (Bihar Agricultural University, Sabour, Bhagalpur) for the financial assistance under research project code BAU/SNP/NRM/Rabi/2019-23. Mr. Sushant Bhanu, Research Scholler was thankfully acknowledged for assistance during investigation.

REFERENCES

1. Camara, I. A. Konan, M. K. Diaman, D. Edia G. and Gourence E.O. Ecology and diversity of freshwater shrimp in Bonco National Park, Cote dlvoire (Banco River Basin) Knowledge and Management of Aquatic Ecosystem, 2009, 393:05.
2. Snyder, M. N. and Freeman M. C. Purucker S. T. and Pringle CM. Using occupancy modeling and logistic regression to assess the distribution of shrimp species in lowland streams Coata Rica Does regional groundwater create favorable habitat. *Freshwater Science*, 2016. 35 (1):80-90.
3. Grave, C. D. Cai, Y, and Anker, A. Global diversity of Shrimps (Crustacea: Decapoda: Caridea) in freshwater. *Hydrobiologia*, 2008.595: 287-293.
4. Prasad, S., Kanaujia, D. R. and Patra, A. K. Diversity Abundance and Composition of large freshwater prawn species in the Ganga river system. *The Bioscan*, 2012. 7 (4):685-689.
5. Pandey, A. K. Upadhyay, A. K. and Lakra, W. S. Diversity of commercially important freshwater prawns and their aquaculture potential in India. *J. Exp. Zool. India*, 2010. 13 (1):121-128.
6. Prasad, S., Khan, M. A. and Kaushal, D. K. Depletion of the Ganga river prawn *Macrobrachium gangeticum*: Need to conservation. *Proc. Zool. Soc. India*. 2010. 9 (2):85-90.
7. De Grave, S. Smith KG, Adeler, NA, Allen, D. J. Alvarez, F. Anker A. Cai V, Carrizo SF. Kiotz W. Mantelatto FL Page TJ, Shy Jy and Villalobos J L Dead Shrimp Blues: A global assessment of extinction risk in freshwater shrimp (Crustacea Decapoda: Caridea) PLOS. ONE 2015.10:e0120198.
8. Susilo, V E, Suranto and Fadillan, N. Narulia E and Wowor, D. Diversity of freshwater shrimp (Decapoda) from bandealit river Meru Betri National Park East Java, Indonesia. *Journal of Physics: 2020, Conference Series* 1465012009.
9. Salathia, S. and Langer, S. Study of freshwater prawn diversity from different rivers of Jammu, India. *Indian Journal of Ecology*. 2022, 49 (1): 183-186.

10. Jhingran, V. G. Fish and Fisheries of India. (Revised 3rd edition) Hindustan Publishing Corporation (India) Delhi, 2003.750.
11. Prasad, S. and Kanaujia, D. R. An observation on reproductive potential of Ganga river prawn *Macrobrachium gangeticum* (Bate). *Progressive Research- An International Journal*. 2014, 09 (Special III):1028-1032.
12. Ranjane, S. S. and Marippan, N. A general and ecological diversity of freshwater prawns *Macrobrachium canarae* and *Caridina gracilirostins* from Kanyakumari Dist. Tamil Nādu, India. *International J. of Genetic Engineering and Biotechnology*, 2011, 2: 23-32.
13. Athiyaman, R and Rajendran. Diversity of freshwater prawns (Crustacea: Decapoda: Paaemonidae and Atyidae) in the river Cauvery. *Elixir Appl. Zoology*, 74 (2014): 26975-26977.
14. Prakash, S. Studies on affinities of Ganga river prawn *Macrobrachium gangeticum* in proceeding of the fifth Indian fisheries forum (Eds) : S. Ayyappan, J. K. Jena and M. Mohan Joseph. Bhubaneswar, India. 2002. 241-244.
15. Prasad, S. Size group, Sex- Ratio, Male and Female Percentage of Indian River Prawn *Macrobrachium malcolmsonii* (Edwads), *Progressive Research - An International Journal*. 2013, 08 (Special Issue):656-659.
16. Prasad, S. Maturation and fecundity of large freshwater prawn *Macrobrachium gangeticum* (Bate) and *Macrobrachium malcolmsonii* (Edwards). In the Ganga river system in India, *Current Journal of Applied Science and Technology*. 2020, 39(21):148-155.
17. Ahmad, J. Freshwater prawn fisheries resource and opportunities for their development: Book Cinrarens, D, 39; Raipur. 1999.Allied Printers New Delhi. Pages 1-136.
18. Kurian, C. V. & Sebastian, V., Prawn and Prawn Fisheries of India. Hindustan Pub. Corp., Delhi, 2001, pp.310.
19. Kanaujia, D. R. Indian River Prawn, *Macrobrachium malcolmsonii* and minor species of commercial important, International Symposium Souvenir on Freshwater prawn 2003, College of fisheries, Agricultural University, Kochi, India, 2003. 51-56.
20. Prasad, S., Ganga Jhinga - Ek Parichay, Daya Publishing House New Delhi, 2007, ISBN. 81-7035-466-8.
21. Kanaujia, D. R. Biology of Freshwater prawn *Macrobrachium malcolmsonii* (Edwards)of river Ganga, Buxar, Bihar. In Proceeding National Seminar on Freshwater Aquaculture CIFA, Bhubaneswar India. 1989, 51-54.

22. Prasad, S. Studies on the freshwater prawn fishery of river Ganga with special reference to the larval biology of larger *Macrobrachium* Species. Ph.D thesis, Utkal University, Bhubaneswar Orissa, 2005.
23. Kanaujia, D. R. Mohanty, A. N., Mitra, G., Prasad, S. Breeding and seed production of the Ganga river prawn *Macrobrachium gangeticum* (Bate) under captive condition. *Asian Fisheries Science*, 2005, 18 (3); 371-381.
24. Prasad, S. Outdoor hatchery larval biology and seed production of Ganga river prawn *Macrobrachium gangeticum* (Bate), *Curr. J. of Appl. Science &Tech.* 2020. 33 (3); 1-7.
25. Jayachandran, K. V. and Indira, B. Sustainable exploitation of freshwater prawn diversity of India for foods and livelihood security with emphasis on planning. *Indian J. Sci. Res.* 2010, 1(2): 127 – 132.
26. Bahuguna, P. Sex population structure of *Macrobrachium assamense peninsularie* (Tiwari) (Crustacea, Decapoda, Palaemonidae) in Khoh River, Uttarakhand, India. *Int. J. Curr. Microbiol. App. Sci.* 2013, 2 (10): 382-390.
27. Ahmed, N. Ambrogi, AO, Muir, J. F. The impact of climatic change on prawn post larvae fishing in coastal Bangladesh. Socioeconomic and ecological perspective, *Mar. Policy*, 2013, 39:224-233.
28. Jewel MAS, Haque, MA, Khatun, R. and Rahman MS. A comparative study of fish assemblage and diversity indices in two different aquatic habitats in Bangladesh: Lakhandaha Wetland and Atari River. *Jordan Journal of Biological Sciences.* 2018, 11:427-434.
29. Prasad, S., Kanaujia, D. R., and Kausal, D. K. A New species *Macrobrachium gandaki* (Bate, 1868) in middle stretch of Ganga river, India. *Proc. Zool. Soc. India*, 2011, 10 (2) : 55-61.
30. Prasad, S, Comparative larval biology of Ganga river prawn species *Macrobrachium malcolmsonii* (Edwards) and *Macrobrachium gangeticum* (Bate) in Hatchery Condition *Progressive Research - An International Journal.* 2015, 10 (Special Issue-8):4426-4431.
31. Kumar, K., Rana, A.R., and Kotnala, C. B. Taxonomic study and identification of freshwater prawn *Macrobrachium assamense peninsularic* (Tiwarly, 1958) in Garhwal region of Central Himalaya, India, *International Journal of Research and Analytical Review*; 2018. 5 (4): 485-491.
32. Ariadi, H. Fadjar, M. Mahmudi, M. The relationships between water quality parameters and the growth rate of white shrimp (*Litopenaeus vannamei*) in intensive ponds. *Aquaculture, Aquarium, Conservation & Legislation*, 2019. 12 (6): 2103-2116.