

A Generalized Common Fixed Point Result for Occasionally Weakly Compatible Four Self-Mappings in a Cone Metric Space

ABSTRACT

In this paper, we prove a unique common fixed point theorem for occasionally weakly compatible four self – mappings in a cone metric space. Our result generalized and improved the recent results existing in the literature.

Keywords: [Fixed point, cone metric space, and occasionally weakly compatible]

1. INTRODUCTION AND PRELIMINARIES

L.G. Huang and X. Zhang, [9] generalized the concept of a cone metric space, and they replacing the real numbers by ordered Banach space and proved fixed theorems in cone metric space, later on my authors inspired with these results and they have been generalized ,extended and improved the results of [9] using with different types of contractive conditions [see, eg. 2-5, 7-10] . And recently Bhatt and Chandra [6] proved some results on occasionally weakly compatible mappings. Our result is generalize and improving the results of [6].

The following definitions are useful to our main results which are of the existing literature .

Definition 1.1. Let S be a real Banach space and P be a subset of S . The set P is called a cone if and only if:

- (a). P is closed, non–empty and $P \neq \{0\}$;
- (b). $a, b \in \mathbb{R}$, $a, b \geq 0$, $x, y \in P \Rightarrow ax + by \in P$;
- (c). $x \in P$ and $-x \in P \Rightarrow x = 0$.

Definition 1.2. Let P be a cone in a Banach space S . Define partial ordering ‘ \leq ’ with respect to P by $x \leq y$ if and only if $y - x \in P$. We shall write $x < y$ to indicate $x \leq y$ but $x \neq y$ while $x \ll y$ will stand for $y - x \in \text{int } P$, where $\text{int } P$ denotes the interior of the set P . This cone P is called an order cone.

Definition 1,3. Let S be a Banach space and $P \subset S$ be an order cone. The order cone P is called normal if there exists $K > 0$ such that for all $x, y \in S$,
 $0 \leq x \leq y$ implies $\|x\| \leq K \|y\|$.

The least positive number K satisfying the above inequality is called the normal constant of P .

Definition 1.4[7] Let X be a nonempty set of S . Suppose that the map $\rho : X \times X \rightarrow S$ satisfies:

- (d1). $0 \leq \rho(x, y)$ for all $x, y \in X$ and $\rho(x, y) = 0$ if and only if $x = y$;
- (d2). $\rho(x, y) = \rho(y, x)$ for all $x, y \in X$;
- (d3). $\rho(x, y) \leq \rho(x, z) + \rho(z, y)$ for all $x, y, z \in X$.

Then ρ is called a cone metric on X and (X, ρ) is called a cone metric space.

It is clear that the concept of a cone metric space in case of the normal case is more general than that of a metric space.

Example 1.5 Let $S = \mathbb{R}^2$, $P = \{(x, y) \in S \text{ such that } : x, y \geq 0\} \subseteq \mathbb{R}^2$, $X = \mathbb{R}$ and $\rho : X \times X \rightarrow S$ such that $\rho(x, y) = (|x - y|, \alpha|x - y|)$, where $\alpha \geq 0$ is a constant. Then (X, ρ) is a cone metric space.

Definition 1.6 Let (X, ρ) be a cone metric space. We say that $\{x_n\}$ is

(i) a convergent sequence if for any $c \gg 0$, there is a natural number N such that for all $n > N$, $\rho(x_n, x) \ll c$, for some fixed x in X . We denote this $x_n \rightarrow x$ (as $n \rightarrow \infty$).

(ii) a Cauchy sequence if for every c in S with $c \gg 0$, there is a natural number N such that for all $n, m > N$, $\rho(x_n, x_m) \ll c$

A cone metric space (X, ρ) is said to be complete if every Cauchy sequence is convergent.

Definition 1.7. Let F and G be self-mappings of a set X . If $u = Fx = Gx$ for some x in X , then x is called a coincidence point of F and G , and u is called a point of coincidence of F and G .

Proposition 1.8 . Let F and G be occasionally weakly compatible (owc) self-mappings of a set X iff there is a point x in X which is coincidence point of F and G at which F and G are commute.

Lemma 1.9 . Let X be a set, F, G are owc self-mappings of X . If F and G have a unique point of coincidence $u = Fx = Gx$, then u is the unique common fixed point of F and g .

2. Main Results

In this section we have generalized the results of [6].

Theorem. Let (X, ρ) be a cone metric space and P be a normal cone. Suppose that f, g, A and B are occasionally weakly compatible self mappings of X and satisfy the following conditions

(i) $\rho(fx, gy) \leq p\rho(Ax, By) + q\rho(fx, Ax) + r\rho(gy, By) + t[\rho(fx, By) + \rho(gy, Ax)]$,

for all $x, y \in X$, for which $fx \neq gy$, $p, q, r, t \in [0, 1]$ and satisfying $p + q + r + 2t < 1$.

(ii) The pairs $\{f, A\}$ and $\{g, B\}$ are occasionally weakly compatible.

Then f, g, A and B have a unique common fixed point.

Proof. Given $\{f, A\}$ and $\{g, B\}$ are occasionally weakly compatible, then there exists $x, y \in X$ such that $fx = Ax$ and $gy = By$. We claim that $fx = gy$. Otherwise by (i)

$\rho(fx, gy) \leq p\rho(Ax, By) + q\rho(fx, Ax) + r\rho(gy, By) + t[\rho(fx, By) + \rho(gy, Ax)]$.

Since $fx = Ax = w$ and $gy = By = z$ are points of coincidence of $\{f, A\}$ and $\{g, B\}$ respectively.

$$\Rightarrow \rho (fx, gy) \leq p \rho (fx, gy) + q \rho (fx, fx) + r \rho (gy, gy) + t[\rho (fx, gy) + \rho (gy, fx)],$$

$$\leq (p + 2t) \rho (fx, gy), \text{ since } p + 2t < 1 ,$$

$\Rightarrow \rho (fx, gy) < \rho (fx, gy)$, which is a contradiction.

Therefore, $fx = gy$, that is, $fx = Ax = gy = By$.

Moreover, if there is another point z such that $fz = Az$, then by (i) $fz = Az = gy = By$ or $fx = fz$.

Hence, $w = fx = gx$ is a unique point of coincidence of f and A . Then by the Lemma (1.9)

w is the unique common fixed point of f and A (1).

Similarly, there is a unique point $z \in X$ such that $z = gz = Bz$. We suppose that $w \neq z$, then by (i), we get that

$$\rho (w, z) = \rho (fw, gz) \leq p \rho (Aw, Bz) + q \rho (fw, Aw) + r \rho (gz, Bz) + t[d(fw, Bz)$$

$$+ \rho (gz, Aw)],$$

$$\leq p \rho (w, z) + q \rho (w, w) + r \rho (z, z) + t[\rho (w, z) + \rho (z, w)],$$

$$\leq (p + 2t) \rho (w, z), \text{ since } p + 2t < 1$$

$\Rightarrow \rho (w, z) < \rho (w, z)$, which is a contradiction.

Therefore, w is the unique common fixed point of f, g, A and B .

3. CONCLUSION: Our result is more general than the results of [6] and it is also improvement of the results of [6].

REFERENCES

- [1] M. Abbas and G. Jungck, Common fixed point results for non commuting mappings without continuity in cone metric spaces, J. Math. Anal. Appl. Vol. 341, issue 1, 2008, 416-420.
- [2] M. Abbas, B. E. Rhoades, Fixed and periodic point results in cone metric spaces, Appl. Math. Lett. 21,2008, 511-515.
- [3] I. Altun, B. Damjanovic, D. Djoric, Fixed point and common fixed point theorems on ordered cone metric spaces, Appl. Math. Lett. 2009 ,doi:10.1016/j.aml.2009.09.016 .
- [4] I. Altun, B. Durmaz, Some fixed point theorems on ordered cone metric spaces, Rend. Circ. Mat. Palermo 58, 2009 319-325.
- [5] M.A.Al-Thagafi and Naseer Shahzad, Generalized I-nonexpansive selfmaps and invariant approximations, Acta Mathematica Sinica, 24, 2008, 867-876.
- [6] Arvind Bhatt and Harish Chandra, Occasionally weakly compatible mappings in cone metric space , Applied Mathematical Sciences, Vol. 6, 2012, no. 55, 2711 – 2717.
- [7] Guangxing Song, Xiaoyan Sun, Yian Zhao, Guotao Wang, New common fixed point theorems for maps on cone metric spaces, Appl. Math. Lett. 32, 2010 1033-1037.
- [8]G. Jungck and B.E.Rhoades, Fixed point theorems for occasionally weakly compatible Mappings, Fixed point theory, 7, 2006 , 286-296.
- [9] L.G. Huang, X. Zhang, Cone metric spaces and fixed point theorems of contractive mappings J. Math. Anal. Appl. 332, No.2, 2007, 1468-1476.
- [10] K. Prudhvi, A Study on Fixed Point Results for OWC Self-Maps on C-Metric Space, Asian Research Journal of Mathematics, 18(9), 2022, 54-57, ARJOM.88770.

UNDER PEER REVIEW