

LEAKY GUT SYNDROME AND ORAL LESIONS - A HYPOTHESIS

ABSTRACT-

There are many oral lesions which run a chronic course and always recur despite being treated multiple times by experts. They cause tremendous damage to multiple tissues. Many oral diseases have orofacial manifestations only, while others are associated with multiple serious systemic manifestations. Various researchers across the globe have been struggling to find out permanent solutions through various therapeutic approaches like, ayurveda and holistic etc but internal root causes should be found out which might give us breakthrough quick therapies in near future.

We need to explore evidence based various hidden internal /systemic causes to achieve permanent cure of such chronic long-standing diseases.

All systems of the body are interconnected and always works in harmony. Oral diseases can never be restricted to orofacial tissues. They are strongly connected to gastro intestinal, ectodermal tissues and all the systems. This article is an attempt to find out same through few hypotheses and connecting oral lesions with other tissues and their disorders .

Keywords-oral diseases, hypothesis, leaky gut syndrome, systemic cause, interstitial fluid, acidic

INTRODUCTION

Dental literature is flooded with the description of oral lesions that have possible cause and effect relationship. When there is single etiological factor associated with the disease ,it is very easy to apply therapeutic measures to treat that disease effectively.(1)However there are oral diseases which are multifactorial in origin which pose challenges to clinicians to plan therapeutic measures .(2) There are wide range of diseases like dental caries , oral potentially malignant disorders , oral cancer etc where there is no single causative etiologic agent (3).Therefore these diseases have been extremely challenging to treat for healthcare providers. Researchers have been trying very hard to narrow down their search in finding out exact cause but with limited success.As we explore and study pathological alterations in these lesions we came across some initial alterations in deeper tissues of these lesions which made us to ask question “**Are there systemic factors responsible for triggering most initial changes in these oral lesions which are then aggravated by external stimuli?**”(4)

DISCUSSION-

We need to find out the exact root cause of oral lesions. There are few hypothesis suggested by authors as follows-

1. **Dental caries** - Dental caries is the most common microbial disease affecting teeth. Various theories have been advanced to explain etiopathogenesis of dental caries. (1) All of them have their own drawbacks as none of them explains as to why there are caries free individuals in spite of consuming sugars in their diet. (1) Most accepted theory is the Chemico-parasitic theory put forth by W.D. Miller which says dental caries is caused by bacteria which ferment sugars present on the tooth surface. Fermentation process leads to formation of acids and results in dissolution of enamel crystals leading to cavity formation. Therefore sandhya et al and larma et al postulated that there is additional internal systemic approach before external caries attack. (5)(6)(7)(8) This internal attack is due to acidic interstitial fluid which flows through dental lymph or dentinal tubules which causes internal tooth porosities due to removal of minerals in an attempt to buffer acidic interstitial fluid. (6)(9)(10)(11)(12) Such porosities will decide whether such individuals are prone to caries or not. (10) Sometimes even if porosities are more but if oral hygiene methods are good then patients would be less prone to caries attack.
2.) **Erosion**-Erosion is considered to be wearing of tooth structure due to chemical factors . Such chemical factors are mostly cold drinks .Now the question arises whether such factors act through external or internal routes. It could also be hypothesized that once the internal porosities are formed such teeth are prone to external attack. If the attack is bacterial then caries will occur. If oral hygiene methods are good then caries occurrence will be less ,but chances of erosion will be more . Patients with erosion will have good oral hygiene methods and personal hygiene too. Inspection of personal hygiene can give a clue to correlate erosion .(13)(14)
3. **Oral premalignancy** -Oral premalignancy such as OSMF ,lichen planus and leukoplakia have mostly due to external carcinogens such as tobacco ,alcohol and areca nut ,medicaments ,filling materials etc. but when histopathological descriptions are put **together** in sequence then initiating changes are seen in basal portion of the epithelium or the **the** stroma .(15)

In **leukoplakia** the cause is tobacco eating it is always in contact of surface mucosa but resultant epithelial dysplasia always starts from basal layer instead of stratum corneum .hyperkeratosis is just a protective mechanism against tobacco but cellular alterations starts from basal region .(16)

In **oral submucous fibrosis** ,the continuous trauma of areca nut chewing is on the surface oral mucosa but the histopathological alterations starts at stroma and it further leads to epithelial dysplasia which again starts at the basal region.(17)

Lichen planus is a mucocutaneous premalignant lesion which is caused by stress, medication ,and filling materials . Histopathological events start at stomal level in the form of juxta epithelial dense inflammatory cell reaction ,liquefaction degeneration of basal cell layer and saw tooth rete pegs civatte bodies etc.(18)(16)

4. **Genetic disorders** -Genetic disorders would be due to a damage to genes through mother and transmitted to fetus.(1)
5. **Joint disorders**- Such escaped molecules can easily lodge in joint spaces too and create inflammation and joint disorders .(19)(4)
6. **Infections**- our body is surrounded by many microbes on the body surfaces and in the environment. Healthy and live body is a bit immune to the infections but the unhealthy and dead body is prone to microbial attack. (4)(20)
7. **Intelligence**- when intracellular interstitial fluid accumulates it leads to neuron functions too and leads to confusion and sometimes disorientation and imbalance.(21)(4)
8. **Weight gain** -sometimes there is sudden weight gain in the absence of heavy eating habits , instead it occurs due to less eating habits, **i** such cases there is accumulation of adipose tissues around the bones and abdomen to prevent acid to withdraw minerals and damage the tissues in an attempt of buffering the acidity.(22)(23)

Question arises what could be the reason for acidic interstitial acidic ph . Literature says, emotional stress , bad eating habits, insufficient sleep, excessive exercise ,vitamin D ,iron and calcium deficiency etc. Such nutritional deficiency can lead to weak muscle contractions which lead to slow peristaltic movements and

slow digestion .Food retains for longer in the body before excretion leads to acid accumulation and micro trauma to gut wall .(24)(25)

9. **Leaky gut syndrome and skin diseases** - It has been mentioned that if our gut is healthy we are healthy. It means if there are injuries to gut tissues due to food in terms of unhealthy ingredients and improper mastication, it leads to overload on soft tissues of the gut wall. This leads to micro trauma on the luminal wall of the gut.(25)

Our gut tissues are under constant trauma and hardly get time for self-healing. They are subjected to continuous food intake and continuous digestion. A traumatized gut will be subjected to more trauma.(26)

Such micro trauma or micro perforation is called a leaky gut. Such leaky gut when subjected to food intake peristalsis starts. But some undigested food molecules enter the circulation through these perforations present in the gut wall . **These escapes undigested food molecules** and perforations could be of various sizes depending on the extent of trauma and injuries.(25)

It could be hypothesized that onces these molecules escapes the gut wall, depending on the size of food molecules it get lodged **within tissues of peripheral tissues**. Bigger molecules will get lodged in deeper tissues and organs while smaller molecules reach up to most peripheral parts of body such as extremities and oral cavity specifically epithelium and connective tissues.

Autoimmune blistering diseases-This could be the base for various **autoimmune blistering diseases** in the oral cavity and skin with and without systemic involvement. (25)

For example **pemphigus** has intraepithelial vesicle formation ,**pemphigoid** has sub epithelial vesicle formation and other diseases have split at anchoring fibril level etc. (2)

It can be hypothesized that blistering diseases with deeper tissues esicles have larger food molecules as self-antigen and diseases with intraepithelial and sub epithelial split have relatively smaller escaped food molecules. Since they have escaped digestion they evoke autoimmune response and leads to tissue destruction in the form of vesicle formation .

Conclusion -Various researchers involved in intensive care unit and ayurveda primarily on focus on buffering this interstitial fluid and healing the traumatic gut as therapeutic measures along with supportive and symptomatic treatment.

References –

1. Shafer, Hine L. Shafer's Textbook of Oral Pathology. 5th ed. In: Rajendran R, Sivapathasundharam B E, editor. New Delhi: Elsevier Ltd; 2006. 404–7 p.
2. Sapp JP, Eversole LR WG. . Contemporary oral and maxillofacial pathology. . 2 nd ed. St. Louis: Mosby; .; 2004.
3. Regezi JA, Sciubba J, Jordan RCK, Joseph A. Regezi, James J. Sciubba RCKJ. Oral Pathology: Clinical Pathologic Correlations Seventh Edition [Internet]. Vol. 01, International Standard Book. 2017. 1689–1699 p. Available from: www.elsevier.com/permissions.
4. Sandhya T. INTERNAL PH IN HEALTH AND DISEASE.
5. Larmas M. Dental caries seen from the pulpal side: A non-traditional approach. *J Dent Res.* 2003;82(4):253–6.
6. Southward K. The systemic theory of dental caries. *Gen Dent.* 2011;59(5):367–73.
7. Gronthos S, Mankani M, Brahim J, Robey PG, Shi S. Postnatal human dental pulp stem cells (DPSCs) in vitro and in vivo. *Proc Natl Acad Sci U S A.* 2000;97(25):13625–30.
8. Sandhya Tamgadge AT. Is Systemic (Interstitial Fluid) Acidosis an Initial Event in the Etiopathogenesis of Dental Caries? A Hypothesis. *Dent Hypotheses.* 2018;9(4):96-100.
9. Özok AR, Wu MK, Ten Cate JM, Wesselink PR. Effect of dentinal fluid composition on dentin demineralization in vitro. *J Dent Res.* 2004;83(11):849–53.
10. Steinman RR, Leonora J. Relationship of Fluid Transport Through the Dentin to the Incidence of Dental Caries. *J Dent Res.* 1971;50(6):1536–43.
11. Neves-Silva R, Alves FA, Antunes A, Goes MF, Giannini M, Tenório MD, et al. Decreased dentin tubules density and reduced thickness of peritubular dentin in hyperbilirubinemia-related green teeth. *J Clin Exp Dent.* 2017;9(5):e622–8.
12. Agematsu H, Abe S, Shiozaki K, Usami A, Ogata S, Suzuki K, et al. Relationship between large tubules and dentin caries in human deciduous tooth. *Bull Tokyo Dent Coll.* 2005;46(1–2):7–15.
13. Joshi M. Techniques to Evaluate Dental Erosion: A Systematic Review of Literature. *J Clin Diagnostic Res.* 2016;1–7.
14. Enam F, Mursalat M, Guha U, Aich N, Anik MI, Nisha NS, et al. Dental erosion potential of beverages and

- bottled drinking water in Bangladesh. *Int J Food Prop* [Internet]. 2017;20(11):2499–510. Available from: <https://doi.org/10.1080/10942912.2016.1242607>
15. Tamgadge S, Tamgadge A. Histopathology of oral submucous fibrosis in third dimension with an additional note on hypothesis of epithelial atrophy. *J Microsc Ultrastruct*. 2020;8(1).
 16. Cawsons RA OE. *Cowson's essentials of oral pathology and medicine*. 8 th ed. Philadelphia: Churchill livingstone,Elsevier; 108–9 p.
 17. Jk S. Review Article Oral Submucous Fibrosis - A review [Part 2]. 2011;2(1):37–48.
 18. Warnakulasuriya S, Johnson NW, Van Der Waal I. Nomenclature and classification of potentially malignant disorders of the oral mucosa. *J Oral Pathol Med*. 2007;36(10):575–80.
 19. Tamgadge S, Tamgadge A. Histology of tooth development in 3D animation video and images - A preliminary report. *J Microsc Ultrastruct*. 2020;0(0):0.
 20. Takahashi N. Microbial ecosystem in the oral cavity: Metabolic diversity in an ecological niche and its relationship with oral diseases. *Int Congr Ser*. 2005;1284:103–12.
 21. Schwalfenberg GK. The alkaline diet: Is there evidence that an alkaline pH diet benefits health? *J Environ Public Health*. 2012;2012.
 22. Rajendran R. No Title. In: In:Rajendran R SB, editor. *Benign and malignant tumors of oral cavity*. 6th ed. Philadelphia: Elsevier Ltd.; 2009.
 23. Kellum JA. Determinants of blood pH in health and disease. *Crit Care*. 2000;4(1):6–14.
 24. Tamgadge S. Emotional Stress and Acidic Intercellular PH is an Initial Event in the Etiopathogenesis of Oral Lesions - A Hypothesis. 2019;3(1):78–9.
 25. Fasano A. Leaky gut and autoimmune diseases. *Clin Rev Allergy Immunol*. 2012;42(1):71–8.
 26. Dietrich CG, Geier A, Oude Elferink RPJ. ABC of oral bioavailability: Transporters as gatekeepers in the gut. *Gut*. 2003;52(12):1788–95.