

## Review Article

### Breeding approaches for quality improvement in fruit crops: strategies and achievements

#### Abstract

As we know that fruit plays important role in the daily human diet for healthy living and is also a commercial commodity in trade and processing industries. The primary factor that customers use to determine whether or not a fruit is acceptable is its quality like appearance, shape, size, colour and taste, etc. Success in a breeding programme depends upon the overall acceptability of fruit quality because most of the developed varieties having desired traits like resistance to biotic and abiotic stresses could not be commercialized and are not in commercial cultivation owing to their poor-quality traits. Therefore, the development of cultivars with desirable quality attributes in fruit crops is challenging. Quality improvement in fruit crops is restricted by several factors such as long juvenility, tall stature, environmental stress and high heterozygosity. Quality traits in fruit crops are polygenic and governed by many genes which makes it difficult to improve particular desirable traits. Many attempts have been made to enhance the qualitative characteristics of annual crops, although perennial fruit crops neatly overlook this issue. Accordingly, the use of both combined conventional and modern breeding techniques could in overcoming these problems. Biotechnological and molecular approaches like marker-assisted selection, transgenics, genomic editing, genomics cis-genics and candidate gene offer precision and reliability to reduce the breeding cycle and are also advantageous when dealing with tedious fruit crops. The challenges with fruit breeding and the state of various breeding techniques for enhancing fruit quality in fruit trees will be the main topics of this review.

Keywords: MAS, Fruit quality, Resistance, Genomics, Shelf life

#### Introduction

Fruit breeding programs have a wide range of specific aims regarding abiotic and biotic stresses resistance, tree architecture, precocity, and productivity, with a common objective to develop high-quality fruit (reference). Fruits quality has a different meaning for different fruit species consisting of diverse attributes (Callahan, M. A. 2003). For instance, in some species, the crisp texture is much more acceptable than the soft vice versa. Few fruits require a balance of acidity and sweetness for taste whereas, for others, it is simply defined by the degree of sweetness (reference).

However, all commercial cultivar releases must have delicious palatable fruits (reference). It does not matter whether the tree is a disease or insect resistant or highly productive but if the quality of the fruit is not acceptable, it would be failed in the commercial market (reference).

It is difficult to define the accurate definition of fruit quality as it varies according to the taste or requirement of the consumer, so the simple definition of fruit quality is: Whatever the consumer desires (Barritt, 2001; Elia, 2001; Kupferman, 2002). Since the nature of people is different, their desires and ideas of quality are different hence breeders need to provide numerous alternative forms to meet the market desire. Quality may also refer to aspects like colour, size, nutrients, shelf life, suitability for processing,

texture, taste, and sweetness (Kader, 2002). However, breeding for quality improvement in fruit crops as they are perennial in nature is hampered by several limitations including the large size of the plant, long juvenile phase, and environmental problems (e.g., fruit drops due to natural calamities). Besides, fruit quality is a polygenic trait, which is quantitatively inherited and thus making the breeding program complicated in the quality improvement of fruit crops. Any property of an individual showing heritable variation is known as a character or trait. A trait that defines some aspect of produce quality is quality traits. Each crop has a specific and often somewhat completely different set of quality traits. Fruits play a significant role in the nutritious diet (Wargovich, 2000) as it has the potential to provide all the essential elements or compounds which are generally deficient in agronomical crops such as fibre, vitamins, minerals, proteins, fats, and carbohydrates which are perfect for curing nutritional disorders.

#### **Bottlenecks in fruit trees breeding and quality improvements**

To date, continuous attempts have been made to improve the quality traits of staple crops whereas very fewer efforts have been made to the improvement of fruit trees as compared to agronomical crops. In the case of fruits, breeding objectives are mainly focused on tree architecture, breeder to be least interested. For example, an apple-breeding program that was undertaken by Dresden-Pillnitz, Germany screened 52,000 (Myles, 2013). Knowledge of genetics is very meagre in the case of tropical fruits as compared to temperate fruits (Arias et al., 2012). Breeding fruit trees are even made complicated by reproductive biology (diurnal flowering in avocado), polyembryony (zygotic and nucellar seedlings in mango), sexual types (dioeciousness in papaya) and apomixes (obligate apomixes in garcinia) (Arias et al., 2012). In addition to this, environmental problems and natural calamities often result in huge commercial and breeding losses due to flower drops, fruit drops, pest and disease infestations etc.

precociousness, yield, resistance against biotic, abiotic stresses, and physiological disorders (Ray, 2002); however, very little attention has been paid to maintaining the quality of fruit. There are many constraints in the breeding of fruit crops such as consumers usually prefer the local or indigenous varieties as compared to improved varieties thereby making the breeding program a failure. For example, in the case of mango, many improved selections and hybrids have been developed, which are regular and heavy bearers and are also free from the problem of spongy tissue, however, local farmers tend to prefer Alphonso mango, though it is the shy and alternate bearer, and suffers from the spongy tissue.

The major constraint in breeding is the large size and perennial behaviour of the fruit trees which makes the breeding program difficult especially while carrying out important operations such as emasculation, bagging, tagging, data recording, harvesting, etc. The majority of fruit trees are propagated through the asexual means of reproduction, which bears flowers and fruit normally after 5 years of age (Arias et al., 2012), whereas those derived from seedlings take more than 10 years, thereby prolonging the breeding cycle. Breeders have to wait for a long time to get the result. In addition, after the long wait, still, the chance of getting undesired results is there and which forced the seedlings for 26 years and eventually, only three varieties were released at the commercial level

Fruit quality improvement needs strong genetic knowledge about its inheritance and variation. Quality characters are inherited quantitatively and regulated by multiple genes. Fruit trees are heterozygous **erops crops**, and a mass population is required for screening to identify the promising genotypes for the breeding of quality fruits (Kenis et al., 2008). The biotechnological approaches provide a precise, reliable, and easy way for addressing some of the problems encountered during conventional breeding. Molecular approaches like marker-assisted selection, candidate genes, genomics (Murovec and Bohanec, 2012), transgenic, and cis-

genics have shown to be advantageous in terms of time, effort and patience required while dealing with the cumbersome crops. There is a negative correlation between yield and quality of fruits.

#### Source of fruit quality traits

1. A cultivated variety: most preferred source.
2. A germplasm lines.
3. A spontaneous or induced mutant.
4. A wild relative.
5. A transgene.

#### Fruit quality traits

##### Appearance

The characteristics that affect appearance are primarily size and colour. During consumer surveys on peaches and apples, it was found that the bigger size is more demanding with bright and clear colour and consumers are willing to pay enough more to have it (Kupferman, 2002).

##### 1. Size and shape

Fruit size has a large genetic component thus selecting for larger fruit is relatively straightforward. Fruit size is a function of cell number, cell volume, and cell density (Janick and Moore, 1996).

##### 2. Colour

Table 1: Pigments responsible for fruit colouration

Colour	Pigments	Example
Orange	Beta-carotene	Mango, Pineapple
Red-purple	Anthocyanins	Grape, Pomegranate, raspberry
Orange	Caricaxanthin	Papaya
Red	Lycopene	Papaya, Guava var. Arka Kiran
Orange-yellow	Flavonoids	Peach, Papaya, Orange, Tangerine
Yellow-green	Lutein & Zeaxanthin	Avocado
Green	Chlorophyll	Guava
Yellow	Xanthophylls	Guava

Source: Singh, J. 2002; Ray, P.K. 2002

##### Taste

When quality is enhanced by breeding by applying breeding methods, it affects the total yield of crop plants for instance if quality is increases yield decreases and vice-versa.

This trait is an important aspect of appearance. The overall colour of fruit is reflected by the colour of the outer pericarp and the flesh colour. Pigments responsible for the colours are various modifications of anthocyanins, lycopene, and carotenoids. Predicting colours is difficult because small modifications or combinations of pigments result in unpredictable colours. Due to that fruit depicts different shades of colour. The de-greening process during ripening exposes the colours in both the pericarp and the flesh (Winkel-Shirley, 2001). The de-greening process is the breakdown of chlorophylls, which is usually done by ethylene. The other pigments are no longer masked by chlorophyll and the fruit 'colours. One of the potential problems of some modern cultivars is that brightly coloured blush in the pericarp has been selected that appears before ripening (Janick and Moore, 1996). This in itself masks the chlorophylls thereby negating the de-greening as an indication of ripeness. For exampleexample, the orange colour of mango is due to Beta-carotene and the red and purple colour of Grapes, Pomegranate, Blackberries, and Blueberries is due to Anthocyanin.

The most important aspect of fruit quality is taste. The fruit may be the most desirable looking, but if it doesn't taste good the consumer will not buy it again. Consumer preference is for higher sweetness, more intense flavors, and firm fruit that softens before consumption (Kader, 2002).

### 1. Sweetness

Major fruit's TSS ranges from 9–20° Brix (Refractometry measure of soluble solids) when ripe. Brix is highly correlated with the amount of sugar contained in the juice. The levels of sucrose, fructose and glucose are what determine sweetness; however, the level of acidity affects the perception of sweetness such that fruit with high sugar and moderate levels of acid will be perceived to be as sweet as fruit with moderate levels of sugar and low acid (Janick and Moore, 1996). The acid levels are primarily based on the concentration of malic or citric acid. Generally, the acid present in fruits is malic acid, tartaric acid, etc. For example, new cultivar development in peach has concentrated on high sugar with low acid to fill a niche in the Asian market (Baldwin, 2002). The fact that sugar accumulation occurs

before final ripening makes it easier to harvest at a time with high sugar.

### 2. Flavour and aroma

Flavour and aroma are determined by a combination of volatiles. There are three main pathways for volatile production; cleavage of lipids followed by alcohol dehydrogenase activity to yield short-chain aldehydes and alcohols, the shikimic acid pathway, and the degradation of terpenoids. Interestingly the colour pigments are also derived from these pathways, anthocyanins from the shikimic acid pathway and  $\beta$ -carotene and lycopene from the degradation of terpenoids (Baldwin, 2002). As fruit ripens there are hundreds of volatiles detected but only some above threshold levels that taste panels can detect. Of those, a few have been shown to determine the characteristic aroma/flavour of particular fruits. For example, p-hydroxyphenylbutan (raspberry), cinnamate derivatives (strawberry), cyanidin-3-rutinoside (litchi), decadienoate esters (pear),  $\gamma$ -decalactone, and linalool (peach), Benzaldehyde (Almond), 2-methyl butyrate (Apple) citral (orange), Isopentyl acetate (Banana), etc (Kumar and Ellis, 2001).

Table 2: Aroma compounds in fruits

Fruits	Compound
Apple-Ripe	Ethyl-2-methylbutyrate
Apple-Green	Hexanal, 2-Hexenal
Banana-Green	2-Hexenal
Banana-Ripe	Eugenol
Banana-overripe	Isopentanol
Grapefruit	Nootakaton
Grape	Methyl Anthranilate
Lemon	Citral
Orange	Valencene
Raspberry	1-(p-hydroxyphenyl)-3-butanone

Source: Singh, J. 2002; Ray, P.K. 2002

### 3. Texture

The texture of the fruit flesh is based on how cells shear in the chewing process, the mouth feels. Texture ranges from crisp to melting and all the stages in between. In melting texture, swelling and softening of the cell wall are evident, but in crisp texture, cell wall swell is not observed

during ripening. Three enzymes, polygalacturonase (PG), the  $\beta$ -subunit of PG, and pectin methyl esterase (PME) have been associated with texture determination. Their substrate is the homogalacturonans or pectin located primarily in the middle lamella of cell walls (Redgwell and Fischer, 2002).

## Keeping quality

Fruit can be harvested at various times concerning their peak quality and that time is dependent on the desired texture, the handling process, and the shelf-life of each commodity. Some fruit (non-climacteric such as blueberry, Grape, and strawberry) are harvested eaten ripe and then stored. Climacteric fruit such as Mango, peach, or apricot is harvested at earlier stages for the fruit to withstand the handling. This fruit will finish ripening during storage and transport. The rate at which the fruit ripens and softens determines when it must be harvested to withstand handling and arrive to the consumer either in the process of ripening or eating ripe. These aspects can be modified postharvest but there is also a large genetic component that can be taken into account in a breeding program.

### 1. Softening

a. Softening is attributed to the disruption of the cellulose/xyloglucan cell. Numerous enzymes have been postulated to be involved including  $\beta$ -galactosidase, expansin (EXP), pectate lyase (PEL), endo-(2-4)  $\beta$ -D-glucanase (EGase), and xyloglucan endotransglycosylase (XET) (Brummell and Harpster, 2001). For example: pectate lyase in strawberries resulted in significantly firmer fruit implying they do have a role in softening (Smith et al., 2002). Polyuronides are depolymerized to a very small size during ripening in avocado. Matrix glycans become highly depolymerized in strawberries but not in avocado.

a. —

b. ~~pectate lyase in strawberries resulted in significantly firmer fruit implying they do~~

~~have a role in softening (Smith et al., 2002).~~

- e. ~~Polyuronides are depolymerized to a very small size during ripening in avocado.~~
- d. ~~Matrix glycans become highly depolymerized in strawberries but not in avocado.~~

### 2. Control of ripening.

1) The expression of quality traits normally is co-ordinately regulated and peaks at ripening. Breeders have been selected for early expression of some of these traits such as skin blush, but the texture, softening flavour development, reduction of acid and phenolic compounds, and colour development peak at the ripe stage (Seymour and Manning, 2002). Sugar accumulation takes place prior to the ripest stage. The problem with harvesting fruit at the peak of quality and ripeness is that the fruit at that stage has practically no shelf life. The fruit needs to be picked before peak ripe, at a stage that combines the maximum development of desirable traits and the maximum shelf life. For example: in climacteric fruit, the increase in the amount of ethylene synthesized triggers final ripening. While, in non-climacteric fruit does not increase ethylene with ripening (Seymour and Manning, 2002).

1) ~~In climacteric fruit, the increase in the amount of ethylene synthesized triggers final ripening.~~

2) ~~Non-climacteric fruit does not increase ethylene with ripening (Seymour and Manning, 2002).~~

Many of the genes involved in those ripening traits are under the control of ethylene. It is unclear whether or not low levels of ethylene in non-climacteric fruit are enough to induce those ripening-related genes or if there are other mechanisms to control ripening. Such as the

Formatted: Indent: Left: 0.5", No bullets or numbering

Formatted: Indent: Left: 0.5", No bullets or numbering

discovery of a MADS-box transcription factor as the gene responsible for early ripening in non-climacteric strawberries (Vrebalov et al., 2002).

### Nutritional Quality traits

Fruits are good sources of fiber, minerals, vitamins, and some beneficial phytochemicals such as carotenoids, phenolics, and glucosinolates. Fruits are a major source of both “macro” nutrients such as fiber and carbohydrates, and “micro” nutrients such as Vitamin C, B complex (thiamin, riboflavin, B6, niacin, folate), A, E, minerals, and the lesser-studied polyphenolics, carotenoids, and

glucosinolates. Nutrients may be classified as either water or lipid soluble—meaning they dissolve in water or a lipid medium (Wargovich, 2000). Water soluble nutrients include Vitamin C, B complex, polyphenolics, and glucosinolates. Fat-soluble nutrients include Vitamin A, D, E, K and other carotenoids such as lycopene and  $\beta$ -carotene. Vitamin C is one of the most sensitive vitamins, being degraded relatively quickly by exposure to heat, light, and oxygen (Vitamin C is unstable) (Oguntibeju et al., 2013). For this reason, it is often used as an index of nutrients Department of Health and Human Services and the degradation. For example:

Table 3: Nutritive value of fruits

Nutrition	Fruits/100gm
Vitamin A ( $\beta$ -carotene)	Mango (4800IU) > Papaya (2020IU)
Vitamin B <sub>1</sub>	Cashew nut (630mg) > Walnut (450mg)
Vitamin B <sub>2</sub>	Bael (1191mg) > Papaya (250mg)
Vitamin C	Barbados cherry (1000-4000mg) > Aonla (600mg) > Guava (199mg)
Carbohydrate	Raisins (77.3%) > Dry Apricot (72.2%)
Protein	Cashew nut (21.2%) > Almond (20.8%)
Fat	Pecan nut (70.4%) > Walnut (64.5%)
Fibre	Fig > Guava (6.9%)
Calcium	Litchi (0.21%)
Phosphorus	Almond (0.49%)
Potassium	Banana
Iron	Dry Karonda (39.1%)
Calorific value	Walnut (687mg) > Almond (655mg) > cashew (596mg)

Source: Singh, J. 2002; Ray, P.K. 2002

### Breeding methods for fruit quality

Breeding for fruit quality can require extended periods, particularly for tree fruits since fruit evaluation cannot be done until the tree is mature and fruiting and the progeny will be in the field for several years before the first evaluations can be done. Secondly, a balance must be achieved

to produce beautiful fruit that has the desirable taste and adequate shelf life to get that fruit to the consumer still beautiful with desirable taste. This task requires the combination of multiple complex traits and precise evaluation. General steps involved in the breeding of fruit crops are depicted in fig.1.




Fig.1 General steps of fruit breeding program.

### Domestication:

Domestication is the process of bringing wild species under human cultivation. The process of domestication started when man started superior plants for use (Simmonds, 1979). Domestication of plants is the change of ideotype to adapt them better to manmade environments (Harlan 1971). It is the first step in the development of cultivated plants. Most of the crops were domesticated by prehistoric man under domestication and the crop species have changed considerably. Examples, Date palm, Olive, Grape, Almond, Fig, and Pomegranate.

### Plant Introduction:

Plant introduction consists of taking a genotype or a group of genotypes of plants into new environments where they were not being grown before. The introduction may involve new varieties of a crop already grown in the area, wild relatives of the crop species or a totally new crop species. Mostly materials are introduced from other countries or continents (Singh et al. 2005). But the movement of crop varieties from one environment into another within a country is also an introduction.

Table 4: Important introductions in different fruit crops

Fruit	Cultivar	Country
<b>Mango</b>	Tommy Atkins, Sensation, Haden- Coloured varieties	USA, Florida
	Sweet	Thailand
	Carabao- Regular bearer	Philippines
<b>Banana</b>	Lady finger (Resistant to Bunchy top)	Australia
	Grand Naine	France
<b>Citrus</b>	Torocco	USA
	Sunramon	Peru
	Kinnow	USA
<b>Grape</b>	Grapefruit	California and Florida
	Thompson Seedless, Perlette, Beauty Seedless	USA

	Kishmish Beli, Kishmish Chorni	USSR
<b>Guava</b>	Beaumont G-135	Australia
<b>Pomegranate</b>	Wonderful	USA
<b>Apple</b>	Red Spur, Oregon Spur	Italy
	Prima, Sir Prize, Jonafree Liberty, Priscilla- Scab resistant	USA
	Vance Delicious, Top Red, Royal Red	USA

Source: Ray, P.K. 2002.

### Selection:

Selection is basic to any crop improvement. Isolation of desirable plant types from the population is known as selection. It is one of the two fundamental steps of any breeding program viz., 1. creation of variation and 2 Selection. There are two agencies involved in carrying out the selection: one is Nature itself (Natural selection) and the other is man (artificial selection). Though both may complement each other in some cases,

they are mostly opposite in direction since their aims are different under the two conditions (nature and domestication). The effectiveness of selection primarily depends upon the degree to which phenotype reflects the genotype (Ray, P.K., 2002) It involves three basic principles: 1) it works on already existing germplasm, 2) it acts only through heritable alterations and 3) it works by favouring some individuals over others in reproduction.

Table 5: Some important varieties with improved fruit quality traits developed through selection

Fruit crop	Varieties	Method of breeding/Parents	Quality traits improved
<b>Aonla</b>	Kanchan (NA-4)	Selection	Suitable for processing
	Krishna (NA-5)	Selection	Suitable for processing
	Goma Aishwariya	Selection	Suitable for processing and export
<b>Cherry</b>	CITH-Cherry-2	Selection	Bold, attractive
<b>Guava</b>	Allahabad Safeda	Selection from Allahabad	White soft pulp, Sweet
	Arka Mridula	Selection From Allahabad Safeda	Soft seeded sweet, good pectin content, Keeping quality
	L-49	Selection from Allahabad Safeda	Highest vitamin C
	Lalit	-do-	Red colour pulp
	Try (G)-1	Selection	Off season, drought, sodicity tolerance
<b>Jackfruit</b>	PLR (J)-2	Selection from Pathirakkotai Local	good quality, fetch more price due to attractive characters and good keeping quality
<b>Litchi</b>	Rose Scented	Selection	Rose Scented, moderately juicy, soft and white flesh
<b>Mandarin</b>	Nagpur mandarin	Selection	Major position in mandarin markets, sweet, juicy and saffron coloured segments
<b>Mango</b>	Alphanso	Selection	Popular variety in domestic and export markets, yellow pleasant pulp, good keeping and

			processing quality
	Dashehari	Selection	Good keeping and table quality
	Pusa Surya	Selection from Eldon	Apricot yellow peel colour
	Langra	Selection	Turpentine flavour,
<b>Papaya</b>	CO-5	Selection	Cultivated mainly for papain production
	Coorg Honey Dew	Selection from Honey Dew	Gynodioecious
<b>Sapota</b>	Kirthabharti	Selection	Pulp is very sweet. Good for transportation to distant places
<b>Walnut</b>	CITH Walnut-1	Selection	Export purpose, bold nuts
<b>Pomegranate</b>	Ganesh	Selection from Alandi	Very soft seed,
	G-137	Selection from Ganesh	Soft seeded
<b>Apple</b>	Granny Smityh	Sel. From Lady Hamilton	
	Cameo	Sel. From Block of Red	--
<b>Peach</b>	Sharbati	Selection	Good flavour

Source: An individual selection and its traits taken from individual institute's website from where these varieties are developed and released and Department of Agriculture and Cooperation, 2012.

Table 6: Important selection in different fruits from pantnagar

Fruit crop	Cultivar	Parents
<b>Mango</b>	Pant Chandra	Seedling selection
	Pant Sindhuri	Clonal selection
<b>Guava</b>	Pant Prabhat	---
<b>Papaya</b>	Pant Papaya-1,2 & 3	----
<b>Bael</b>	Pant Urvashi, Pant sujata, Pant Aparna, Pant Shivani	---
<b>Peach</b>	Pant Peach-1	Seedling sel. From Sharbati
<b>Pear</b>	Pant Pear-3	---
<b>Plum</b>	Fla-12	Exotic type
	Pant Plum-1	---
<b>Aonla</b>	Pant Aonla-1	Seedling selection
<b>Karonda</b>	Pant Manohar, Pant Sudarshan, Pant Suvarna	Clonal selection
<b>Jackfruit</b>	Pant Mahima, Pant Garima	Clonal selection
<b>Lemon</b>	Pant Lemon-1	Clonal sel. from Kagazi Kalan

Source: Individual variety is taken from GBPUAT, Pantnagar Website ([www.gbpuat.ac.in](http://www.gbpuat.ac.in)).

### Hybridization

Hybridization involves the crossing of desired parents and a further selection of progenies. These desired parents are generally obtained after an appropriate screening of the natural populations and crop wild relatives being conserved under in-situ and ex-situ conditions hence, germplasm conservation is the most

important step, particularly for the utilization of wild species in breeding programs (Sharma et al. 2015). It helps in the selection of elite parents for the crossing and development of superior varieties. Conventional breeding involving approaches like hybridization, bridge crossing, distant hybridization, sibmating, half-sib mating, etc. had been advantageous in framing breeding strategies

for various crops and perennial fruit crops. Also, an admirable approach like doubled haploids could overcome some of the breeding limitations and is extremely useful for genetic studies such as gene mapping and genomics (Murovec and Bohanec, 2012). However, suitable technology for the development of superior fruit quality has to be adopted. Conventional breeding has contributed to fruit quality improvement with quite a good number of fruit trees. Many improved varieties have evolved through inter-specific hybridization or polyploidization or a combination of both (Jalilop 2015).

**Objectives:**

- To create genetic variation.
- Transfer of one or few qualitative characters.
- To develop biotic and abiotic resistant varieties.

**Types of hybridization**




Table 7: Some important varieties with improved fruit quality traits developed through hybridization breeding approaches

Fruit crop	Varieties	Method of breeding/Parents	Quality traits improved
<b>Mango</b>	Amrapali	Dashehari x Neelum	Fibreless, excellent taste, high carotenoids
	ArkaAnmol	Alphonso x Janardhan Pasand	Attractive skin colour, free from spongy tissue, good keeping quality, good sugar/acid blend
	Sindhu	Ratna x Alphonso	Seedless
	Konkan Ruchi	Neelum x Alphonso	Suitable for pickle making
<b>Papaya</b>	Arka Puneet	Alphanso x Banganpalli	Free from spongy tissue
	Arka Surya	Sunrise Solo x Pink Flesh Sweet	Red pulp, sweet
	CO-4	CO-1 x Washington	Purple pigment Variety
<b>Pomegranate</b>	Phule Arakta	Ganesh x Gul-e-shah Red	Dark red aril
	Amlidana	Ganesh x Nana	Suitable for anardana
	Mridula	Ganesh x Gulshah Red	Rose Pink, blood red aril, red

			rind, soft seeded
	Ruby	Ganesh x Kabul x Yercaud x Gulshah Rose Pink	The rind is pinkish yellow to reddish yellow. Fruit contains red and bold aril. It is soft seeded variety.
<b>Lime</b>	Rasraj	Kagzi Lime x Nepali Round	Good quality, juicy
<b>Guava</b>	ArkaAmulya	Allahabad Safeda x triploids	Round, firm, white fleshed, soft seeded, good keeping quality
<b>Sapota</b>	PKM-3	Guthi x Cricket Ball	Cluster Bearing habit
	Hisar Surkha	Apple Colour x Banarsi Surkha	Pink flesh
	ArkaKiran	Kamasari x purple local	Soft, yellow peel
<b>Custard apple</b>	Arka Sahan	<i>A. Atemoya</i> x <i>A. Squamosa</i>	Waxy skin, pleasant aroma
<b>Apple</b>	Ambred	Red Delicious x Ambri 157	Crisp, aromatic, juicy, shelf life
	Honeycrisp	MN447 x Northern Spy x unknown	Red peel, Rich flavor

Source: Individual hybrid is taken from their respective institutional website from where these varieties are developed and released

Table 8: Inter-generic hybrids of citrus

Crop	Parent	Name of hybrid
Citrange	<i>C.sinensis</i> x <i>P. trifoliata</i>	Troyer, Morton, Carrizo, Rusk
Citrumelo	<i>C.paradisii</i> x <i>P.trifoliata</i>	Swingle
Lemonimes	<i>C.limon</i> x <i>C.aurantifolia</i>	Parrine
Tangor	<i>C.reticulata</i> x <i>C.sinensis</i>	Temple, Clementine, Monreal
Tangerin	Robinson x Osceola	Sunbrust
Tangelo	<i>C.reticulata</i> x <i>C.paradisii</i>	Orlando, Sampson, Minneola

Source: Individual hybrid is taken from their respective institutional website from where these varieties are developed and released

### Mutation breeding:

Mutations are the heritable changes in the DNA sequence that are not derived from genetic segregation or recombination' (VanHarten 1998). The occurring mutations might be spontaneous or induced. The majority of mutation studies were concentrated upon annual crops mainly flowers and ornamental crops and special attention is required for attempting mutation in woody perennial crops. Induced mutations are highly effective in enhancing natural genetic resources, and have significantly assisted in developing improved fruit cultivars. Many commercial varieties have evolved mainly from spontaneous

mutations and chance seedlings. Bud mutations are a valuable source of variation (Usman et al. 2001), which could result in variants having characteristics including good fruit quality. Spontaneous mutants are reported in citrus (Liu et al. 2007) and mango (Medina 1997). For example, the variety Rosica has been reported in mango, which is the bud mutant of the Rosado-de-Ica variety and has large and good quality fruits.

### Mutagens

#### A. Physical mutagens

1. Ionizing radiations

(a) Particulate radiations-  $\alpha$ -rays, fast neutrons, thermal neutrons.

(b) Non-Particulate radiations- X-rays,  $\gamma$ -rays.

2. non-ionizing radiations- UV rays.

### B. Chemical mutagens.

1. Alkylating agents – Sulphur mustard, EMS (Ethyl Methyl Sulphonate), Ethylene Imine.

2. Acridine dyes – Acridine orange, ethidium bromide, acriflavine.

3. Base analogues – 5- bromouracil, 5-chlorouracil.

4. Others – Nitric acid, hydroxyl amine.

### Steps in mutation

- Well defined objective of the program.
- Selection of a variety of treatments.
- Part of a plant treated- seed, pollen grain, bud, cutting.
- Dose of mutagen- LD50
- Treatment- method and treatment.

Table 9: Important varieties with improved fruit quality traits developed through Mutation Breeding

Fruit crops	Cultivar	Year	Mutagens	Improved fruit traits
<b>Mango</b>	Rosica	1966	Spontaneous mutation	Large and good quality
<b>Apple</b>	Golden Haidegg	1986	$\gamma$ -rays	Fruit size
	McIntosh 8F-2-32	1970	$\gamma$ -rays	Skin colour
	Senbatsu-Fuji-2-Kei	1985	$\gamma$ -rays	Fruit colour
<b>Grapefruit</b>	Rio Red	1984	thN	Fruit colour
	Star Ruby	1970	thN	Seedless
<b>Indian jujube</b>	Mahong	1986	thN	Round, pink rose sweeter taste
<b>Loquat</b>	Shiro-mogi	1981	$\gamma$ -rays	Fruit size
<b>Orange</b>	Xuegan 9-12-1	1983	$\gamma$ -rays	Seedless
	Hongju 420	1986	$\gamma$ -rays	Seedless
	Eureka 22	1987	X-rays	Fruit set, fruit quality
	Valencia 2	1987	X-rays	Fruit quality
<b>Peach</b>	Magnif	1968	$\gamma$ -rays	Large, red skin
	Plovdiv	1981	$\gamma$ -rays	Large, fruit quality
<b>Sweet cherry</b>	Lapins	1983	X-rays	Larger size, firmer
	Compact Lambert	1964	X-rays	Compact growth
	Ferrovía spur	1992	X-rays	Dwarfness
<b>Pear</b>	Fuxiangyanghongdli	1983	$\gamma$ -rays	Eating, cooking quality
<b>Almond</b>	Supernova	1987	$\gamma$ -rays	Late maturity
<b>Fig</b>	Bol	1979	$\gamma$ -rays	Not specified
<b>Banana</b>	Novaria	1993	$\gamma$ -rays	Earliness
	Al-beely	2007	$\gamma$ -rays	High yield
<b>Japanese pear</b>	Gold Nijisseiki	1993	$\gamma$ -rays	Disease resistance
<b>Papaya</b>	Pusa Nanha	1986	$\gamma$ -rays	Dwarfness
<b>Plum</b>	Spurdente-Ferco	1988	$\gamma$ -rays	Earliness
<b>Pomegranate</b>	Karabakh	1979	$\gamma$ -rays	--
<b>Sea</b>	zyriank			

## buckthorn

Source:(Predieri, 2001)

Table 10: Other important varieties developed through mutation breeding

Fruit crop	Variety	Parents	Nature of mutation
Mango	Rosica	Rosa-do-delca (Peru)	Natural
	Davis Haden	Haden	Natural
Grape	Marvel Seedless	Delight	Induced
Banana	High Gate	Gros Michel	Natural
	Motta Poovan	Poovan	Natural
	Krishna Vazahi	Virupakshi	Natural
Orange	Washington Novel	Navel Orange	Natural
Grape fruit	Foster	Walter	Natural
	Red Blush	Thompson	Natural
	Thompson	Marsh Seedless	Natural
	Star Ruby	Hudson	Induced

Source: Ray, P.K. 2002

### Polyploidy breeding

Polyploidy refers to the presence of more than two complete sets of chromosomes per cell nucleus, which has been considered a ubiquitous phenomenon in plant evolution and diversification (Ray, P.K. 2002). A polyloid individual arising within one or between populations of a single species is denominated autopolyploid, while the term allopolyploid refers to individuals of hybrid origin. Allopolyploids are often divided into two sub-classes: true and segmental allopolyploids. The formation of true allopolyploids involves hybridization between distantly related species. In this case, the divergent chromosome complements do not pair with each other, resulting in the formation of bivalents during meiosis and a disomic inheritance pattern. On the other hand, segmental allopolyploids originate from hybridization between closely related species with partially differentiated genomes (Shukla and Shukla, 2004). Therefore, segmental allopolyploids may undergo univalent, bivalent and/or multivalent pairing during meiosis, being considered intermediate types between true allopolyploids and

auto-polyploids. Now day Colchicine @ 0.2-0.8% is mostly used for the induction of polyloids.

### Morphological and cytological aspects of polyloids

1. Slow growth and delayed flowering.
2. Increased flesh weight but reduced dry weight.
3. Increase in size but the decrease in the number of leaves, flowers, and fruits.
4. Larger cell size.
5. Stomatal gourd cell larger (stomata count/unit area lesser).
6. Variation in ploidy from species to species.

### Application of polyploidy

- Quality improvement: e.g., Seedlessness
- Direct use as new var. or species.
- Inter-specific gene transfer.
- Widening genetic base of existing allopolyploids.
- Tracking the origin of natural allopolyploids.

Table 11: Varieties developed through polyploidy breeding in fruit crops

Ploidy level	Autopolyploidy type	Crop
3x	Auto triploid	Banana, Apple, Tahiti lime
4x	Auto tetraploid	Aonla, beal, Litchi, Phalsa, Jackfruit, grapes, Ber (cv. Umran)
6x	Auto hexaploid	Kiwifruit, Persimmon
8x	Auto octaploid	Ber (cv. Gola, Illaichi)
	<b>Allo-polyploidy types</b>	<b>Crop</b>
	Amphidiploids/Allo-tetraploid	Mango
	Allo-hexaploid	European plum
	Allo-octaploid	Strawberry
<b>Aneuploidy</b>		
	Aneuploid-82	Pusa Srijan (Guava dwarf rootstock)

Source: Ray. P.K. 2002.

Table 12: Ploidy level of varieties and their origin in different fruit crops

Fruit crop	Common interest	Polyploidy level	Origin	Reference
Banana	Edible fruit	Autopolyploidy/ allopolyploidy	Synthetic/natural	Silvaetal.(2001)
Grape	Edible fruit	Autopolyploidy/ allopolyploidy	Synthetic/natural	Motosugietal.(2002)
Apple	Edible fruit	Autopolyploidy/allopolyploidy	Synthetic/natural	Janicketal.(1996)
Strawberry	Edible fruit	Allopolyploidy	Natural	Whitaker(2011)
Kiwifruit	Edible fruit	Autopolyploidy	Natural	Hopping(1994)
Tahiti Lime	Edible fruit	Allopolyploid	Natural	Morton(1987)
Plum	Edible fruit	Allopolyploid	Natural	BennettandLeitch(1995)

Source: Ray. P.K. 2002.

### Somatic hybridization

Somatic hybridization involves the fusion of partial and complete genome exchange, which might result in the development of novel varieties (Loredana et al. 2012). That means a combination of the nuclear, chloroplast and mitochondrial genomes in a novel arrangement (Singh and Rajam 2009) might aid in the development of novel varieties with desirable fruit quality traits. Since the transgenic approaches are limited by social barriers, these approaches might be useful to

surpass this limitation in the development of new cultivars particularly having improved polygenic fruit quality. Somatic hybridization has widely been attempted for developing inter-specific/inter-generic crosses in the Citrus group, to overcome problems like sexual incompatibility, polyembryony and pollen or ovule sterility (Singh and Rajam 2009). Loredana et al. (2012) developed a hybrid and two cybrids by protoplast fusion of sweet orange (*Citrus sinensis* L. Osbeck) and lemon (*C. limon* L. Burm.). These cybrids

exhibited improved essential oil composition, responsible for imparting aroma in citrus.

### **Molecular Approaches**

The work related to molecular biology in the case of perennial crops is very scarce as compared to annuals (Pena and Seguin 2001). Molecular approaches have proved to overcome some of the breeding problems in fruit trees. Prediction of colour, taste, shelf-life behaviour, texture, and nutrition characteristics by detection of marker genes before the tree even starts to bear fruit would be of much practical utility. Germplasm maintenance and evaluation, an integral part of conventional breeding, is more expensive in terms of time, labour, money and other inputs (Myles 2013). The approaches of genomics and marker-assisted selection (MAS) are more advantageous, particularly in woody perennials, including fruit crops (Kumar et al. 2012). Besides, advanced technologies in genetic transformation have been proven to shorten the juvenile phases of the tree, resistance to biotic stresses, and phytoremediation in perennial trees (Pena and Seguin 2001). Some of the major molecular based technologies and significant attempts made for fruit quality improvement are discussed hereunder.

### **Marker assisted selection and QTLs what this? All words and abbreviations**

The term QTL was first coined by Gelderman (1975). The markers tightly linked with the trait of interest are very informative in utilizing them in the identification and further selection process. Molecular markers have been in vogue for identifying the trait of interest, SSR and SNPs being the most preferred. QTLs provide this information with manifold applications in breeding for complex fruit quality traits. The close linkage with the other quality parameters is advantageous when both are of superior and acceptable traits by breeders. However, if one is not acceptable in hybrids then it becomes difficult to segregate them apart in the progeny. To know this, series of crosses between cultivars with superior and poor traits are made and their population information

provides QTL, which could be used as markers in MAS. Such analyses for fruit texture were conducted by Longhi et al. (2012), wherein the located QTLs by SSR and SNP markers. The most common method of QTL is interval mapping. For examples:

1. Costa et al. (2005) in their study, utilized gene-specific molecular markers (ACS genes) for studying their effects on ethylene production and shelf life in apple and could position them on a linkage map. They reported the marker for the identification of apple cultivars having a good shelf life.
2. Huan et al. (2012) developed SSR and AFLP markers linked to major gene loci involved in the fruit shape index of apples (*Malus domestica*). It was also reported that the fruit shape is controlled by five genes.
3. Davey et al. (2006) identified QTLs related to ascorbic acid in fruit skin and flesh in apple.

### **Candidate genes approach and source to sink interaction in fruits**

According to the concept that a molecular polymorphism is directly connected to phenotype, the candidate gene technique includes breeding in reverse, going from the gene to the trait of interest (Pflieger et al. 2001). The candidate gene approach for understanding the signalling mechanism and biosynthetic pathway might be a useful approach because fruit quality is directly proportional to the metabolites present in it. Flavonoids, antioxidants, phenolics, active ingredients, pigments, etc. give the good flavour, taste and colour to the fruits. Secondary metabolites are produced using particular or common metabolic pathways. Enzymes that are connected to these pathways are encoded by structural and regulatory genes. Source-to-sink relation is very complicated for improving the traits (Nookaraju et al. 2010). Biosynthetic pathways of pigments have been extensively studied in fruit crops like apple (Honda et al. 2002), peach (Li et al. 2012), sweet cherry (Wei et al. 2015), litchi (Zan-wei et al. 2011), mangosteen (Palapol et al. 2009) and citrus (Kato et al. 2004). For examples:

1. Anthocyanin-related gene expressions in fruit pericarp of many cultivars of litchi (Zan-wei et al., 2011).
2. The integrated approach of metabolomics and transcriptomics related to mango fruit peel colouration.
3. Candidate genes involved in sugar and organic acid metabolism in apple and 12 candidate genes were allocated to 4 linkage groups of the peach genome (Etienne et al., 2002).
4. Candidate genes are involved in taste development in the citrusfruit (Kato et al., 2004).
5. Candidate genes are responsible for the texture, ethylene production Costa et al. (2005), sugar and organic acid content (Etienne et al., 2002) and polyphenols (Chagne et al., 2012) in apple have been studied.

type of marker-assisted selection in which genetic markers available on the whole genome expresses its phenotypic variation Myles, (2013). The application of genomics and its function in battling few of the breeding difficulties has also been studied in apple and grapes by Myles (2013). The Apple Breed Data Base offers easy admittance for the documentation of molecular markers linked to fruit quality traits (e.g., skin colour, shape, or taste) (Antofie et al. 2007). Genomic analysis related to produce quality at developmental stages has been assessed in fruit trees like bayberry, citrus, apple and peach (Zhu et al. 2013; Cer-cos et al. 2006; Terol et al. 2008;).

### Transgenic approach

Transgenic or genetically altered fruit crops involve introduction of desired genes (Dias and Ortiz 2014) principally for the development of elite varieties with improved quality traits. *Agrobacterium*-mediated transformation, fruit quality gene was successfully transferred through protoplast transformation with the retrieval of transgenic plants in citrus (Guo et al., 2005). The first transgenic plant was developed is Tobacco in 1983.

### Genomics approach:

Genomics includes study of whole genetic information related to particular fruit crop. The term genomics was coined by Dr.T. Roderick. It can be used for selection of better parents and for hybrid development. Genomic selection (GS) is a

Table 13: Transgenic varieties in fruit crops

Fruit crop	Variety	Trait	Year	References
Papaya	Rainbow, Sunup	Rainbow, Sunup Resistant to papaya ring spot virus	1998	Gonsalves et al. (1998)
Grapes	Pinot Noir			Franks et al. (1998)
Banana	DH-Pahang	Hepatitis B	-	Kumar et al. (2006)
Plum	Honey Sweet	Resistant to Plum Pox Virus (PPV)	2009	Scorza et al., (1994)
Apple	Arctic Apple	Resistant to browning	2015	Murata et al. (2000)
	Arctic golden delicious	Resistant to browning		
	Arctic granny smith	Resistant to browning		
Strawberry	Apel	fruit firmness	-	Jimenez-Bermudez et al. (2002)

Table 14: Transgenic fruit crops developed by *Agrobacterium* mediated transformation

Species	Traits	Plasmid	Transgenes	References
---------	--------	---------	------------	------------

<b>Orange</b>				
<i>Citrus sinensis</i>	Method optimization	<i>pGA482GG</i>	<i>gusA, nptII</i>	Oliveira <i>et al.</i> , 2009
<i>Citrus aurantifolia</i>	Resistance to virus	<i>pBin19-sgfp</i>	<i>nptII, sgfp, p23</i>	Fagoaga <i>et al.</i> , 2006
<i>Poncirus trifoliata</i>	Enhanced salt tolerance	<i>pBin438</i>	<i>nptII, AhBADH</i>	Fu. <i>et.al.</i> , 2011
<b>Papaya</b>				
<i>Carica papaya</i>	Resistance to PRSV	<i>pRPTW</i>	PRSV replicase gene, neo	Chen <i>et al.</i> ,2001
<b>Pomegranate</b>				
<i>Punica granatum</i>	Method optimization	<i>pBIN19-sgfp</i>	<i>nptII, gfp</i>	Terakami <i>et al.</i> , 2007
<i>Fragaria spp.</i> (Strawberry)	Modulation of fruit softening	<i>pBI121</i>	antisense of endo- <i>b1,4</i> -glucanase	Lee & Kim,2011

Table 15: Permits and notifications of transgenic fruits approved

Crop	Trait	Genes
<b>Apple</b>	Reduced polyphenol oxidase	PPO suppression transgene, <i>nptII</i>
	Non-browning reduced polyphenol oxidase	Polyphenol oxidase antisense, <i>PGAS1, PGAS2</i>
<b>Banana</b>	Bunchy top resistance	Replicase associated protein, replicase inverted repeat, <i>nptII</i>
<b>Grape rootstock</b>	Grapevine fan leaf virus resistance	Coat protein gene, heat shock 90 homologous genes, <i>nptII</i>
<b>Grapefruit</b>	Aphid resistance	Agglutin coat protein, <i>GUS</i> .
<b>Papaya</b>	Female to male (or) hermaphrodite	<i>EST116, ESTS, FSH11, FSH19</i>



Image 1 : Pro Enzyme induced apple

### Problems and prospects

The intake of fresh market fruits in the United Nationlike stone fruits and apple,has been stable or decreasing (Perez *et al.*, 2001). Surveys propose that, primary reason is that the consumers

are not pleased with quality offruits(Stockwin, 1996). Our present strategy for quality as breeders has been to choose subtle alterations in a high-quality cultivar to fill a specific niche. A little alteration that will solve an issue in the consumer market may be a different ripening duration, a change in colour, or a greater sugar to acid ratio. Although this breeding strategy has had great

success, it hasn't necessarily improved customer access to fruit of the utmost quality. Our understanding of the factors influencing fruit quality has improved from the use of molecular approaches to the study of fruit ripening (Giovannoni, 2001; Knee, 2002) yet, it is not sufficient to make molecular selection and modification at practical level as a primary approach to improve quality traits for consumers. Long juvenile phases, prolonged gestation periods, plant architecture and inadequate planting material are the key issues with fruit breeding programmes aimed at enhancing the qualitative attributes of fruit crops.

Breeders may have to think outside the box, at least until we have a better knowledge of fruit quality traits. Fruit quality traits are defined by cultivars that may perhaps be centuries old such as 'Montmorency' cherry (17th century), 'Bing' cherry (1800s), 'Golden Delicious', 'Delicious', 'Granny Smith', and 'Jonathan' apples (18th and 19th century), 'Chinese Cling' peach (18th century), 'Belle of Georgia', and 'Elberta' peach (1850s), 'Bartlett' and 'Bosc' Pear (18th century), Smooth Cayenne pineapple (1819) and 'Fuerte' avocado (1911), all are very dominant germplasm in breeding programs. One strategy for breeding for better fruit quality is to diligently screen for better fruit quality and not be deterred by poor appearance (Janick and Moore, 1996). Perhaps we

should have to select quality attributes as per future thrust to improve. More varieties of small, firm fruit, like the recently released "Pixie Crunch" apple, or fruit that can be consumed without creating a mess, like the Zee Sweet series of peaches, nectarines, and pluots, which can be consumed while still crisp due to their high sugar and low acid ratios, may be included in successful releases. In present, fruit processing industry is growing and now major challenges for fruit breeder is to improve the variety in way that have good flavour, texture and taste after cut into the pieces and lightly processed.

### Conclusions

Fruit crops need integrated techniques that combine traditional and non-conventional methods for breeding varieties with positive characteristics since they are constrained by a variety of restrictions compared to short-lived and seasonal crops. As polygenes are in control of controlling fruit quality, integration becomes essential for achieving the desired improvement. The biotechnological approaches deliver an accurate, reliable and easy way for breeding fruit trees for improvement of fruit quality. Also, it lessens the time, effort and patience of the breeder, when dealing with breeding related activities of fruit trees. These methods might aid in the development of high-quality fruits to fulfil both domestic and global fruit demand.

### References

- Antofie, A., Lateur, M., Oger, R., Patocchi, A., Durel, C. E., & Van de Weg, W. E. (2007). A new versatile database created for geneticists and breeders to link molecular and phenotypic data in perennial crops: The applebreed database. *Bioinformatics*, 23(7), 882–891.
- Arias, R. S., Borrone, J. W., Tondo, C. L., Kuhn, D. N., Irish, B. M., & Schnell, R. J. (2012). Genomics of Tropical Fruit Tree Crops. *Genomics of Tree Crops*, 209–239. [://doi.org/10.1007/978-1-4614-0920-57](https://doi.org/10.1007/978-1-4614-0920-57).
- Baldwin, E.A. (2002). Fruit flavour, volatile metabolism and consumer perceptions. p. 89-106. In: M. Knee (ed.), *Fruit Quality and its Biological Basis*. Sheffield Academic Press, Sheffield, UK.
- Barritt, B.H. (2001). Apple quality for consumers. *International Dwarf Fruit Tree Assoc.* 34:54-46.
- Bennett, M. (1995). Nuclear DNA amounts in angiosperms. *Annals of Botany*, 76(2), 113–176. <https://doi.org/10.1006/anbo.1995.1085>.

- Breeding of fruits and plantation crops - researchgate.* (n.d.). Retrieved March 2, 2023, [https://www.researchgate.net/publication/361410582\\_Breeding\\_of\\_Fruits\\_and\\_Plantation\\_Crops](https://www.researchgate.net/publication/361410582_Breeding_of_Fruits_and_Plantation_Crops)
- Brummell, D. A., & Harpster, M. H. (2001). Cell wall metabolism in fruit softening and quality and its manipulation in transgenic plants. *Plant Mol. Biol.*, 47(1-2): 311–340.
- Callahan, M.A. (2003). Breeding for fruit quality. *Acta Horticulture*, 622: 255-302, <http://dx.doi.org/10.17660/ActaHortic.2003.622.27>.
- Cercós, M., Soler, G., Iglesias, D. J., Gadea, J., Forment, J., & Talón, M. (2006). Global analysis of gene expression during development and ripening of citrus fruit flesh. A proposed mechanism for citric acid utilization. *Plant Molecular Biology*, 62(4-5), 513–527.
- Chagné, D., Krieger, C., Rassam, M., Sullivan, M., Fraser, J., André, C., Pindo, M., Troglio, M., Gardiner, S. E., Henry, R. A., Allan, A. C., McGhie, T. K., & Laing, W. A. (2012). QTL and candidate gene mapping for polyphenolic composition in Apple Fruit. *BMC Plant Biology*, 12(1), 12. <https://doi.org/10.1186/1471-2229-12-12>.
- Chen, G., Ye, C. M., Huang, J. C., Yu, M., & Li, B. J. (2001). Cloning of the papaya ringspot virus (PRSV) replicase gene and generation of PRSV-resistant papayas through the introduction of the PRSV replicase gene. *Plant Cell Reports*, 20(3), 272–277.
- Costa, F., Stella, S., Van de Weg, W. E., Guerra, W., Cecchinell, M., Dallavia, J., Koller, B., & Sansavini, S. (2005). Role of the genes MD-ACO1 and MD-ACS1 in ethylene production and shelf life of Apple (*Malus domestica* Borkh). *Euphytica*, 141(1-2), 181–190.
- Davey, M. W., Kenis, K., & Keulemans, J. (2006). Genetic control of fruit vitamin C contents. *Plant Physiology*, 142(1), 343–351. <https://doi.org/10.1104/pp.106.083279>
- de-Oliveira, M. L., Febres, V. J., Costa, M. G., Moore, G. A., & Otoni, W. C. (2009). High-efficiency agrobacterium-mediated transformation of citrus via sonication and vacuum infiltration. *Plant Cell Reports*, 28(3), 387–395.
- Dias, J.S., & Ortiz, R. (2014). Advances in transgenic vegetable and fruit breeding. *Agricultural Sciences*, 05(14), 1448–1467.
- Elia, L. (2002). *Producing a profitable peach*. Fruit Grower June:6.
- Etienne, C., Rothan, C., Moing, A., Plomion, C., Bodénès, C., Svanella-Dumas, L., Cosson, P., Pronier, V., Monet, R., & Dirlwanger, E. (2002). Candidate genes and qtls for sugar and organic acid content in peach [*Prunus persica* (L.) Batsch]. *Theoretical and Applied Genetics*, 105(1), 145–159.
- Fagoaga, C., López, C., de Mendoza, A. H., Moreno, P., Navarro, L., Flores, R., & Peña, L. (2007). Post-transcriptional gene silencing of the p23 silencing suppressor of citrus tristeza virus confers resistance to the virus in transgenic Mexican lime. *Biotechnology and Sustainable Agriculture 2006 and beyond*, 211–213.

- Franks, T., He, D.G., & Thomas, M.R. (1998). Regeneration of transgenic *Vitis vinifera* L. Sultana plants: genotypic and phenotypic analysis. *Molecular Breeding* 4: 321–333.
- Fu, X.-Z., Khan, E. U., Hu, S.-S., Fan, Q.-J., & Liu, J.-H. (2011). Overexpression of the betaine aldehyde dehydrogenase gene from *Atriplex hortensis* enhances salt tolerance in the transgenic trifoliolate orange (*Poncirus Trifoliata* L. Raf.). *Environmental and Experimental Botany*, 74, 106–113.
- Giovannoni, J. 2001. Molecular biology of fruit maturation and ripening. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 52:725–49.
- Gonsalves, D., Gonsalves, C., Ferreira, C., & Fitch, M. (1998). Transgenic virus-resistant papaya: From hope to reality in controlling papaya ringspot virus in Hawaii. *APSnet Feature Articles*.
- Guo, W., Duan, Y., Olivares-Fuster, O., Wu, Z., Arias, C. R., Burns, J. K., & Grosser, J. W. (2005). Protoplast transformation and regeneration of transgenic valencia sweet orange plants containing a juice quality-related pectin methylesterase gene. *Plant Cell Reports*, 24(8), 482–486.
- Harlan, J. R. (1971). Agricultural Origins: Centers and noncenters. *Science*, 174(4008), 468–474. <https://doi.org/10.1126/science.174.4008.468>
- Honda, C., Kotoda, N., Wada, M., Kondo, S., Kobayashi, S., Soejima, J., Zhang, Z., Tsuda, T., & Moriguchi, T. (2002). Anthocyanin biosynthetic genes are coordinately expressed during red coloration in Apple Skin. *Plant Physiology and Biochemistry*, 40(11), 955–962.
- Hopping, M. E. (1994). Flow cytometric analysis of *Actinidia* species. *New Zealand Journal of Botany*, 32(1), 85–93.
- Jalikop, S.H. (2015). Allied species utilization in fruit improvement—way forward. Souvenir of lead papers and abstracts of National meet on distant hybridization in horticultural crop improvement. 25–35.
- Janick, J., & Moore, J. N. (1998). *Fruit breeding vol. I tree and tropical fruits*. Wiley India Pvt. Ltd.
- Jiménez-Bermúdez Silvia, Redondo-Nevaldo José, Muñoz-Blanco Juan, Caballero José L., López-Aranda José M., Valpuesta, V., Pliego-Alfaro, F., Quesada, M. A., & Mercado José A. (2002). Manipulation of strawberry fruit softening by antisense expression of a pectate lyase gene. *Plant Physiology*, 128(2), 751–759.
- Kader, A.A. (2002). Fruits in the global market. p. 1–16. In: M. Knee (ed.), *Fruit Quality and its Biological Basis*. Sheffield Academic Press, Sheffield, UK.
- Kato, M., Ikoma, Y., Matsumoto, H., Sugiura, M., Hyodo, H., & Yano, M. (2004). Accumulation of carotenoids and expression of carotenoid biosynthetic genes during maturation in citrus fruit. *Plant Physiology*, 134(2), 824–837.
- Kenis, K., Keulemans, J., & Davey, M. W. (2008). Identification and stability of qtls for fruit quality traits in Apple. *Tree Genetics & Genomes*, 4(4), 647–661.

- Kumar, A., & Ellis, B. E. (2001). The phenylalanine ammonia-lyase gene family in raspberry: structure, expression, and evolution. *Plant Physiology*, 127(1), 230–239.
- Kumar, G. B. S., Srinivas, L., Ganapathi, T. R., & Bapat, V. A. (2006). Hepatitis B surface antigen expression in transgenic tobacco (*Nicotiana tabacum*) plants using four different expression cassettes. *Plant Cell, Tissue and Organ Culture*, 84(3), 315–323.
- Kupferman, E. (2002). Critical aspects of harvest and quality management. Postharvest Information Network. [www.postharvest.tfrec.wsu.edu/EMK2002A.pdf](http://www.postharvest.tfrec.wsu.edu/EMK2002A.pdf).
- Lee, Y. K., & Kim, I.-J. (2011). Modulation of fruit softening by antisense suppression of endo- $\beta$ -1,4-glucanase in strawberry. *Molecular Breeding*, 27(3), 375–383.
- Li, L., Ban, Z.-J., Li, X.-H., Wu, M.-Y., Wang, A.-L., Jiang, Y.-Q., & Jiang, Y.-H. (2012). Differential expression of anthocyanin biosynthetic genes and transcription factor PCMYB10 in Pears (*Pyrus communis* L.). *PLoS ONE*, 7(9).
- Longhi, S., Moretto, M., Viola, R., Velasco, R., & Costa, F. (2011). Comprehensive QTL mapping survey dissects the complex fruit texture physiology in Apple (*Malus X domestica* borkh.). *Journal of Experimental Botany*, 63(3), 1107–1121.
- Loredana, A., Nicasio, T., Sergio, B., Tonia, S., Agatino, R., & Giuseppe, R. (2012). Genetic improvement of Citrus fruits: new somatic hybrids from *Citrus sinensis* (L.) Osb. And *Citruslimon* (L.) Burm. *F. D. Res. Intl.* 48:284–290.
- Motosugi, H., Okudo, K., Kataoka, D., & Naruo, T. (2002). Comparison of growth characteristics between diploid and colchicine-induced tetraploid grape rootstocks. *Engei Gakkai Zasshi*, 71(3), 335–341.
- Murata, M., Nishimura, M., Murai, N., Haruta, M., Homma, S., & Itoh, Y. (2001). A transgenic apple callus showing reduced polyphenol oxidase activity and lower browning potential. *Bioscience, Biotechnology, and Biochemistry*, 65(2), 383–388.
- Murovec, J., & Bohanec, B. (2012). Haploids and doubled haploids in plant breeding. *Plant Breeding*. <https://doi.org/10.5772/29982>
- Myles, S. (2013). Improving fruit and wine: What does genomics have to offer? *Trends in Genetics*, 29(4), 190–196.
- Nookaraju, A., Upadhyaya, C. P., Pandey, S. K., Young, K. E., Hong, S. J., Park, S. K., & Park, S. W. (2010). Molecular approaches for enhancing sweetness in fruits and vegetables. *Scientia Horticulturae*, 127(1), 1–15.
- Oguntibeju, O., J., E., & J., A. (2013). The role of fruit and vegetable consumption in human health and disease prevention. *Diabetes Mellitus - Insights and Perspectives*. <https://doi.org/10.5772/50109>

- Palapol, Y., Ketsa, S., Lin-Wang, K., Ferguson, I. B., & Allan, A. C. (2009). A MYB transcription factor regulates anthocyanin biosynthesis in mangosteen (*Garcinia mangostana* L.) fruit during ripening. *Planta*, 229(6), 1323–1334.
- Pena, L., & Seguin, A. (2001). Recent advances in the genetic transformation of trees. *Trends Biotechnology*, 19(12):500–506.
- Perez, A., Lin, B-H. & Allshouse, J. (2001). Demographic profile of apple consumption in the United States. *Fruit and Tree Nuts S&O/FTS* 292:37–47.
- Pflieger, S., Lefebvre, V., & Causse, M. (2001). The candidate gene approach in plant genetics: a review. *Molecular Breeding*, 7(4):275–291.
- Predieri, S., & Gatti, E. (2000). Effects of gamma radiation on plum (*Prunus salicina* Lindl.) „Shiro”. *Advances in Horticultural Science*, 14: 215–223.
- Ray, P.K. (2002). *Breeding Tropical and Subtropical Fruits*. Springer Berlin, Heidelberg publishers, ISBN,978-3-540-42855-8.
- Redgwell, R.J., & Fischer, M. (2002). Fruit texture, cell wall metabolism and consumer perceptions. p. 46–88. In: M. Knee (ed.), *Fruit Quality and its Biological Basis*. Sheffield Academic Press, Sheffield, UK.
- Richard, H. (2014). *GMO Fruit Crops*, Extension Pecan & Pistachio Specialist, <https://mvd.iaea.org/>.
- Scorza, R., Ravelonandro, M., Callahan, A. M., Cordts, J. M., Fuchs, M., Dunez, J., & Gonsalves, D. (1994). Transgenic plums (*Prunus domestica* L.) express the plum pox virus coat protein gene. *Plant Cell Reports*, 14(1), 18–22.
- Seymour, G.B., & Manning, K. (2002). Genetic control of fruit ripening. p. 253–274. In: M. Knee (ed.), *Fruit Quality and its Biological Basis*. Sheffield Academic Press, Sheffield, UK.
- Sharma, T.V.R.S., Singh, S., Singh, A.K., & Singh, P.K. (2015). A note on possibility of utilizing wild relatives of crop plants present in Andaman & Nicobar Islands for creating new variability. Souvenir of lead papers and abstracts of National meet on Distant Hybridization in Horticultural crop improvement. 1-1.
- Shukla, A.K., & Vashishtha, B.B. (2004). *Fruit Breeding: Approaches and Achievements*. International Book Distribution Co., Charbagh, Lucknow.
- Silva, S. O., Souza Junior, M. T., Alves, É. J., Silveira, J. R. S., & Lima, M. B. (2001). Banana breeding program at Embrapa. *Cropps Breeding and Applied Biotechnology*, 1(4), 399–436.
- Simmonds, N.W. (1979b). *Principles of Crop Improvement*. Longman, London.
- Singh, J. (2002). *Basic Horticulture*. Kalyani Publishers, New Delhi.
- Singh, R.V., Chand, D., Tyagi, V., Singh, A.K., Singh, S.P., Singh, S. & Binda P.C. (2005). Priorities for introduction of fruit crops in India. *Indian Journal of plant genetic resources*, 18(1): 67-68.
- Singh, S., & Rajam, M. V. (2009). Citrus biotechnology: Achievements, limitations and Future Directions. *Physiology and Molecular Biology of Plants*, 15(1).

- Smith, D. L., Abbott, J. A., & Gross, K. C. (2002). Down-regulation of tomato  $\beta$ -galactosidase 4 results in decreased fruit softening. *Plant Physiology*, 129(4), 1755–1762.
- Stockwin, W. (1996). Boost stone fruit quality. *Fruit Grower*, June 13–16.
- Sun, H. H., Zhao, Y. B., Li, C. M., Chen, D. M., Wang, Y., Zhang, X. Z., & Han, Z. H. (2011). Identification of markers linked to major gene loci involved in determination of fruit shape index of Apples (*Malus domestica*). *Euphytica*, 185(2), 185–193.
- Terakami, S., Matsuta, N., Yamamoto, T., Sugaya, S., Gemma, H., & Soejima, J. (2007). Agrobacterium-mediated transformation of the dwarf pomegranate (*punica granatum* L. var. *nana*). *Plant Cell Reports*, 26(8), 1243–1251.
- Terol, J., Conesa, A., Colmenero, J.M., Cercos, M., Tadeo, F., Agustí, J., Alos, E., & Andres, F. (2008). The areas of flavor, health, color and ripening. *BMC Genomics* 9: 351.
- Usman, M., Fatima, B., & Muhammad, J.J. (2001). Breeding in mango. *International Journal of Agriculture and Biology*, 4:522–526.
- Van Harten, A.M. (1998). Mutation breeding: theory and practical applications. Cambridge Univ. Press, Cambridge.
- Vrebalov, J., Ruezinsky, D., Padmanabhan, V., White, R., Medrano, D., Drake, R., Schuch, W., & Giovannoni, J. (2002). A MADS-box gene necessary for fruit ripening at the tomato *ripening-inhibitor* (*rin*) locus. *Science*, 296(5566), 343–346.
- Wargovich, M.J. (2000). Anticancer properties of fruits and vegetables. *Hort. Sci.*, 35573575.
- Wei, H., Chen, X., Zong, X., Shu, H., Gao, D., & Liu, Q. (2015). Comparative transcriptome analysis of genes involved in anthocyanin biosynthesis in the red and yellow fruits of sweet cherry (*Prunus avium* L.). *PLOS ONE*, 10(3).
- Whitaker, V. M. (2011). Applications of molecular markers in strawberry. *Journal of Berry Research*, 1(3), 115–127.
- Winkel-Shirley, B. (2001). Flavonoid biosynthesis. A colorful model for genetics, biochemistry, Cell Biology, and biotechnology. *Plant Physiology*, 126(2), 485–493.
- Zan-Wei, Y.-Z., Hu, F.-C., Hu, G.-B., Li, X.-J., Huang, X.-M., & Wang, H.-C. (2011). Differential expression of anthocyanin biosynthetic genes in relation to anthocyanin accumulation in the pericarp of *Litchi Chinensis* sonn. *PLoS ONE*, 6(4), 1-11.
- Zhu, C., Feng, C., Li, X., Xu, C., Sun, C., & Chen, K. (2013). Analysis of expressed sequence tags from Chinese bayberry fruit (*Myrica rubra* sieb. and zucc.) at different ripening stages and their association with Fruit Quality Development. *International Journal of Molecular Sciences*, 14(2), 3110–3123.