

Fixed Points of ξ - (α, β) -Contractive Mappings in b -Metric Spaces

Abstract: In the paper [*Some new observations on Geraghty and Ćirić type results in b -metric spaces*, Mathematics, **7**, (2019), doi: 10.3390/math7070643] Mlaiki *et al.* introduced (α, β) -type contraction in order to generalize the contraction mapping defined by Pant and Panicker. Also, in the paper [*Some fixed point results in b -metric spaces and b -metric-like spaces with new contractive mappings*, Axioms, **10(2)**, (2021), 15 pages, doi: 10.3390/axioms10020055] Jain and Kaur presented the concepts of ξ -contractive mappings. Now, the aim of the present article is to introduce ξ - (α, β) -contractive mappings in b -metric spaces by combining the concepts (α, β) -type contraction and ξ -contractive mappings. Also, we establish some fixed point results for newly defined mappings. Our results generalize various theorems in literature. In support, we provide an example.

Keywords: b -metric space; (α, β) -admissible mappings; (α, β) -type contraction; ξ -contractive mappings; Cauchy Sequence; fixed point.

MSC: 47H10;54H25.

1 Introduction and Preliminaries

In 1922, Banach [1] initiated the study of constructive theory in metric space. Banach Contraction Principle (BCP) is one of the most useful and important theorems in classical functional analysis. This theorem has been extended and generalized in many directions. In 1969, Nadlar [2] proved BCP for a multivalued mapping. Various other extensions of this

principle can be found in [3, 4, 5, 6, 7, 8]. Also, in the last decade Samet *et al.* [9] defined α - ψ -contractive mapping to generalize BCP. Later, in 2015, Chandok [10] introduced the concept of (α, β) -admissible mappings. In 2016, Pant and Panicker [11] obtained some results for (α, β) -Geraghty type contractive mapping. Mlaiki *et al.* [12] generalize the results of Pant and Panicker in 2019. Recently, Jain and Kaur [13] presented the concept of ξ -contractive mappings. As an extension of metric space, Bakhtin [14] has studied the concept of b -metric space (by weakening the triangle inequality in metric space) as:

Definition 1.1. [14] Let X be a non-empty set. Then a mapping $d : X \times X \rightarrow [0, +\infty)$ is called a b -metric, if for all $x, y, z \in X$, the following conditions hold:

- (bm1) $d(x, y) = 0$ if and only if $x = y$;
- (bm2) $d(x, y) = d(y, x)$;
- (bm3) $d(x, y) \leq 2(d(x, z) + d(z, y))$.

The pair (X, d) is called a b -metric space.

In 1998, Czerwik [15] presented the concept of b -metric space in the more general form as shown below:

Definition 1.2. [15] Let X be a non-empty set and $s \geq 1$ be a given real. Then a mapping $d : X \times X \rightarrow [0, +\infty)$ is called a b -metric, if for all $x, y, z \in X$, the following conditions hold:

- (b1) $d(x, y) = 0$ if and only if $x = y$;
- (b2) $d(x, y) = d(y, x)$;
- (b3) $d(x, y) \leq s(d(x, z) + d(z, y))$.

The pair (X, d) is called a b -metric space. In 2010, Khamsi and Hussain [16] used the notion of b -metric space under the name metric type space. **More on b -metric spaces can be studied in [17, 18, 19, 20, 21, 22, 23, 24, 25].**

To prove our results, we need the following concepts and results of literature.

Definition 1.3. [26] Let (X, d) be a b -metric space. Then, a sequence $\{x_n\}$ in X is called :

- (i) *Cauchy sequence*, if for each $\epsilon > 0$ there exist $n_0 \in \mathbb{N}$ such that $d(x_n, x_m) < \epsilon$, for all $n, m \geq n_0$.
- (ii) *Convergent*, if there exists $l \in X$ such that for each $\epsilon > 0$ there exist $n_0 \in \mathbb{N}$ such that $d(x_n, l) < \epsilon$, for all $n \geq n_0$. In this case, sequence $\{x_n\}$ is said to *converge* to l .

Definition 1.4. [26] A b -metric space (X, d) is said to be *complete* if every Cauchy sequence is convergent in it.

Lemma 1.5. [27] Let (X, d) be a b -metric space with $s \geq 1$ and suppose that sequences $\{x_n\}$ and $\{y_n\}$ converge to x and $y \in X$, respectively. Then

$$\frac{1}{s^2}d(x, y) \leq \liminf_{n \rightarrow +\infty} d(x_n, y_n) \leq \limsup_{n \rightarrow +\infty} d(x_n, y_n) \leq s^2d(x, y).$$

In particular, if $x = y$, then $\lim_{n \rightarrow +\infty} d(x_n, y_n) = 0$.

Moreover, for any $z \in X$,

$$\frac{1}{s}d(x, z) \leq \liminf_{n \rightarrow +\infty} d(x_n, z) \leq \limsup_{n \rightarrow +\infty} d(x_n, z) \leq sd(x, z).$$

Lemma 1.6. [28, 29] Every sequence $\{x_n\}$ of elements from a b -metric space (X, d) , having the property that there exists $\lambda \in [0, 1)$ such that $d(x_n, x_{n+1}) \leq \lambda d(x_{n-1}, x_n)$ for every $n \in \mathbb{N}$, is Cauchy.

Definition 1.7. [10] Let (X, d) be a b -metric space, $T : X \rightarrow X$ and $\alpha, \beta : X \times X \rightarrow [0, +\infty)$. Then mapping T is said to be (α, β) -admissible, if $\alpha(x, y) \geq 1$ and $\beta(x, y) \geq 1$ implies $\alpha(Tx, Ty) \geq 1$ and $\beta(Tx, Ty) \geq 1$, for all $x, y \in X$.

Definition 1.8. [10] Let (X, d) be a b -metric space, $T : X \rightarrow X$ and $\alpha, \beta : X \times X \rightarrow [0, +\infty)$. Then X is said to be (α, β) -regular, if $\{x_n\}$ is a sequence in X such that $\{x_n\}$ converges to x , $\alpha(x_n, x_{n+1}) \geq 1$, $\beta(x_n, x_{n+1}) \geq 1$ for all n , there exists a subsequence $\{x_{n_k}\}$ of $\{x_n\}$ such that $\alpha(x_{n_k}, x_{n_k+1}) \geq 1$, $\beta(x_{n_k}, x_{n_k+1}) \geq 1$ for all k and $\alpha(x, Tx) \geq 1$, $\beta(x, Tx) \geq 1$.

Definition 1.9. [10] Let Ψ be the set of all functions $\psi : [0, +\infty) \rightarrow [0, +\infty)$ such that ψ is continuous, strictly increasing and $\psi(0) = 0$.

Definition 1.10. [10] Let Θ be the set of all functions $\theta : [0, +\infty) \rightarrow [0, 1)$ such that for any bounded sequence $\{t_n\}$ of positive reals, $\theta(t_n) \rightarrow 1$ implies $t_n \rightarrow 0$.

In 2016, Pant and Panicker [11] introduced the following contractive mapping.

Definition 1.11. [11] Let (X, d) be a b -metric space with $s \geq 1$. Then a mapping $T : X \rightarrow X$ is said to be (α, β) -Geraghty type contractive mapping, if there exists $\alpha, \beta : X \times X \rightarrow [0, +\infty)$, $\psi \in \Psi, \theta \in \Theta$ such that

$$\alpha(x, Tx)\beta(y, Ty)\psi(s^3d(Tx, Ty)) \leq \theta(\psi(N(x, y)))\psi(N(x, y)), \quad (1.1)$$

for all $x, y \in X$, where $N(x, y) = \max \left\{ d(x, y), d(x, Tx), d(y, Ty), \frac{d(x, Ty)+d(Tx, y)}{2s} \right\}$.

In 2019, Mlaiki *et al.* [12] established the following contraction mapping which generalizes the contractive mapping defined by Pant and Panicker.

Definition 1.12. [12] Let (X, d) be a b -metric space with $s > 1$. Then mapping $T : X \rightarrow X$ is said to be an (α, β) -type contraction, if there exists $\alpha, \beta : X \times X \rightarrow [0, +\infty)$, $\psi \in \Psi$, $\epsilon > 1$ such that

$$\alpha(x, Tx)\beta(y, Ty)\psi(s^\epsilon d(Tx, Ty)) \leq \psi(N(x, y)), \quad (1.2)$$

for all $x, y \in X$, where $N(x, y) = \max \left\{ d(x, y), d(x, Tx), d(y, Ty), \frac{d(x, Ty)+d(Tx, y)}{2s} \right\}$.

Theorem 1.13. [12] Let (X, d) be a complete b -metric space with $s > 1$ and $T : X \rightarrow X$ be a mapping such that the following hold:

- (i) T is (α, β) -type contraction;
- (ii) T is (α, β) -admissible;
- (iii) there exists $x_0 \in X$ such that $\alpha(x_0, Tx_0) \geq 1$ and $\beta(x_0, Tx_0) \geq 1$;
- (iv) either T is continuous or X is (α, β) -regular;

Then T has a unique fixed point.

Following concepts and results were introduced by Jain and Kaur [13].

Definition 1.14. [13] For any $m \in \mathbb{N}$, define Ξ_m to be the set of all functions $\xi : [0, +\infty)^m \rightarrow [0, +\infty)$ such that

- (ξ_1) $\xi(t_1, t_2, \dots, t_m) < \max\{t_1, t_2, \dots, t_m\}$ if $(t_1, t_2, \dots, t_m) \neq (0, 0, \dots, 0)$;
- (ξ_2) if $\left\{ t_i^{(n)} \right\}_{n \in \mathbb{N}}$, $1 \leq i \leq m$, are m sequences in $[0, +\infty)$ such that

$$\limsup_{n \rightarrow +\infty} t_i^{(n)} = t_i < +\infty \text{ for all } i = 1 \text{ to } m, \text{ then}$$

$$\liminf_{n \rightarrow +\infty} \xi \left(t_1^{(n)}, t_2^{(n)}, \dots, t_m^{(n)} \right) \leq \xi(t_1, t_2, \dots, t_m).$$

Definition 1.15. [13] Let (X, d) be a b -metric space with $s \geq 1$. The mapping $T : X \rightarrow X$ is said to be an ξ -contractive mapping of type-I, if there exists $\xi \in \Xi_4$ and

$$d(Tx, Ty) \leq \frac{1}{s} \xi \left(d(x, y), d(x, Tx), d(y, Ty), \frac{d(x, Ty) + d(Tx, y)}{2s} \right), \quad (1.3)$$

for all $x, y \in X$.

Theorem 1.16. [13] Let (X, d) be a complete b -metric space with $s \geq 1$ and $T : X \rightarrow X$ be an ξ -contractive mapping of type-I. Then T has a unique fixed point.

Definition 1.17. [13] Let (X, d) be a b -metric space with $s \geq 1$. The mapping $T : X \rightarrow X$ is said to be an ξ -contractive mapping of type-II, if there exists $\xi \in \Xi_5$ and

$$d(Tx, Ty) \leq \frac{1}{s} \xi \left(d(x, y), d(x, Tx), d(y, Ty), \frac{d(x, Ty)}{2s}, d(Tx, y) \right), \quad (1.4)$$

for all $x, y \in X$.

Theorem 1.18. [13] Let (X, d) be a complete b -metric space with $s \geq 1$ and $T : X \rightarrow X$ be an ξ -contractive mapping of type-II. Then T has a unique fixed point.

2 Main Results

In the present work, first we introduce ξ - (α, β) -contractive mapping of type-I and ξ - (α, β) -contractive mapping of type-II in b -metric spaces. After that, we prove some unique fixed point theorems which generalize various results in literature.

Definition 2.1. Let (X, d) be a b -metric space with $s \geq 1$ and $T : X \rightarrow X$ be a mapping. We say that T is an ξ - (α, β) -contractive mapping of type-I, if there exists $\alpha, \beta : X \times X \rightarrow [0, +\infty)$, $\psi \in \Psi$, $\xi \in \Xi_4$, $\epsilon > 1$ such that

$$\begin{aligned} & \alpha(x, Tx)\beta(y, Ty)\psi(s^\epsilon d(Tx, Ty)) \\ & \leq \psi \left(s^{\epsilon-1} \xi \left(d(x, y), d(x, Tx), d(y, Ty), \frac{d(x, Ty) + d(Tx, y)}{2s} \right) \right), \end{aligned} \quad (2.1)$$

for all $x, y \in X$.

Definition 2.2. Let (X, d) be a b -metric space with $s \geq 1$ and $T : X \rightarrow X$ be a mapping. We say that T is an ξ - (α, β) -contractive mapping of type-II, if there exists $\alpha, \beta : X \times X \rightarrow [0, +\infty)$, $\psi \in \Psi$, $\xi \in \Xi_5$, $\epsilon > 1$ such that

$$\begin{aligned} & \alpha(x, Tx)\beta(y, Ty)\psi(s^\epsilon d(Tx, Ty)) \\ & \leq \psi \left(s^{\epsilon-1} \xi \left(d(x, y), d(x, Tx), d(y, Ty), \frac{d(x, Ty)}{2s}, d(Tx, y) \right) \right), \end{aligned} \quad (2.2)$$

for all $x, y \in X$.

Theorem 2.3. Let (X, d) be a complete b -metric space with $s \geq 1$ and $T : X \rightarrow X$ be a mapping such that the following hold:

- (i) T is ξ - (α, β) -contractive mapping of type-I;
- (ii) T is (α, β) -admissible;
- (iii) there exists $x_0 \in X$ such that $\alpha(x_0, Tx_0) \geq 1$ and $\beta(x_0, Tx_0) \geq 1$;
- (iv) either T is continuous or X is (α, β) -regular with respect to T ;

Then T has a fixed point. If $Tx = x$ implies $\beta(x, x) \geq 1$, then fixed point of T is unique.

Proof. Define a sequence $\{x_n\}$ in X as $x_n = Tx_{n-1}$ for all $n \geq 1$. Assume that any two consecutive terms of sequence $\{x_n\}$ are distinct, otherwise T has a fixed point. Now by condition (ii), (iii) and using induction, we get

$$\alpha(x_n, x_{n+1}) \geq 1 \quad \text{and} \quad \beta(x_n, x_{n+1}) \geq 1 \quad \text{for all } n \geq 1.$$

First, we prove that $\{x_n\}$ is a Cauchy sequence. For this, let $n \in \mathbb{N}$. Consider

$$\begin{aligned} & \psi(s^\epsilon d(x_n, x_{n+1})) \\ & \leq \alpha(x_{n-1}, x_n) \beta(x_n, x_{n+1}) \psi(s^\epsilon d(x_n, x_{n+1})) \\ & = \alpha(x_{n-1}, Tx_{n-1}) \beta(x_n, Tx_n) \psi(s^\epsilon d(Tx_{n-1}, Tx_n)) \\ & \leq \psi \left(s^{\epsilon-1} \xi \left(d(x_{n-1}, x_n), d(x_{n-1}, x_n), d(x_n, x_{n+1}), \frac{d(x_{n-1}, x_{n+1}) + d(x_n, x_n)}{2s} \right) \right) \end{aligned}$$

which implies that

$$s^\epsilon d(x_n, x_{n+1}) \leq s^{\epsilon-1} \xi \left(d(x_{n-1}, x_n), d(x_{n-1}, x_n), d(x_n, x_{n+1}), \frac{d(x_{n-1}, x_{n+1})}{2s} \right),$$

i.e.,

$$d(x_n, x_{n+1}) \leq \frac{1}{s} \xi \left(d(x_{n-1}, x_n), d(x_{n-1}, x_n), d(x_n, x_{n+1}), \frac{d(x_{n-1}, x_{n+1})}{2s} \right) \quad (2.3)$$

$$\begin{aligned} & < \frac{1}{s} \max \left\{ d(x_{n-1}, x_n), d(x_{n-1}, x_n), d(x_n, x_{n+1}), \frac{d(x_{n-1}, x_{n+1})}{2s} \right\} \\ & = \frac{1}{s} \max \left\{ d(x_{n-1}, x_n), \frac{d(x_{n-1}, x_{n+1})}{2s} \right\} \\ & \leq \frac{1}{s} \max \left\{ d(x_{n-1}, x_n), \frac{d(x_{n-1}, x_n) + d(x_n, x_{n+1})}{2} \right\}, \end{aligned} \quad (2.4)$$

which implies that

$$d(x_n, x_{n+1}) < \frac{1}{s} d(x_{n-1}, x_n), \quad \text{for all } n \geq 1. \quad (2.5)$$

Case 1: If $s > 1$, then by Lemma 1.6 in view of (2.5), $\{x_n\}$ is a Cauchy sequence.

Case 2: If $s = 1$, then by (2.5), sequence $\{d(x_n, x_{n+1})\}$ is monotonically decreasing and also it is bounded below, therefore, $d(x_n, x_{n+1}) \rightarrow k$ for some $k \geq 0$. Suppose that $k > 0$. Now taking $\liminf_{n \rightarrow +\infty}$ in (2.3), we have $k \leq \xi(k, k, k, k')$,

where

$$k' = \limsup_{n \rightarrow +\infty} \frac{d(x_{n-1}, x_{n+1})}{2} \leq \limsup_{n \rightarrow +\infty} \frac{d(x_{n-1}, x_n) + d(x_n, x_{n+1})}{2} = k.$$

Now, $k \leq \xi(k, k, k, k') < \max\{k, k, k, k'\} = k$, a contradiction, therefore,

$$\lim_{n \rightarrow +\infty} d(x_n, x_{n+1}) = 0. \quad (2.6)$$

Suppose that $\{x_n\}$ is not a Cauchy sequence, then there exists $\varepsilon > 0$ such that for any $r \in \mathbb{N}$, there exists $m_r > n_r \geq r$ such that

$$d(x_{m_r}, x_{n_r}) \geq \varepsilon. \quad (2.7)$$

Also, assume that m_r is smallest natural greater than n_r such that (2.7) holds. Now

$$\begin{aligned} \varepsilon &\leq d(x_{m_r}, x_{n_r}) \\ &\leq d(x_{m_r}, x_{m_r-1}) + d(x_{m_r-1}, x_{n_r}) \\ &< d(x_{m_r}, x_{m_r-1}) + \varepsilon \\ &< d(x_r, x_{r-1}) + \varepsilon, \end{aligned}$$

so by using (2.6) and taking $\lim r \rightarrow +\infty$, we get

$$\lim_{r \rightarrow +\infty} d(x_{m_r}, x_{n_r}) = \varepsilon. \quad (2.8)$$

Consider

$$\begin{aligned} &\psi(d(x_{m_r+1}, x_{n_r+1})) \\ &\leq \alpha(x_{m_r}, x_{m_r+1})\beta(x_{n_r}, x_{n_r+1})\psi(d(x_{m_r+1}, x_{n_r+1})) \\ &= \alpha(x_{m_r}, Tx_{m_r})\beta(x_{n_r}, Tx_{n_r})\psi(d(Tx_{m_r}, Tx_{n_r})) \\ &\leq \psi\left(\xi\left(d(x_{m_r}, x_{n_r}), d(x_{m_r}, x_{m_r+1}), d(x_{n_r}, x_{n_r+1}), \frac{d(x_{m_r}, x_{n_r+1}) + d(x_{m_r+1}, x_{n_r})}{2}\right)\right), \end{aligned}$$

which implies that

$$\begin{aligned} &d(x_{m_r+1}, x_{n_r+1}) \\ &\leq \xi\left(d(x_{m_r}, x_{n_r}), d(x_{m_r}, x_{m_r+1}), d(x_{n_r}, x_{n_r+1}), \frac{d(x_{m_r}, x_{n_r+1}) + d(x_{m_r+1}, x_{n_r})}{2}\right). \end{aligned}$$

Now

$$\begin{aligned} &d(x_{m_r}, x_{n_r}) \\ &\leq d(x_{m_r}, x_{m_r+1}) + d(x_{m_r+1}, x_{n_r+1}) + d(x_{n_r+1}, x_{n_r}) \\ &\leq d(x_{m_r}, x_{m_r+1}) + d(x_{n_r+1}, x_{n_r}) + \\ &\quad \xi\left(d(x_{m_r}, x_{n_r}), d(x_{m_r}, x_{m_r+1}), d(x_{n_r}, x_{n_r+1}), \frac{d(x_{m_r}, x_{n_r+1}) + d(x_{m_r+1}, x_{n_r})}{2}\right), \end{aligned}$$

thus, by taking $\liminf r \rightarrow +\infty$ on both sides and also using (2.6) and (2.8), we get $\varepsilon \leq 0 + 0 + \xi(\varepsilon, 0, 0, \varepsilon')$, where

$$\begin{aligned} \varepsilon' &= \limsup_{r \rightarrow +\infty} \frac{d(x_{m_r}, x_{n_r+1}) + d(x_{m_r+1}, x_{n_r})}{2} \\ &\leq \limsup_{r \rightarrow +\infty} \frac{d(x_{m_r}, x_{n_r}) + d(x_{n_r}, x_{n_r+1}) + d(x_{m_r+1}, x_{m_r}) + d(x_{m_r}, x_{n_r})}{2} \\ &= \frac{\varepsilon + 0 + 0 + \varepsilon}{2} = \varepsilon. \end{aligned}$$

Thus, $\varepsilon \leq \xi(\varepsilon, 0, 0, \varepsilon') < \max\{\varepsilon, 0, 0, \varepsilon'\} = \varepsilon$, a contradiction. Hence $\{x_n\}$ is a Cauchy sequence in (X, d) . But (X, d) be a complete b -metric space, therefore, there exists $x \in X$ such that $x_n \rightarrow x$.

If T is continuous, then $x_n \rightarrow x$ implies $x_{n+1} = Tx_n \rightarrow Tx$. But, as limit is unique, therefore, $Tx = x$.

If X is (α, β) -regular with respect to T , then $\alpha(x, Tx) \geq 1$ and $\beta(x, Tx) \geq 1$. Suppose that $Tx \neq x$. Now

$$\begin{aligned} \psi(s^\varepsilon d(Tx_n, Tx)) &\leq \alpha(x_n, Tx_n)\beta(x, Tx)\psi(s^\varepsilon d(Tx_n, Tx)) \\ &\leq \psi\left(s^{\varepsilon-1}\xi\left(d(x_n, x), d(x_n, x_{n+1}), d(x, Tx), \frac{d(x_n, Tx) + d(x, Tx_n)}{2s}\right)\right), \end{aligned}$$

which implies that

$$s^\varepsilon d(Tx_n, Tx) \leq s^{\varepsilon-1}\xi\left(d(x_n, x), d(x_n, x_{n+1}), d(x, Tx), \frac{d(x_n, Tx) + d(x, x_{n+1})}{2s}\right),$$

i.e.,

$$d(x_{n+1}, Tx) \leq \frac{1}{s}\xi\left(d(x_n, x), d(x_n, x_{n+1}), d(x, Tx), \frac{d(x_n, Tx) + d(x, x_{n+1})}{2s}\right).$$

Taking $\liminf n \rightarrow +\infty$ on both sides and using Lemma 1.5, we get

$$\frac{1}{s}d(x, Tx) \leq \frac{1}{s}\xi(0, 0, d(x, Tx), l),$$

i.e.,

$$d(x, Tx) \leq \xi(0, 0, d(x, Tx), l),$$

where,

$$l = \limsup_{n \rightarrow +\infty} \frac{d(x_n, Tx) + d(x, x_{n+1})}{2s} \leq \limsup_{n \rightarrow +\infty} \frac{sd(x, Tx) + 0}{2s} = \frac{d(x, Tx)}{2}.$$

Hence,

$$d(x, Tx) \leq \xi(0, 0, d(x, Tx), l) < \max\{0, 0, d(x, Tx), l\} = d(x, Tx),$$

which is a contradiction, therefore, $Tx = x$.

If $Ty = y$ for some $y \in X$, then by the given condition, $\beta(y, y) \geq 1$. Now suppose that $x \neq y$, and consider

$$\begin{aligned} \psi(s^\varepsilon d(x, y)) &= \psi(s^\varepsilon d(Tx, Ty)) \\ &\leq \alpha(x, Tx)\beta(y, Ty)\psi(s^\varepsilon d(Tx, Ty)) \\ &\leq \psi\left(s^{\varepsilon-1}\xi\left(d(x, y), d(x, Tx), d(y, Ty), \frac{d(x, Ty) + d(y, Tx)}{2s}\right)\right), \end{aligned}$$

which implies that

$$s^\varepsilon d(x, y) \leq s^{\varepsilon-1}\xi\left(d(x, y), d(x, Tx), d(y, Ty), \frac{d(x, Ty) + d(y, Tx)}{2s}\right),$$

i.e.,

$$\begin{aligned}
d(x, y) &\leq \frac{1}{s} \xi \left(d(x, y), 0, 0, \frac{d(x, y)}{s} \right) \\
&< \frac{1}{s} \max \left\{ d(x, y), 0, 0, \frac{d(x, y)}{s} \right\} \\
&= \frac{d(x, y)}{s},
\end{aligned}$$

a contradiction, therefore, $x = y$. \square

Example 2.4. Let $A = \left\{ \frac{1}{\sqrt{n}} \mid n \in \mathbb{N} \right\} \cup \{0\}$, $B = \{n + 1 \mid n \in \mathbb{N}\}$ and $X = A \cup B$. Define $d : X \times X \rightarrow [0, +\infty)$ by $d(x, y) = |x - y|^2$ for all $x, y \in X$. Then d is a b -metric on X with $s = 2$.

Define $T : X \rightarrow X$ by $T\left(\frac{1}{\sqrt{n}}\right) = \frac{1}{\sqrt{2(n+1)}}$, $T(n + 1) = n + 2$ for all $n \in \mathbb{N}$ and $T(0) = 0$.

Define $\alpha, \beta : X \times X \rightarrow [0, +\infty)$, $\psi \in \Psi$, $\xi \in \Xi_4$ as:

$$\alpha(x, y) = \beta(x, y) = \begin{cases} 1, & \text{if } x, y \in A, \\ 0, & \text{otherwise,} \end{cases} \quad (2.9)$$

$\psi(t) = t$, and

$$\xi(t_1, t_2, t_3, t_4) = \begin{cases} \frac{\max\{t_1, t_2, t_3, t_4\}}{1+t_1}, & \text{if } t_1 > 0, \\ \frac{1}{2} \max\{t_2, t_3, t_4\}, & \text{otherwise.} \end{cases}$$

Now for all $x, y \in X$, (2.1) is satisfied. Also T is (α, β) -admissible, X is (α, β) -regular with respect to T and for $x_0 = 1 \in X$, $\alpha(x_0, Tx_0) \geq 1$ and $\beta(x_0, Tx_0) \geq 1$. Thus, all the conditions of Theorem 2.3 are satisfied and 0 is only fixed point of T .

Now following remark improves our main result Theorem 2.3.

Remark 2.5. Theorem 2.3 is also valid, if term $\frac{d(x, Ty) + d(Tx, y)}{2s}$ in (2.1) is replaced by $\frac{d(x, Ty) + d(Tx, y)}{\delta s}$, where δ is a real number defined by

$$\delta = \begin{cases} 2, & \text{if } s = 1, \\ \delta', & \text{if } 1 < s \leq 2, \\ 1, & \text{if } s > 2, \end{cases}$$

where δ' is any number in $\left(\frac{2}{s}, 1 + \frac{1}{s}\right)$.

Corollary 2.6. (Theorem 1.13) Let (X, d) be a complete b -metric space with $s > 1$ and $T : X \rightarrow X$ be a mapping such that the following hold:

- (i) T is (α, β) -type contraction;
- (ii) T is (α, β) -admissible;
- (iii) there exists $x_0 \in X$ such that $\alpha(x_0, Tx_0) \geq 1$ and $\beta(x_0, Tx_0) \geq 1$;
- (iv) either T is continuous or X is (α, β) -regular;

Then T has a unique fixed point.

Proof. Take $\xi \in \Xi_4$ defined by $\xi(t_1, t_2, t_3, t_4) = \frac{1}{s^{\epsilon-1}} \max\{t_1, t_2, t_3, t_4\}$, then by Theorem 2.3, we get the result. \square

Corollary 2.7. (Theorem 1.16) Let (X, d) be a complete b -metric space with $s \geq 1$ and $T : X \rightarrow X$ be an ξ -contractive mapping of type-I. Then T has a unique fixed point.

Proof. Take $\alpha, \beta : X \times X \rightarrow [0, +\infty)$ such that $\alpha(x, y) = \beta(x, y) = 1$, for all $x, y \in X$, and $\psi \in \Psi$ by $\psi(t) = t$. Then by Theorem 2.3, T has a unique fixed point. \square

Remark 2.8. Example 2.4 is applicable for Theorem 2.3, but not for Corollary 2.7.

Remark 2.9. In view of Remark 2.5, Corollaries 2.6 and 2.7 are also valid, if term $\frac{d(x, Ty) + d(Tx, y)}{2s}$ is replaced by $\frac{d(x, Ty) + d(Tx, y)}{\delta s}$, where δ is same as defined in Remark 2.5.

Proof of our next result is on a similar manner as the proof of Theorem 2.3.

Theorem 2.10. Let (X, d) be a complete b -metric space with $s \geq 1$ and $T : X \rightarrow X$ be a mapping such that the following conditions hold:

- (i) T is ξ - (α, β) -contractive mapping of type-II;
 - (ii) T is (α, β) -admissible;
 - (iii) there exists $x_0 \in X$ such that $\alpha(x_0, Tx_0) \geq 1$ and $\beta(x_0, Tx_0) \geq 1$;
 - (iv) either T is continuous or X is (α, β) -regular with respect to T ;
- Then T has a fixed point. If $Tx = x$ implies $\beta(x, x) \geq 1$, then fixed point of T is unique.

Corollary 2.11. (Theorem 1.18) Let (X, d) be a complete b -metric space with $s \geq 1$ and $T : X \rightarrow X$ be an ξ -contractive mapping of type-II. Then T has a unique fixed point.

Proof. Take $\alpha, \beta : X \times X \rightarrow [0, +\infty)$ such that $\alpha(x, y) = \beta(x, y) = 1$, for all $x, y \in X$, and $\psi \in \Psi$ by $\psi(t) = t$. Then by Theorem 2.10, T has a unique fixed point. \square

Author Contributions: Both the authors have equally contributed in the planning, execution and analysis of study.

Funding: This research received no external funding.

Conflicts of interest: The authors declare no conflict of interest.

References

- [1] Banach, S.; *Sur les operations dans les ensembles abstraits et leur application aux equations integrales*, Fundamenta Mathematicae, **3**, (1922), 133-181.
- [2] Nadler, Jr.S.B.; *Multivalued contraction mappings*, Pacific Journal of Mathematics, **30**, (1969), 475-488.
- [3] A. Meir and E. Keeler, *A thorem on contractive mappings*, Journal of Mathematical Analysis and applications, **28**, (1969), 326-329.
- [4] R. Kannan, *Some results on fixed points*, Bulletin of the Calcutta Mathematical Society, **25**, (1972), 727-730.
- [5] L. B. Ciric, *A generalization of Banachs contraction principle*, Proceeding of the American Mathematical Society, **45**, (1974), 267-273.
- [6] B. E. Rhoades, *A comparison of various definitions of contractive mappings*, Transactions of The American Mathematical Society, **226**, (1977).

- [7] B. E. Rhoades, *Some theorems on weakly contractive maps*, Non-linear Analysis TMA, **47(4)**, (2001), 2683-2694.
- [8] D. Wardowski, *Fixed points of a new type of contractive mappings in complete metric spaces*, Fixed Point Theory and Applications, **94**, (2012), doi: 10.1186/1687-1812-2012-94.
- [9] Samet, B.; Vetro, C.; and Vetro, P.; *Fixed point theorems for α - ψ -contractive type mappings*, Nonlinear Analysis: Theory, Methods and Applications, **75(4)**, (2012), 21542165.
- [10] Chandok, S.; *Some fixed point theorems for (α, β) -admissible Geraghty type contractive mappings and related results*, Mathematical Science, **9**, (2015), 127-135.
- [11] Pant, R.; and Panicker, R.; *Geraghty and Ćirić type fixed point theorems in b-metric spaces*, Journal of Nonlinear Science and Applications, **9**, (2016), 5741-5755.
- [12] Mlaiki, N.; Dedovic, N.; Aydi, H.; Filipović, M.G.; Mohsin, B.B.; and Radenović, S.; *Some new observations on Geraghty and Ćirić type results in b-metric spaces*, Mathematics, **7**, (2019), doi: 10.3390/math7070643.
- [13] Jain, K.; and Kaur, J.; *Some fixed point results in b-metric spaces and b-metric-like spaces with new contractive mappings*, Axioms, **10(2)**, (2021), 15 pages, doi: 10.3390/axioms10020055.
- [14] Bakhtin, I.A.; *The contraction mapping principle in almost metric spaces*, Functional Analysis, **30**, (1989), 26-37.
- [15] Czerwik, S.; *Nonlinear set-valued contraction mapping in b-metric spaces*, Atti del Seminario Matematico e Fisico dell'Universita di Modena e Reeggio Emilia, **46**, (1998), 263-276.
- [16] Khamsi, M.A.; and Hussain, N.; *KKM mappings in metric type spaces*, Nonlinear Analysis, **73**, (2010), 3123-3129.
- [17] Aydi, H.; Bota, M.F.; Karapinar, E.; and Mitrović, S.; *A fixed point theorem for set-valued quasi-contractions in b-metric spaces*, Fixed Point Theory and Applications, (2012), **2012:88**.
- [18] Kir, M.; and Kiziltunc, H.; *On some well known fixed point theorems in b-metric spaces*, Turkish Journal of Analysis and Number Theory, **1**, (2013), 13-16.
- [19] Latif, A.; Parvaneh, V.; Salimi, P.; and Al-Mazrooei, A.E.; *Various Suzuki type theorems in b-metric spaces*, Journal of Nonlinear Science and Applications, **8(4)**, (2015), 363-377.
- [20] Aleksić, S.; Mitrović, Z.D.; and Radenović, S.; *On some recent fixed point results for single and multivalued mappings in b-metric spaces*, Fasciculi Mathematici, (2018), doi: 10.1515/fascmath-2018-0013.
- [21] Karapinar, E.; Czerwik, S.; and Aydi, H.; *(α, ψ) -Meir-Keeler contraction mappings in generalized b-metric spaces*, Journal of Function Spaces, (2018), doi: 10.1155/2018/3264620.
- [22] Karapinar, E.; Mitrović, Z.D.; Öztürk, A.; and Radenović, S.; *On a theorem of Ćirić in b-metric spaces*, Rendiconti del Circolo Matematico di Palermo Aeries 2, March 2020, doi: 10.1007/s12215-020-00491-9.

- [23] Mitrović, Z.D.; Parvaneh, V.; Mlaiki, N.; Hussain, N.; and Radenović, S.; *On some new generalizations of Nadler contraction in b-metric spaces*, Cogent Mathematics and Statistics, (2020), **7:1**, 1760189.
- [24] Kanwal, S.; Hanif, U.; Noorwali, M.E.; and Alam, M.A.; *On fixed-point results of generalized contractions*, Journal of Function Spaces, (2022), 11 pages, doi: 10.1155/2022/9167716.
- [25] Nallaselli, G.; Gnanaprakasam, A.J.; Mani, G.; and Ege, O.; *Solving integral equations via admissible contraction mappings*, Filomat, (2022), 4947-4961, doi: 10.2298/FIL2214947N.
- [26] Boriceanu, M.; *Fixed point theory for multivalued generalized contraction on a set with two b-metrics*, Studia Universitatis Babeş-Bolyai Mathematica, **LIV(3)**, (2009), 1-14.
- [27] Aghajani, A.; Abbas, M.; and Roshan, J.R.; *Common fixed point of generalized weak contractive mappings in partially ordered b-metric spaces*, Mathematica Slovaca, **64(4)**, (2014), 941-960, doi: 10.2478/s12175-014-0250-6.
- [28] Miculescu, R.; and Mihail, A.; *New fixed point theorems for set-valued contractions in b-metric spaces*, Journal of Fixed Point Theory and Applications, **19(3)**, (2017), 2153-2163.
- [29] Suzuki, T.; *Basic inequality on a b-metric space and its applications*, Journal of Inequalities and Applications, (2017), **2017:256**, doi: 10.1186/s13660-017-1528-3.