

## Original Research Article

### **Microbiology of Chronic Suppurative Otitis Media amongst Children attending a Tertiary Health Facility in north-western Nigeria.**

#### **Abstract:**

#### **Background:**

Chronic suppurative otitis media (CSOM) is a global middle ear disease worst in developing nations of the world. The burden in our environment of repeated episodes of childhood CSOM otorrhoea, and the resultant difficulty in the management of these patients because of improper or inadequate treatment is quite worrisome.

**Aim:** The aim of this study was to prospectively analyze the aspirated middle ear exudates of paediatric patients with CSOM for both facultative aerobic bacteria and fungi, with their antimicrobial sensitivity patterns.

**Materials and Methods:** A total of 166 children clinically diagnosed with CSOM were enrolled in the study, of which 219 middle ear discharge samples were obtained using sterile disposable plastic Pasteur pipettes and cultured for microbial flora. Drug susceptibility testing for the isolated microbes was conducted using the Kirby-Bauer disc diffusion method.

#### **Results:**

The ratio of facultative/aerobic bacteria to fungi was 2.1: 1, while the ratio of Gram-positive to Gram-negative aerobic bacteria was 1: 1.7. The most common causative aerobic organisms isolated were *Staphylococcus aureus* 41/202 (20.3%), *Pseudomonas aeruginosa* 30/202 (14.9%), *Klebsiella pneumonia* 23/202 (11.4%), and *Proteus mirabilis* 21/202 (10.4%). *Aspergillus species* 47/202 (23.3%) and *Candida albicans* 21/202 (10.4%) were the commonest fungal isolates. The antimicrobial profile of aerobic isolates revealed maximum sensitivity to gentamycin (91.1%), ceftriaxone (90.1%), and ciprofloxacin (85.6%); while the isolated fungi were completely sensitive to all the antifungals.

**Conclusion:** The main isolates were Gram-negative aerobic bacteria, though Gram-positive aerobes and fungi were also isolated. Their varying antimicrobial susceptibility brings to the fore, the therapeutic significance of these drugs in CSOM management.

**Keywords:** *Chronic suppurative otitis media, aerobes, fungi, Sokoto*

## **Introduction:**

The ear as an organ receives minimal attention but is vulnerable to a variety of infections.<sup>1</sup> By definition, chronic suppurative otitis media (CSOM) otherwise referred to as active chronic otitis media (active COM) is characterized by chronic inflammatory changes of the mucoperiosteum of the middle ear cleft with the presence of permanent perforation of the tympanic membrane and recurrent or persistent mucoid or mucopurulent otorrhoea.<sup>2,3</sup> The otorhinolaryngologists adopt >12 weeks (3 months) of active otorrhoea for CSOM.<sup>3</sup> But generally, patients with tympanic membrane perforations who continue to discharge mucoid material for periods from 6-12 weeks, despite medical treatment, are recognized as CSOM cases.<sup>2</sup> For urgent referrals from primary health care facilities, and to prevent complications associated with the protraction of the disease, the World Health Organization (WHO) adopted 2 weeks of otorrhoea for the diagnosis of CSOM.<sup>4</sup>

Chronic suppurative otitis media is a global middle ear disease with a serious health-economic burden, especially in developing nations, including Nigeria where the disease prevalence could be very high.<sup>5,6</sup> The highest prevalence of the disease among children in the world ranges from 7% to 46% and is reported among the Inuits of Alaska, Canada and Greenland, American Indians, and Australian Aborigines.

Unfortunately, many local studies examine commonly the bacteriology of CSOM, avoiding mycology; thus, leaving a dearth of knowledge in this regard. *Pseudomonas aeruginosa* and *Staphylococcus aureus* are the most common aerobic microbial isolates in patients with CSOM, followed by *Proteus vulgaris*, *Klebsiella pneumoniae*, and *Diphtheroids*.<sup>2,7</sup> The anaerobes and fungi complete the spectrum of colonizing organisms responsible for this disease.<sup>8</sup> These bacteria and fungi may grow concurrently in a symbiotic relationship, exhibiting an increased virulence of infection.<sup>2,9-11</sup> Previous studies have highlighted CSOM as one of the common otorhinolaryngological diseases among children in

ORL clinics.<sup>12,13</sup> The World Health Organization (WHO) estimates that each year 51,000 children aged below 5 years die from complications of CSOM in developing countries.<sup>4</sup> Thus, the burden in our environment of repeated episodes of childhood CSOM, and the resultant difficulty in the management of these patients, is the rationale for this study.

This study is intended to prospectively analyze the aspirated middle ear exudates of paediatric patients with CSOM for both aerobic bacteria and fungi, with their antimicrobial sensitivity patterns. This knowledge will guide clinicians in empirical prescriptions for a more scientifically based treatment plan for CSOM.

#### **Materials and methods:**

**Study design:** This was a descriptive, cross-sectional hospital-based prospective study conducted among children aged < 18 years that attended a tertiary hospital in North-western Nigeria with a clinical diagnosis of CSOM between June 2016 and June 2018. The criteria for establishing the diagnosis of CSOM was the presence of a tympanic membrane perforation with active otorrhoea lasting for  $\geq 2$  weeks.<sup>4,14</sup>

**Study sampling:** Using consecutive sampling techniques, a total of 166 paediatric patients clinically diagnosed with CSOM, who had not received any antimicrobials within the preceding two weeks, were recruited.

**Study procedure:** The ears were first inspected, and those with dry crust or discharge on the concha were cleaned and sterilized with gauze soaked in 70% alcohol to remove contaminants. Gentle suction toileting or dry mopping was performed to clear the external auditory canal of any discharge. Otomicroscopy was done using an operating microscope with a 250 mm focal length to view and locate perforation on the tympanic membrane, and aid aspiration of the middle ear exudate using sterile disposable plastic Pasteur pipettes. The exudates were expressed onto well-labeled sterile swab sticks (in Amies transport media)

enclosed in airtight plastic tubing, and sent for microbiologic analysis within 30 minutes to 2 hours of collection. Bilaterally discharging ears had their samples collected separately.

***Isolation of Bacteria:*** The middle ear discharges were inoculated on well-labeled and dried Chocolate, MacConkey, and 5% Sheep Blood agar plates using a sterilized inoculated wire loop, which was also used to streak the smear. The agar plates were incubated at 37<sup>0</sup>c for 18-72 hours aerobically; except for the Chocolate agar that was incubated in a candle jar carbondioxide enriched atmosphere. Gram's staining was carried out to ascertain the morphology and Gram's reaction behavior of the isolates after air drying and heat fixing the specimen. Visual examination was done for bacterial growth after 18-72 hours. Plates with no growth were discarded, and the sample was labeled sterile for bacteria. But where there were growths, discrete colonies were selected for morphology. Thereafter, various biochemical tests such as catalase and coagulase tests for Gram positive bacteria, while for Gram negative bacteria motility testing, oxidase, indole, citrate, urease and kligler iron agar screening were carried out to identify the different species as appropriate. Interpretation was done according to standard bacteriological methods to identify pathogens.<sup>15</sup>

***Antibiotic Susceptibility Testing:*** Antibiotic disc susceptibility testing of commonly used antibiotics was carried out using the Kirby Bauer disk diffusion method. The diameter of the zone of inhibition was measured and interpreted as resistant or susceptible as described in Clinical and Laboratory Standards Institute (CLSI) guidelines.<sup>16</sup> The commercially available Oxoid, United Kingdom antibiotic disc comprising of Gentamicin (10µg), Chloramphenicol (30µg), Amoxicillin-Clavulanic acid (30µg), Erythromycin (15µg), Cotrimoxazole (25µg), Ciprofloxacin (5µg), Ceftriaxone (30 µg) and Amoxicillin (10µg).

***Isolation of Fungi:*** The swab sticks were streaked directly on the well-labeled Sabouraud's Dextrose Agar (SDA) plates and incubated at room temperature aerobically for 3-7 days. Following proper masking and gloving, the growths were isolated, identified and

characterized based on their morphological features and cultural characteristics, and a microscopic examination using the lactophenol blue staining technique.<sup>17</sup> Germ tube test was used to differentiate *Candida albicans* from other species of *Candida*.

***Antifungal Susceptibility Testing:*** The antifungal susceptibility test was performed using common antifungal agents, according to Clinical and Laboratory Standards Institute (CLSI) guidelines, and interpreted accordingly.<sup>16</sup> The commercially available Oxoid, United Kingdom antifungal disc comprising Clotrimazole (10µg), Nystatin (100U), Fluconazole (25µg), Griseofulvin (2µg), Ketoconazole (10µg), and Terbinafine (10µg).

***Ethical Considerations:*** Institutional ethical approval was obtained and informed consent was taken from the parents/guardians of the children before enrolling in the study.

***Data Management:*** Data were analyzed using the Statistical Package for Social Sciences (SPSS) version 23 and presented in simple tables.

## **Results:**

A total of 166 paediatric patients were studied, of which 219 middle ear exudates were collected from only the right ear [69 patients (41.6%)], only the left ear [44 patients (26.5%)], or both ears [53 patients (31.9%)]. The ratio of unilateral to bilateral ear discharges was 2.1:1. Patients' ages ranged from 9 months to 17 years with a mean age of 5.5 years. Those aged 1-5 years constituted 42.8%, while 4.2% comprised less than 1 year. Males accounted for 61.5% of the patients, with a male-to-female ratio of 1.6:1. About 15.1% of the patients, were yet to be enrolled in school, while 49.4% of them were in primary schools. Most of the patients (73.0%) were brought to the hospital by their mothers, whereas only 20.7% were brought by their fathers. The commonest presenting symptoms were otalgia (46.4%) and itchy ears (35.5%). Only 7.8% of the patients complained of fever.

Table 1: Sociodemographic characteristics of patients

<b>SOCIO-DEMOGRAPHIC DATA</b>	<b>FREQUENCY (%)</b>
<b>Age group (years) n=166:</b>	
< 1	7 (4.2)
1 – 5	71 (42.8)
6 – 10	43 (25.9)
11 – 15	33 (19.9)
> 15	12 (7.2)
<b>Sex (n=166):</b>	
Male	102 (61.5)
Female	64 (38.6)
<b>Patient's educational level (n=166):</b>	
Yet to be enrolled in school	25 (15.1)
Daycare	0 (0.0)
Nursery	23 (13.9)
Primary	82 (49.4)
Secondary	36 (21.7)




Figure 1: Caregivers that accompanied the child to the hospital.




Figure 2: Presenting symptoms

Out of the 219 aspirates collected, 17 (7.8%) were sterile culturing neither bacteria nor fungi. The remaining 202 (92.2%) cultured different species of aerobic bacteria and fungi. The ratio of aerobic bacteria: fungi was 2.1:1, while the ratio of gram-positive: gram-negative aerobic bacteria was 1: 1.7. Staphylococcus species (27.3%) was the most isolated gram-positive aerobe, followed by Streptococcus species (7.9%), while Klebsiella species (19.7%), followed by Pseudomonas aeruginosa (14.9%) were the most isolated gram-negative aerobe. Aspergillus species (49.0%) and Candida albicans (21.8%) were the commonest fungal isolates.

Gentamycin (91.1%), ceftriaxone (90.1%), and ciprofloxacin (85.6%) showed the highest sensitivity to the isolated aerobes, while cotrimoxazole (10.4%) and amoxicillin (5.5%) were the least; while the isolated fungi were completely susceptible to all the antifungals tested.

Table 2: Microbial isolates

MICROBES	FREQUENCY (%) (N=202)	PREVALENCE (N=219)
<b>A). Aerobic bacteria</b>		92.2
<b>Gram-positive</b>		
1. <i>Staphylococcus aureus</i>	41 (20.3)	
2. <i>Staphylococcus epidermidis</i>	11 (5.5)	
3. <i>Staphylococcus lugdunensis</i>	3 (1.5)	34.7
4. <i>Streptococcus pneumonia</i>	9 (4.5)	
5. <i>Streptococcus pyogenes</i>	7 (3.5)	
6. <i>Corynebacterium diphtheria</i>	5 (2.5)	
<b>Gram-negative</b>		
1. <i>Moraxella catarrhalis</i>	3 (1.5)	
2. <i>Pseudomonas aeruginosa</i>	30 (14.9)	
3. <i>Proteus mirabilis</i>	21 (10.4)	
4. <i>Proteus vulgaris</i>	1 (0.5)	
5. <i>Escherichia coli</i>	17 (8.4)	57.5
6. <i>Enterobacter</i> species	13 (6.4)	
7. <i>Citrobacter</i> species	1 (0.5)	
8. <i>Klebsiella pneumonia</i>	23 (11.4)	
9. <i>Klebsiella oxytoca</i>	17 (8.4)	
<b>B). Fungal isolates</b>		
1. <i>Aspergillus niger</i>	36 (17.8)	
2. <i>Aspergillus fumigatus</i>	11 (5.5)	
3. <i>Candida albicans</i>	21 (10.4)	
4. <i>Mucor</i> species	12 (5.9)	43.8
5. <i>Penicillium</i> species	3 (1.5)	
6. <i>Cephalosporium</i> species	6 (3.0)	
7. <i>Trichosporon</i> species	7 (3.5)	
<b>C). Sterile</b>	-	7.8

Table 3: *In vitro* antibiotic sensitivity pattern of the isolated aerobes

S/No.	Isolated aerobes (n)	<i>In vitro</i> antibiotic sensitivity pattern (%)							
		Gent (10µg)	Chlr (30µg)	Amx-Clav (30µg)	Eryth (15µg)	Cotr (25µg)	Cipro (5µg)	Ceftr (30µg)	Amx (10µg)
1.	<i>Staphylococcus aureus</i> (41)	39 (95.1)	29 (70.7)	21 (51.2)	12 (29.2)	4 (9.8)	30 (73.1)	35 (85.4)	3 (7.3)
2.	<i>Staphylococcus epidermidis</i> (11)	11 (100.0)	-	11 (100.0)	-	-	-	11 (100.0)	-
3.	<i>Staphylococcus lugdunensis</i> (3)	-	3 (100.0)	3 (100.0)	3 (100.0)	-	-	-	-
4.	<i>Streptococcus pneumoniae</i> (9)	5 (55.6)	3 (33.3)	7 (77.8)	5 (55.6)	3 (33.3)	7 (77.8)	9 (100.0)	2 (22.2)
5.	<i>Streptococcus pyogenes</i> (7)	5 (71.4)	3 (42.9)	7 (100.0)	5 (71.4)	1 (14.3)	6 (85.7)	7 (100.0)	-
6.	<i>Corynebacterium diphtheria</i> (5)	4 (80.0)	1 (20.0)	1 (20.0)	1 (20.0)	1 (20.0)	5 (100.0)	3 (60.0)	-
7.	<i>Moraxella catarrhalis</i> (3)	3 (100.0)	1 (33.3)	1 (33.3)	-	-	3 (100.0)	3 (100.0)	-
8.	<i>Pseudomonas aeruginosa</i> (30)	30 (100.0)	12 (40.0)	19 (63.3)	11 (36.7)	7 (23.3)	30 (100.0)	26 (86.7)	2 (6.7)
9.	<i>Proteus mirabilis</i> (21)	21 (100.0)	13 (61.9)	17 (81.0)	5 (23.8)	2 (9.5)	21 (100.0)	18 (85.7)	-
10.	<i>Proteus vulgaris</i> (1)	1 (100.0)	-	1 (100.0)	-	-	1 (100.0)	-	-
11.	<i>Escherichia coli</i> (17)	17 (100.0)	6 (35.3)	12 (70.6)	7 (41.2)	2 (11.8)	16 (94.1)	17 (100.0)	2 (11.8)
12.	<i>Enterobacter</i> species (13)	12 (92.3)	7 (53.9)	9 (69.2)	4 (30.8)	-	13 (100.0)	13 (100.0)	-
13.	<i>Citrobacter</i> species (1)	-	-	1 (100.0)	-	-	1 (100.0)	-	-
14.	<i>Klebsiella pneumoniae</i> (23)	21 (91.3)	7 (30.4)	16 (69.6)	8 (34.8)	1 (4.4)	23 (100.0)	23 (100.0)	1 (4.4)
15.	<i>Klebsiella oxytoca</i> (17)	15 (88.2)	4 (23.5)	12 (70.6)	3 (17.7)	-	17 (100.0)	17 (100.0)	1 (5.9)
	Total (202)	184 (91.1)	89 (44.1)	138 (68.3)	64 (31.7)	21 (10.4)	173 (85.6)	182 (90.1)	11 (5.5)

KEY: Gent – Gentamycin, Chlr – Chloramphenicol, Amx-Clav – Amoxicillin clavulanic acid, Eryth – Erythromycin, Cotr – Cotrimoxazole, Cipro – Ciprofloxacin, Ceftr – Ceftriaxone, Amx – Amoxicillin

Table 4: *In vitro* antifungal sensitivity pattern of the isolated fungi

S/No.	Isolated fungi (n)	<i>In vitro</i> antifungal sensitivity pattern (%)					
		Clot (10µg)	Nyst (100U)	Fluc (25µg)	Gsf (2µg)	Keto (10µg)	Terb (10µg)
1.	<i>Aspergillus niger</i> (36)	36 (100.0)	36 (100.0)	36 (100.0)	36 (100.0)	36 (100.0)	36 (100.0)
2.	<i>Aspergillus fumigatus</i> (11)	11 (100.0)	11 (100.0)	11 (100.0)	11 (100.0)	11 (100.0)	11 (100.0)
3.	<i>Candida albicans</i> (21)	21 (100.0)	21 (100.0)	21 (100.0)	3 (100.0)	21 (100.0)	21 (100.0)
4.	<i>Mucor</i> species (12)	12 (100.0)	12 (100.0)	12 (100.0)	12 (100.0)	12 (100.0)	12 (100.0)
5.	<i>Penicillium</i> species (3)	3 (100.0)	3 (100.0)	3 (100.0)	3 (100.0)	3 (100.0)	3 (100.0)
6.	<i>Cephalosporium</i> species (6)	6 (100.0)	6 (100.0)	6 (100.0)	6 (100.0)	6 (100.0)	6 (100.0)
7.	<i>Trichosporon</i> species (7)	7 (100.0)	7 (100.0)	7 (100.0)	7 (100.0)	7 (100.0)	7 (100.0)
	Total (96)	96 (100.0)	96 (100.0)	96 (100.0)	96 (100.0)	96 (100.0)	96 (100.0)

KEY: Clot - Clotrimazole, Nys - Nystatin, Flu - Fluconazole, Gsf - Griseofulvin, Ket - Ketoconazole, Ter - Terbinafine

## DISCUSSION

Chronic suppurative otitis media is a common problem confronted not only by the Otorhinolaryngologist but also by General medicine practitioners, Family physicians, and Paediatricians.<sup>18</sup> Despite advances in public health and medical care, CSOM is still prevalent around the world. The disease can have irreversible untoward complications if poorly or lately treated, and thus an early bacteriological and mycological diagnosis with prompt treatment using appropriate antimicrobials is essential in combating it. Unfortunately, the susceptibility of CSOM-causing microbes to antimicrobials changes considerably from time to time and is worsened by the misuse of antibiotics. This tends to create multidrug resistance among the microbes, thereby making management even more difficult.<sup>19</sup>

CSOM was commonest among children aged 1 – 5 years (42.8%) in our study. These findings were similar to those reported in other studies, emphasizing CSOM as a childhood disease.<sup>11,13</sup> This study revealed the prevalence of unilateral over bilateral ear discharges in a ratio of 2.1: 1. Similarly, other studies in the south-western and south-eastern regions of Nigeria observed the prevalence of unilateral ear discharge over the bilateral.<sup>11,20</sup> This is important in the management of patients because there is a strong association between bilateral infection and disease severity.<sup>21</sup> It is only reasonable to consider bilateral ear involvement as a more severe disease that requires more attentive and specific management protocol, than unilateral. Outside otorrhoea which was present in all the patients, otalgia (46.4%), itchy ears (35.5%), rhinorrhea (29.5%), and hearing loss (26.5%) were the most common presenting symptoms. Haider and Iseh *et al* noted otalgia, itching, and hearing loss as the commonest presenting symptoms in CSOM.<sup>13,22</sup>

Samples in this study were processed to recover aerobic bacteria and fungi. With regard to the morphotypes, this study yielded more mono-aerobic bacterial cultures (48.4%).

Poly-microbial cultures involving aerobes and fungi accounted for 43.8%; while only 7.8% of the cultures were sterile with neither bacterial nor fungal growths. In India, Yousuf *et al* reported 12.0% sterile cultures, 62.8% mono-microbial, and 25.2% poly-microbial cultures, while Pokharnikar *et al* reported 11.67% of sterile cultures and only 2.5% of poly-microbial culture.<sup>23,24</sup> These dissimilarities in the results may be attributed to differences in the patient population studied and geographical variations. The role of prior empirical antibiotics in sterile cultures of actively discharging CSOM has been reiterated; invariably making treatment of CSOM.<sup>11</sup>

A trend towards Gram-negative aerobes (Gram-positive to Gram-negative ratio of 1:1.7) was noted. This suggests that CSOM infection in our study is more Gram-negative than Gram-positive aerobes. Yousuf *et al* highlighted a Gram-positive to Gram-negative ratio of 1.08:1.<sup>23</sup> Our study revealed that active CSOM infection in the Sokoto children population was mainly caused by the following aerobes: *Staphylococcus aureus* 41/202 (20.3%), *Pseudomonas aeruginosa* 30/202 (14.9%), *Klebsiella pneumonia* 23/202 (11.4%), *Proteus mirabilis* 21/202 (10.4%), *Escherichia coli* 17/202 (8.4%) and *Klebsiella oxytoca* 17/202 (8.4%). Similarly, *Pseudomonas aeruginosa*, *Staphylococcus aureus*, *Proteus* species, *Klebsiella* species, and *Diphtheroids* are the most reported aerobic bacteria isolates in CSOM globally, even by the World Health Organisation.<sup>2,4,24</sup> Shrivastava *et al* in their study, isolated a comparable number of *Staphylococcus aureus* (29.24%) and *Pseudomonas aeruginosa* (28.30%) as the first and second most isolated aerobic bacteria.<sup>25</sup> Isolation of coliforms bacteria such as *Klebsiella* species and *Escherichia coli* in this study was similar to those in India.<sup>23,24</sup> An earlier study in this same center had cultured an equal number of *Pseudomonas aeruginosa* and *Staphylococcus aureus* as the main isolates with very few *Proteus* species.<sup>13</sup> Interestingly, a study in South Africa and south-western Nigeria, cultured *Haemophilus influenzae* as an isolate in CSOM.<sup>26,27</sup> Many have argued that the bacteria isolated from

chronically discharging middle ears are secondary invaders of mucosa that is inflamed due to other factors rather than they are the primary cause of the disease because in CSOM the middle ear environment is thought to be more tolerant to unusual organisms.<sup>3</sup> This view is strengthened by the fact that *Pseudomonas* and *Proteus* which are common inhabitants of even healthy external auditory canals were isolated in this study. *Pseudomonas aeruginosa* is a unique microbe because it survives competition with other pathogens due to its minimal nutritional requirements and its arsenal of antibacterials like pyocyanin and bacteriocin.<sup>25</sup> In local infections like CSOM, the organism can create a secure environment for itself through the necrotizing activities of its extracellular enzymes.<sup>28</sup> A combination of the physical characteristics of the niche, a damaged epithelium, interrupted circulation, devitalized tissue and opportunistic nature protects it from normal host defense mechanisms, and antibiotic agents, causing it to flourish in the external auditory canal. *Staphylococcus aureus* is ubiquitous with highly resistant strains; thus a likely reason for its high prevalence in our study.<sup>9</sup> The isolation of fecal bacteria like *Escherichia coli* and *Klebsiella* species, and water bacteria like *Pseudomonas aeruginosa* may indicate that children in this study area are at risk of infection due to poor hygiene conditions; a reflection of their low socio-economic status.

The prevalence of fungi in our study was 43.8%. Ibekwe *et al* and Prakash *et al* had 25.0% and 12.25% respectively of fungal isolates in their studies.<sup>8,29</sup> Interestingly, opportunistic fungal infections have gained greater importance in human medicine, possibly because of the increasing number of immunocompromised patients; even though they can also produce infection in immunocompetent hosts.<sup>30-32</sup> The fungi isolated from actively discharging CSOM of the Sokoto children population included *Aspergillus niger* (17.8%), *Candida albicans* (10.4%), and *Aspergillus fumigatus* (5.5%). These findings further reiterate the role of fungal infections in CSOM. Some have postulated that the prolonged use of topical broad-spectrum antibiotics may lead to the suppression of bacterial flora and the subsequent

emergence of opportunistic fungal flora in the oral cavity, gastrointestinal tract, vaginal tract, and even in the middle ear.<sup>9,33</sup> This occurs following the entry of fungal spores from the external environment into the moist and alkaline medium of middle ear discharge, which finally leads to the development of mycotic otitis media causing intractable otorrhoea.<sup>33</sup> In CSOM, the development of otalgia, itching, and the presence of hyphae indicate the presence of fungi.<sup>8</sup>

Overall, the aerobes isolated from the actively discharging CSOM of the Sokoto children population were sensitive to gentamycin (91.1%), ceftriaxone (90.1%), and ciprofloxacin (85.6%). Earlier studies in Nigeria, Iran, and India where similar organisms were isolated, reported high *in vitro* sensitivity to aminoglycoside and fluoroquinolones.<sup>11,13,29,34</sup> Regarding individual aerobe susceptibility to antimicrobials, *Pseudomonas* had all isolates completely sensitive to aminoglycoside and fluoroquinolone, with 86.7% and 63.3% sensitivity to cephalosporin and amoxicillin-clavulanic acid respectively. A study in India by Srivastava et al also reported a similar antibiotic sensitivity pattern with *Pseudomonas*.<sup>35</sup> *Klebsiella species* in our study displayed complete sensitivity to fluoroquinolone and cephalosporine, with 90.0% and 70.0% sensitivity to aminoglycoside and amoxicillin-clavulanic acid. *Staphylococcus* did not show complete sensitivity to any antibiotic. Even amoxicillin-clavulanic acid, which enjoys high patronage in this study area, being the most often prescribed empirical antibiotic in otitis media did not express excellent *in vitro* sensitivity when compared with gentamycin, ceftriaxone, and ciprofloxacin. This goes to illustrate that variations in the degree of susceptibility of microbes to antimicrobials occur, and emphasizes the need for local studies in this respect to guide the rational use of the existing antibiotics. Some isolates displayed multidrug resistance which reflects the degree of misuse of antibiotics in this study area. Resistance of pathogenic organisms to over-the-counter antibiotics from recurrent use has become a worldwide problem with serious

consequences for the treatment of diseases including CSOM.<sup>36</sup> In this study, many aerobes showed some form of resistance to the antibiotics tested, especially amoxicillin and cotrimoxazole, further buttressing the multidrug resistance phenomenon. The anaerobes displayed minimal multidrug resistance when compared with the aerobes.

Beta-lactams (penicillins, cephalosporins, etc) are by far the most widely used and efficacious of all antibiotics.<sup>9</sup> Over the past few decades, widespread resistance because of the production of beta-lactamase enzyme has evolved among the most common pathogens to these drugs. The beta-lactamase-producing bacteria isolated in this study were *Staphylococcus*, *Proteus*, *Pseudomonas aeruginosa*, *Streptococcus*, *Klebsiella*, *Citrobacter*, and *Escherichia coli*. These isolates expressed either partial or complete resistance to amoxicillin or ceftriaxone, both of which are beta-lactams. Even though clavulanic acid is a weak broad-spectrum antibiotic, its addition at low concentrations to these beta-lactamase-labile penicillins has demonstrated marked improvement in their activity. Thus, the concept behind amoxicillin-clavulanic acid. These beta-lactamase-producing bacteria, though demonstrated better susceptibility to amoxicillin-clavulanic acid than to only amoxicillin in this study, only a few of the regular bacteria isolated in this study showed complete (100.0%) *in vitro* sensitivity to the drug. This finding may support the concluding statements of Reading *et al*, who noted that the improvement of the activity of beta-lactams by clavulanic acid is very marked for strains of bacteria that owe their resistance to the production of beta-lactamase, but only occurs where this enzyme is readily inhibited by the acid.<sup>37</sup> From this statement, therefore, it is possible that the clavulanic acid in the amoxicillin-clavulanic acid does not readily inhibit all the beta-lactamases produced by the different bacteria strains isolated in this study area, which likely explains this pattern of sensitivity to amoxicillin-clavulanic acid.

With regards the antifungals studied, the isolated fungi showed complete *in vitro* sensitivity to clotrimazole, nystatin, fluconazole, griseofulvin, ketoconazole, and terbinafine. This is unlike the studies in Rajasthan and Uyo, where their isolated fungi showed resistance to Fluconazole and Ketoconazole respectively.<sup>38,39</sup> The most likely explanation for this complete *in vitro* antifungal sensitivity, is that these antifungals are not routinely prescribed in our study center. Thus, the reason for absence of resistant fungal strains. Topical ketoconazole is preferred because of its hepatotoxic effects, while its less toxic synthetic and generally more effective triazole antifungal (fluconazole) is preferred for systemic use. Terbinafine can be used both systemic and topically. However, the cautious use of these antifungals ototopically have been emphasized because of their ototoxic potential.<sup>40</sup>

..

### **Conclusion:**

Aerobic bacteria and fungi were identified in this study to cause CSOM. The main isolates were Gram-negative aerobic bacteria. The bacteria *in vitro* antibiotic sensitivity patterns from this study were gentamycin, ceftriaxone, and ciprofloxacin all exhibiting good gram-negative coverage. The isolated fungi were completely susceptible to all the *in vitro* antifungals tested. This brings to the fore, the role of antimicrobials in the management of CSOM. As first-line drugs in children with CSOM, systemic cephalosporins, ad-hoc ototopic constitution of ceftriaxone, or ototopic fluoroquinolones may be used. In cases of recalcitrant suppuration, mycotic otorrhoea should be suspected and daily aural wick dressing with any of the antifungals studied can be added to the treatment plan.

### **References**

1. Monasta L, Ronfani L, Marchetti F, Montico M, Brumatti L, Bavcar A, et al. Burden of disease caused by otitis media: Systematic review and global estimates. Vol. 7, PLoS ONE. Public Library of Science; 2012. p. e36226.
2. Mittal R, Lisi C V., Gerring R, Mittal J, Mathee K, Narasimhan G, et al. Current concepts in the pathogenesis and treatment of chronic suppurative otitis media. *Journal of Medical Microbiology*. 2015.
3. Browning G, Merchant S, Kelly G, Swan I, Canter R, McKerrow S. Chronic Otitis Media. In: Gleeson M, Browning G, Bruton M, (eds) et al, editors. *Scott-Brown's Otorhinolaryngology Head and Neck Surgery*. 7th ed. Edward Arnold (Publisher) Ltd; 2008. p. 3410.
4. World Health Organisation. Chronic suppurative otitis media - Burden of Illness and Management Options. WHO Library Cataloguing-in-Publication Data. 2004.
5. Lasisi AO, Sulaiman OA, Afolabi OA. Socio-economic status and hearing loss in chronic suppurative otitis media in Nigeria. *Ann Trop Paediatr*. 2007 Dec;27(4):291–6.
6. Ibekwe T, Nwaorgu O. Classification and management challenges of otitis media in a resource-poor country. *Niger J Clin Pract*. 2011 Jul;14(3):262–9.
7. Hamilton J. Chronic Otitis Media in Childhood. In: Gleeson M, Jones N, Clarks R, (eds.) et al, editors. *Scott-Brown's Otolaryngology, Head and Neck Surgery*. 7th ed. London: Hodder Arnold Publishers; 2008. p. 928–64.
8. Ibekwe AO, Shareef ZAL, Benayam A. Anaerobes and fungi in chronic suppurative otitis media. *Annals of Otology, Rhinology and Laryngology*. 1997 Jun 28;106(8):649–52.
9. Qureishi A, Lee Y, Belfield K, Birchall JP, Daniel M. Update on otitis media - Prevention and treatment. Vol. 7, *Infection and Drug Resistance*. Dove Press; 2014. p. 15–24.
10. Nnebe-Agumadu U, Okike O, Orji I, Ibekwe RC. Childhood suppurative otitis media in Abakaliki: Isolated microbes and in vitro antibiotic sensitivity pattern. *Niger J Clin Pract*. 2011 Apr;14(2):159–62.
11. Orji F, Dike B. Observations on the current bacteriological profile of chronic suppurative otitis media in South Eastern Nigeria. *Ann Med Health Sci Res*. 2015;
12. Iseh KR, Adegbite T. Pattern and bacteriology of acute suppurative otitis media in Sokoto, Nigeria. *Ann Afr Med*. 2004;3(4):164–6.
13. Iseh K, Abubakar T. Chronic Suppurative Otitis Media: A Clinical Profile in Sokoto, Nigeria. *Sahel Medical Journal*. 2004;
14. Adoga AS, Maan EN, Malu D, Badung BP, Obiesie I V., Nwaorgu OGB. Swab and aspiration specimen collection methods and antibiogram in chronic suppurative otitis media at Jos University Teaching Hospital: Which is superior. *Ann Afr Med*. 2010;9(4):230–4.
15. Barrow GI, Feltham RKA, editors. *Cowan and Steel's Manual for the Identification of Medical Bacteria*. 3rd ed. Cambridge University Press. Cambridge; 1993.

16. CLSI. Performance Standards for Antimicrobial Susceptibility Testing. CLSI supplement M100. Performance standards for antimicrobial susceptibility testing. 30th ed. CLSI supplement M100. Wayne, PA: Clinical and Laboratory Standards Institute. 2020. p. 332.
17. Collins CH, Lyne PM, Grange IM, Falkinham III JO, editors. Collins and Lyne's Microbiological Methods. 8th ed. London: Arnold; 2004.
18. Iqbal K, Khan M, Satti L. Microbiology of chronic otitis media: experience at Dera Ismail Khan. Gomal J Med Sci. 2011;9(2):189–93.
19. Lee SK, Park DC, Kim MG, Boo SH, Choi YJ, Byun JY, et al. Rate of Isolation and Trends of Antimicrobial Resistance of Multidrug Resistant Pseudomonas Aeruginosa from Otorrhea in Chronic Suppurative Otitis Media. Clin Exp Otorhinolaryngol. 2012 Mar;5(1):17.
20. Akinpelu O V., Amusa YB, Komolafe EO, Adeolu AA, Oladele AO, Ameye SA. Challenges in management of chronic suppurative otitis media in a developing country. J Laryngol Otol. 2008 Jan;122(1):16–20.
21. Nwokoye NN, Egwari LO, Coker AO, Olubi OO, Ugoji EO, Nwachukwu SCU. Predisposing and bacteriological features of otitis media. Afr J Microbiol Res. 2012;6(3):520–5.
22. Haider A. Chronic Suppurative Otitis Media (CSOM): Bacteriological Study. The ORION medical J. 2002;13:13–4.
23. Yousuf A, Malik M, Shamas JU, Beigh Z, Kumari S, Pampori PRA. Bacteriological profile of ear discharge and their antibiotic sensitivity In chronic suppurative otitis media in Kashmir,India. Bangladesh Journal of Medical Science. 2012;11(3):212–6.
24. Pokharnikar S, Souza D'. Bacteriological study of chronic suppurative otitis media (CSOM) amongst adult patients. MedPulse International Journal of Microbiology. 2019;12(1):14–8.
25. Shrivastava A, Singh R, Varshney S, Gupta P, Bist S, Bhagat S et al. Microbiological Evaluation of an Active Tubotympanic Type of Chronic Suppurative Otitis Media. Nepalese J of ENT Head and Neck Surg. 2010;1(2):14–5.
26. Okesola A, Fasina O. Trends in the Resistance Pattern of Bacterial Pathogens of Otitis Media in Ibadan, Nigeria. African Journal of Clinical and Experimental Microbiology. 2011;13(1):46–50.
27. Tiedt NJ, Butler IRT, Hallbauer UM, Atkins MD, Elliott E, Pieters M, et al. Paediatric chronic suppurative otitis media in the Free State Province: Clinical and audiological features. South African Medical Journal. 2013;
28. Vartiainen E, Vartiainen J. Effect of aerobic bacteriology on the clinical presentation and treatment results of chronic suppurative otitis media. J Laryngol Otol. 1996;110(4):315–8.
29. Prakash R, Juyal D, Negi V, Pal S, Adekhandi S, Sharma M, et al. Microbiology of chronic suppurative otitis media in a tertiary care setup of Uttarakhand State, India. N Am J Med Sci. 2013;

30. Carney A. Otitis externa and Otomycosis. In: Gleeson M, Jones N, Clarke R, (eds) et al, editors. *Scott-Brown's Otolaryngology, Head and Neck Surgery*. 7th edn. London: Hodder Arnold Publishers; 2008. p. 3351–7.
31. Kumar H, Seth S. Bacterial and Fungal Study of Chronic Suppurative Otitis Media. *Journal of Clinical and Diagnostic Research*. 2011;5(6):1224–7.
32. Viswanath B. Otomycosis in Immunocompetent Patients – Comparative Study and Literature Review. *ENT Journal*. 2012;91(3):114–21.
33. Vaidya K, Madhup SK, Shrestha BL, Gautam A, Tuladha NR. Bacteriological and Mycological profile of Chronic Suppurative Otitis Media among patients visiting Dhulikhel Hospital. *Annals of Clinical Chemistry and Laboratory Medicine*. 2015 Mar 19;1(1):37–41.
34. Ettehad G, Refahi S, Nemmati A, Pirzadeh A, Daryani A. Microbial and antimicrobial susceptibility patterns from patients with chronic otitis media in Ardebil. *Int J Trop Med*. 2006;1:62–5.
35. Srivastava A, Singh R, Varshney S, Gupta P, Bist S, Bhagat S, et al. Microbiological Evaluation of an Active Tubotympanic Type of Chronic Suppurative Otitis Media. *Nepalese Journal of ENT Head and Neck Surgery*. 2010 Jan 1;1(2):14–6.
36. Alekshun MN, Levy SB. Molecular Mechanisms of Antibacterial Multidrug Resistance. *Cell*. 2007.
37. Reading C, Cole M. Clavulanic acid: a beta lactamase inhibiting beta lactam from *Streptomyces clavuligerus*. *Antimicrob Agents Chemother*. 1977;
38. Agrawal R, Pradhan R, Khatri PK, Shah H, Marothi Y. Mycological profile of chronic suppurative otitis media in a tertiary care center in Rajasthan. *IP International Journal of Medical Microbiology and Tropical Diseases* [Internet]. 2021 Feb 15 [cited 2023 Jan 17];2(4):142–4. Available from: <https://www.ijmmt.org/article-details/3415>
39. Akinjogunla O, Eghafona N. Mycological Investigation in Patients with Acute Otitis Media. *Scientific Journal of Microbiology* [Internet]. 2012 [cited 2021 Mar 8];1(1):19–26. Available from: [https://www.researchgate.net/publication/266007301\\_Mycological\\_investigation\\_in\\_patients\\_with\\_acute\\_otitis\\_media](https://www.researchgate.net/publication/266007301_Mycological_investigation_in_patients_with_acute_otitis_media)
40. Sagit M, Somdas MA, Korkmaz F, Akcadag A. The ototoxic effect of intratympanic terbinafine applied in the middle ear of rats. *J Otolaryngol Head Neck Surg* [Internet]. 2013 [cited 2023 Jan 17];42(1). Available from: <https://pubmed.ncbi.nlm.nih.gov/23663536/>