

## Evaluation of Gene Based Markers Against Fusarium wilt Disease in Chickpea (*Cicer Arietinum* L.)

### ABSTRACT

Chickpea is an important legume crop owing to its nutritional worth. Due to increasing population issue it is needed to maintain productivity of chickpea to fulfil the requirements. At present, the main constraints in chickpea productions are the Fusarium wilt disease. Forty genotypes of chickpea (*Cicer arietinum* L.) were screened against Fusarium wilt resistance using ~~6~~ six gene-based markers. Out of which ~~05~~ five STMS markers showed polymorphism and amplified the alleles linked to resistance and susceptibility to Fusarium wilt disease in chickpea genotypes. The highest PIC value was obtained with STMS Marker TR-29 and the least with STMS Marker TR-19. Based on molecular characterization, genotypes RSVTK-2019-203, RSVT PS -2019-214, RSVT PS -2019-213, RSVTK-2019-101, RSVT PS -2019-215, RSVTK-2019-104, RSVT PS -2019-212 and RSVTK-2019-105 were found to be like RVG -203 (check variety) based on the amplification of same allele with these all. The results open a window to use these genotypes for the development of Fusarium wilt resistant chickpea varieties.

**KEYWORDS:** Characterization, Polymorphic Information content, wilt, molecular markers

### INTRODUCTION

Chickpea (*Cicer arietinum* L.) is the 2<sup>nd</sup> most important legume crop [1] after common bean (*Phaseolus vulgaris* L.). Chickpea consists of about 20.8% protein, 5.6% fat, 1.2% fibre, 59.8% carbohydrate, 4.8-5.5% oil, 3% ash, 0.2% calcium and 0.3% phosphorus. Alongside protein, it is also wealthy in fibre, folic acid and minerals like phosphorus, zinc, iron, calcium and magnesium. Chickpeas are a multifunctional grain legume that is widely utilised around the world, particularly as a protein source [2,3].

Chickpea production is predominantly affected by numerous biotic and abiotic stresses. Biotic stresses consist of disease - Fusarium wilt, collar rot, dry root rot, Ascochyta blight *etc.* Among them Fusarium wilt caused by deuteromycetes fungal pathogen *Fusarium oxysporum* f. sp. ciceris (FOC) is one of the widely disbursed diseases of chickpea [4] and cause yield loss up to the level of 100% depending on varietal susceptibility and climatic conditions [5,6].

**Comment [a1]:** Field score also may be included to confirm its resistance. Because only DNA score will not confirm its resistance

**Comment [a2]:** Where?

Developing wilt-resistant chickpea varieties necessitates labour and money intensive field level phenotyping of numerous germplasm and breeding lines against pathogen races. It also takes a lot of effort. Additionally, such phenotyping using sick plots is likely to run into issues like uneven inoculum distribution and the prevalence of other soil-borne fungi [7]. In order to screen a large number of genotypes, characterising wilt resistance using established DNA markers related to wilt resistance genes is the best method.

Molecular breeding involves molecular markers for selection as well as characterization of crop genotypes [8,9,10]. These markers have immense potential to increase the efficiency and precision of traditional plant breeding [11]. Genomic tools in the form of molecular markers have been developed by molecular biology to identify certain DNA variants that can be used to assist crop improvement programmes [12,13,14]. In chickpea various markers have been identified with their linkage to resistance genes responsible to produce resistance against different races of *F. oxysporum* [15,16]. However, it is important to use these markers for screening of different chickpea genotypes to identify source of resistance against Fusarium wilt. Considering this background, a study was performed to screen out the chickpea genotypes against Fusarium wilt disease resistance using gene-based molecular markers.

## MATERIALS AND METHODS

**Plant material:** A total 40 chickpea genotypes (Table 1) were screened using gene-based STMS markers (Table 2) against Fusarium wilt disease at Plant Molecular Biotechnology Laboratory, Department of Plant Molecular Biology and Biotechnology, College of Agriculture, RVSKVV, Gwalior. These genotypes were collected from Jawaharlal Nehru Krishi Vishwa Vidyalaya, Jabalpur, Madhya Pradesh, India, and RAK College of Agriculture, Sehore, RVSKVV, Gwalior, Madhya Pradesh, India.

S. no.	Name of genotype	S. no.	Name of genotype
1	RVSVT PS-2019- 201	21	RVSVTK-2019-106
2	RVSVT PS-2019- 202	22	RVSVTK-2019-107
3	RVSVT PS-2019- 203	23	RVSVTK-2019-108
4	RVSVT PS-2019- 204	24	RVSVTK-2019-109

5	RVSVT PS-2019- 205	25	RVSVTK-2019-110
6	RVSVT PS-2019- 206	26	RVSVTD-2019-1
7	RVSVT PS-2019- 207	27	RVSVTD-2019-2
8	RVSVT PS-2019- 208	28	RVSVTD-2019-3
9	RVSVT PS-2019- 209	29	RVSVTD-2019-4
10	RVSVT PS-2019- 210	30	RVSVTD-2019-5
11	RVSVT PS-2019- 211	31	RVSVTD-2019-6
12	RVSVT PS-2019- 212	32	RVSVTD-2019-7
13	RVSVT PS-2019- 213	33	RVSVTD-2019-8
14	RVSVT PS-2019- 214	34	RVSVTD-2019-9
15	RVSVT PS-2019- 215	35	RVSVTD-2019-10
16	RVSVTK-2019-101	36	RVSVTD-2019-11
17	RVSVTK-2019-102	37	RVSVTD-2019-12
18	RVSVTK-2019-103	38	RVSSG91-13
19	RVSVTK-2019-104	39	RVSSG96-14
20	RVSVTK-2019-105	40	RVG-203

**DNA extraction:** Leaf samples were collected from one month old seedling from experimental field. The collected samples were placed in cooling pads to transfer and then stored at -80 °C deep freezer. High quality genomic DNA was extracted from 8-10 days old young and fresh leaves by employing CTAB method as proposed by Doyle and Doyle [17] with some modifications as suggested by Tiwari *et al.* [18]. Extracted DNA was quantified through electrophoresis on 0.8% agarose gel and compared after loading a known quantity DNA marker ( $\lambda$  DNA) on the same gel as a standard. Apart from it a Spectrophotometer was also used for quantification of DNA.

**Markers analysis:** The polymerase chain reaction was performed in 10 µl reaction mixture comprising of 1X PCR buffer, 0.1 µl *Taq* DNA polymerase, 1 µl dNTP (1 mM), 0.5 µl of primers (10 pM) and 20 ng/µl of genomic DNA in a thermocycler (Bio-Rad, USA). The PCR protocol comprised of initial denaturation step of 94 °C for 3 min followed by 35 cycles of 94 °C for 1 min, annealing cycles (from 52 °C to 57 °C) varied for different markers system for 30 sec, elongation at 72 °C for 1 min with final extension at 72 °C for 7 min. PCR amplified products of STMS markers along with standard markers (100 bp) were separated through electrophoresis on 3% agarose gel respectively at 100 V for two hrs. The agarose gels were stained with Ethidium Bromide (1µg/ml). After electrophoresis the agarose gels were visualized under UV light and photographed under Bio-Rad Gel documentation system.

**Band scoring and data analysis:** The genetic profile of 40 chickpea genotypes was obtained based on differences in allele size using five STMS reported markers (Table 2). The scoring was done using a standard size ladder and a banding pattern. The data sheet was produced based on allele pattern A/A and used for further analysis. The major allele frequency, polymorphism information content (PIC) and genetic distance-based clustering was performed with Unweighted Pair Group Method for Arithmetic average (UPGMA) tree using Power Marker v3.25 software [19].

<b>Table 2</b> Details of primers used for screening of Fusarium wilt in chickpea genotypes				
<b>Primer Name</b>	<b>Category</b>	<b>Primer sequence</b>		<b>Reference</b>
TA-59	STMS	F: ATC TAA AGA GAA ATC AAA ATT GTC GAA	R: GCA AAT GTGAAG CAT GTA TAG ATA AAG	[20]
TA-96	STMS	F: TGT TTT GGA GAA GAG TGA TTC	R: TGT GCA TGC AAA TTC TTA CT	[20]
TR- 19	STMS	F: TCA GTA TCA CGT GTA ATT CGT	R: CAT GAA CAT CAA GTT CTC CA	[20]
TA194	STMS	F:TTTTGGCTTATTAGA CTGAC TT	R:TTGCCATAAAA TACAAAATCC	[20]
TR29	STMS	F:GCCCACTGAAA AATAAAAAG	R:ATTTGAACCTCA AGTTCTCG	[20]

TR31	STMS	F:CTTAATCGCACATTT ACTCTAAA ATCA	R:ATCCATTAACA CGGTTACCTATAA	[20]
------	------	------------------------------------	--------------------------------	------

## RESULTS AND DISCUSSION

Due to the minimal polymorphism in the chickpea genome, the development of gene linked molecular markers has been relatively slow. The markers associated to various wilt resistance genes have been found and mapped. In the past few decades, chickpea breeders have developed an array of varieties that have performed well under field conditions. The last ten years have seen advancements in the study of chickpeas using molecular breeding tools. However, the fusion of traditional and advanced molecular breeding has accelerated study in cereals like wheat and rice. So, similar advancements must be made in the improvement of chickpea crop to meet complete requirements. Utilizing molecular markers for identification of desired genotypes may help in planning of molecular breeding experiment of intergression of targeted gene(s) in desired genotype [21,22]. These approaches may help in deciding gene-pyramid schemes as well [23,24,25,26].

During the present investigation 40 chickpea genotypes were evaluated with Fusarium wilt resistance gene linked markers. All STMS markers *viz.*, TA-59, TA-96, TR-19, TA-194 and TR-29 showed polymorphism in the chickpea genotypes and amplified alleles associated with resistance and susceptibility. The results found in our present study are like with the earlier studies. Sahu *et al.* [27] investigated chickpea genotypes and used gene-based molecular markers to screen them against Fusarium wilt disease. Amadabade *et al.* [28] investigated six chickpea genotypes, each with a distinct Fusarium wilt response, using DNA-based genetic markers associated with disease resistance/susceptibility. Padaliya *et al.* [29] employed seven molecular markers previously linked to disease resistance/susceptibility with six chickpea genotypes. In the current study, a total of 28 alleles were identified with an average of 5.6 alleles per locus for different markers (Table 3). However, Solanki *et al* [13] reported an average of 1.65 alleles per locus while working on diversity analysis in chickpea genotypes using different markers. Previously, Bhardwaj *et al.* [30] reported an average 2.49 alleles per locus during their study on diversity assessment in chickpea genotypes using STMS markers.

Comment [a3]: TA 31 is missing only 5 are listed

Comment [a4]: Repetition of wordss may be avoided

Comment [a5]: Repetition of words in the same meaning may be avoided in the same para

Marker	Major allele frequency	Allele no.	Gene diversity	PIC
TA-59	0.400	6	0.753	0.719
TA-96	0.500	5	0.631	0.569
TR-19	0.525	5	0.625	0.567
TA-194	0.350	5	0.734	0.687
TR-29	0.2750275	7	0.790	0.759
<b>Mean</b>	<b>0.410</b>	<b>5.6</b>	<b>0.707</b>	<b>0.660</b>

Comment [a6]: What about TR 31


The gene diversity arrayed between 0.625 to 0.790 for the markers TR-19 and TR-29 correspondingly with an average of 0.707 and Polymorphic Information Content (PIC) values varied between 0.5666-0.567 to 0.759 for the markers TR-19 and TR-29 respectively with a mean value of 0.660 respectively. ~~The primer which showed highest gene diversity and PIC values was TR29 while the lowest gene diversity and PIC values was detected for the primer TR-19.~~ The major allele frequency ranged between 0.2750-275 (TR29) to 0.525 (TR-19) with a mean worth of 0.410.

The genetic relationships among chickpea genotypes are presented in molecular based UPGMA tree. All the genotypes were grouped into 4 clusters. First cluster had 9 genotypes including RVSVD-2019-1, RVSVD-2019-2, RVSVD-2019-4, RVSVD-2019-5, RVSVD-2019-6, RVSVD-2019-9, RVSVD PS 2019-203, RVSSG 96-14 and RVSVD-2019-10. These genotypes are found to resemble to each other at molecular level. Further, the second cluster also contained 9 genotypes *i.e.*, RVSVDK-2019-103, RVSVD PS-2019-214, RVSVD PS-2019-213, RVSVDK- 2019- 101, RVSVD PS -2019-215, RVSVD PS -2019-104, RVSVD PS-2019-212, RVSVDK- 2019-105 along with check variety for Fusarium wilt resistance RVG-203. Grouping of these chickpea genotypes in the same cluster with the check variety (RVG-203) indicates the presence of similar segment of DNA in them. Due to which the applied markers were able to amplify similar banding pattern with these genotypes. A crucial step in selecting effective sources of high resilience for breeding programmes is the identification of genotypes that have high stability for low disease severity. However, the presence of resistance against Fusarium wilt in chickpea genotypes grouped with the check variety RVG-203 should be

confirmed before their selection as a donor in a breeding programme. The job of testing breeding lines and germplasm for disease resistance is extensive and involves different methods like field trials and laboratory-based screening. However, field level screening has few limitations because of association of difficulties in development and maintenance of uniform sick plot.

Consequently, third cluster had 11 chickpea genotypes including RSVTK-2019-109, RSVTK-2019-108, RSVT PS-2019-205, RSVT PS-2019-210, RSVT PS- 2019-204, RSVTK-2019-102, RSVT PS- 2019-201, RSVT PS- 2019-202, RSVTK-2019-110, RSVTK-2019-106 and RSVTK-2019-107. Forth cluster also contained 11 genotypes namely RSVT PS-2019-208, RSVT PS-2019-209, RSVT PS-2019-207, RSVTD -2019-12, RVSSG 91-13, RSVTD-2019-11, RSVT PS-2019-206, RSVTD-2019-07, RSVTD-2019-08, RSVT PS-2019-211, RVSTVD-2019-03.

UNDER PEER REVIEW



**Fig.1** Dendrogram of 40 Chickpea genotypes showing clusters based on similarity using UPGMA relationship.

The best method for controlling chickpea wilt disease in India and other emerging nations continues to be the use of resistant cultivars. Identification of markers closely linked with Fusarium wilt resistance gene(s) is the prerequisite before performing screening of chickpea genotypes. It is important to combine field as well as laboratory-based methods in order to create linkage maps, find stable QTLs associated with Fusarium wilt resistance, and investigate novel markers in order to accurately select resistant genotypes from segregating populations. It will be easier to introduce resistant genes from Fusarium wilt resistant cultivars carrying targeted genes

into different chickpea genotypes/varieties on availability of closely-related markers for those genes [31].

## CONCLUSION

For validation of marker against Fusarium wilt in chickpea genotypes, high quality DNA was extracted from 40 genotypes and six STMS markers were screened in selected genotypes and five markers produced polymorphism. Highly resistant genotypes including RVSVTK-2019-203, RVSVT PS -2019-214, RVSVT PS -2019-213, RVSVTK-2019-101, RVSVT PS -2019-215, RVSVTK-2019-104, RVSVT PS -2019-212 and RVSVTK-2019-105 were validated at genomic level using STMS markers which could be used in hybridization program for chickpea improvement.

Comment [a7]: Resistance field score may be included to confirm its resistance

## REFERENCES

1. Asati, R., Tripathi, M.K., Tiwari, S., Yadav, R.K. and Tripathi, N. (2022). Molecular breeding and drought tolerance in chickpea. *Life* 12:1846.
2. Sahu, V.K., Tiwari, S., Gupta, N., Tripathi, M.K. and Yasin, M. (2020a). Evaluation of physiological and biochemical contents in desi and Kabuli chickpea. *Legume Research* DOI:10.18805/LR-4265.
3. Gupta, N., Tiwari, S., Tripathi, M.K. and Bhagyawant, S.S. (2021). Antinutritional and protein-based profiling of diverse desi and wild chickpea accessions. *Current Journal of Applied Science and Technology* 40(6):7-18
4. Jimenez-Gasco, M., Navas-Cortes, J. A. and Jimenez-Diaz, R. M., (2010). The *Fusarium oxysporum* f. Sp. *Ciceris* /*Cicer arietinum* pathosystem: a case study of the evolution of plant pathogenic fungi into races and pathotypes. *International Microbiology*. 7(2): 95-104.
5. Patil, M., Om, G., Pawar, M., and Chobe, D. R. (2015). Effect of culture media, pH and temperature on the mycelial growth and sporulation of *Fusarium oxysporum* f. sp. *ciceris* isolates of chickpea from Central Zone of India. *Res. J. Appl. Sci.* 49, 54–60.
6. Yadav, S., Shah, V. and Mod, B. (2019). Genetic diversity analysis between different varieties of chickpea (*Cicer arietinum* L.) using SSR markers. *Int. J. Appl. Sci. Biotechnol.* 7(2): 236-242
7. Babu, D.R. and Ravikumar, R.L. (2013). Characterization of Fusarium Wilt Resistance in Diverse Chickpea (*Cicer arietinum* L.) Genotypes using Established Molecular Markers. *International Journal of Bio-resource and Stress Management*, 4(4):631-634.
8. Shyam, C., Tripathi, MK, Tiwari, S, Tripathi, N. and Ahuja, A. (2020). Molecular characterization and identification of *Brassic* genotype(s) for low and higher glucosinoid content using SSR markers. *Global J Biosci Biotechnol.* 9(2):56-66.
9. Upadhyay, S., Singh, A.K., Tripathi, M.K., Tiwari, S. and Tripathi, N. (2020). Validation of simple sequence repeats markers for charcoal rot and *Rhizoctonia* root rot resistance in soybean genotypes. *I.J.A.B.R.* 10(2):137-144.

10. Verma, R., Tripathi, M. K., Tiwari, S., Pandya, R. K., Tripathi, N. and Parihar, P. (2021). Screening of pearl millet [*Pennisetum glaucum* (L.) R. Br.] genotypes against blast disease on the basis of disease indexing and gene specific SSR markers. *Int J Curr Microbiol Appl Sci.* 10(02):1108-1117.
11. Mandloi, S., Tripathi, M.K., Tiwari, S. and Tripathi, N. (2022). Genetic diversity analysis among late leaf spot and rust resistant and susceptible germplasm in groundnut (*Arachis hypogea* L.). *Israel Journal of Plant Sciences* doi: [https://doi.org/ 10.1163/22238980-bja10058](https://doi.org/10.1163/22238980-bja10058).
12. Tiwari, S., Tripathi, N., Tsuji, K. and Tantwai, K. (2019). Genetic diversity and population structure of Indian soybean (*Glycine max* (L.) Merr.) as revealed by microsatellite markers. *Physiology and Molecular Biology of Plants* 25: 953-964.
13. Solanki, R.S., Babbar, A. and Tripathi, N. (2022) Genetic diversity analysis in *kabuli* chickpea (*Cicer arietinum* L.) genotypes based on quantitative traits and molecular markers. *Bangladesh Journal of Botany* 51(3): 581-587.
14. Tripathi, M.K., Tripathi, N., Tiwari, S. Mishra, N., Sharma, A., Tiwari, S. and Singh, S. 2023. Identification of Indian soybean (*Glycine max* [L.] Merr.) genotypes for drought tolerance and genetic diversity analysis using SSR markers *SCI- 3(3);2023;31-46*
15. Cobos, M.J., Fernandez, M., Rubio, J., Kharrat, M., Moreno, M.T., Gil, J. and Millan, T. (2005). A linkage map of chickpea (*Cicerarietinum* L.) based on populations from Kabuli x Desi crosses: location of genes for resistance to Fusarium wilt race 0. *TheorAppl Genet* 110:1347–1353.
16. Jha, U.C., Bohra, A., Pandey, S. and Parida, S.K. (2020). Breeding, Genetics, and Genomics Approaches for Improving Fusarium Wilt Resistance in Major Grain Legumes. *Front. Genet.* 11:1001. doi: 10.3389/fgene.2020.01001
17. Doyle, J.J. and Doyle, J.L. (1987). A rapid DNA isolation procedure for small quantities of fresh leaf tissue. *Phytochemical Bulletin*, 19: 11-15.
18. Tiwari, S., Tomar, R.S, Tripathi, M.K. and Ahuja, A. (2017). Modified protocol for plant genomic DNA isolation. *Indian Res J Genet & Biotech.* 9(4):478–485
19. Liu, K. and Muse, S.V. (2005). Power marker integrated analysis environment for genetic marker data. *Bioinformatics* 21: 2128-2129.
20. Winter, P., Benko-Iseppon, A.M., Huttel, B., Ratnaparkhe, M., Tullu, A., Sonnante, G., Pfaff, T., Tekeoglu, M., Santra, D. and Sant, V. (2000), A linkage map of the chickpea (*Cicer arietinum* L.) genome based on recombinant inbred Page 14/19 lines from a *C. arietinum* × *C. reticulatum* cross: localization of resistance genes for Fusarium wilt races 4 and 5. *Theor Applied Genet* 101: 1155-1167.
21. Mishra, N., Tripathi, M.K., Tiwari, S., Tripathi, N. and Trivedi, H.K. (2020). Morphological and molecular screening of soybean genotypes against yellow mosaic virus disease. *Legume Research* DOI: 10.18805/LR-4240.
22. Tripathi, N., Tripathi, M.K., Tiwari, S. and Payasi, D.K. (2022). Molecular breeding to overcome biotic stresses in soybean: update. *Plants (Basel).* 11(15):1967.
23. Baghel, R., Sharma, A.K., Tiwari, S., Tripathi, M.K. and Tripathi, N. (2020). Genetic diversity analysis of Indian mustard (*Brassica spp*) germplasm lines using SSR molecular markers. *Int. J. Curr. Microbiol. App. Sci.* 9(12):137-143.
24. Parmar, A., Tripathi, M.K., Tiwari, S., Tripathi, N., Parihar, P. and Pandya, R.K. (2022). Characterization of pearl millet [*Pennisetum glaucum* (L.) R Br.] genotypes against downey mildew disease employing disease indexing and ISSR markers. *Octa J. Biosci.* 10 (2):134-142

25. Rajpoot, P., Tripathi, M.K., Tiwari S., Bimal S. S., Tripathi N., Parihar P. Pandya R. K. and Satyavathi, C.T. (2023). Characterization of pearl millet [*Pennisetum glaucum* (L.) R br.] Genotypes against blast disease employing disease scoring and gene specific SSR markers. *SCI- 3(3)*16-30
26. Bhawar PC, Tiwari S, Tripathi MK, Tomar RS, Sikarwar RS. Screening of groundnut germplasm for foliar fungal diseases and population structure analysis using gene based SSR markers. *Curr J Appl Sci Technol.* 2020;39(2):75-84.
27. Sahu, V.K., Tiwari, S., Tripathi, M.K., Gupta, N., Tomar, R.S. and Yasin, M. (2020b). Morpho-physiological and biochemical traits analysis for Fusarium wilt disease using gene-based markers in desi and Kabuli genotypes of chickpea (*Cicer arietinum* L.). *Indian J. Genet* 80: 16
28. Amadabade, J., Arora, A., Sahu, H. and Singh, M.K., (2016). Molecular characterization of fusarium wilt resistance and confirmation of hybridity in chickpea genotypes. *Ecology, Environment and Conservation.* 22: 47-51.
29. Padaliya, R.V., Suthar, K.P., Singh, D., Mahatma, M.K. and Patil, V.R. (2013). Marker assisted characterization of chickpea genotypes for wilt resistance. *African Journal of Biotechnology*, 12: 6907-6912.
30. Bharadwaj, C., Chauhan, S.K., Rajguru, G., Srivastava, R., Satyavathi, C.T. and Yadav, S. (2010). Diversity analysis of chickpea (*Cicer arietinum* L.) using STMS markers. *Indian Journal Agricultural Science* 80: 947–51.
31. Jha, U., Jha, R., Bohra, A., Manjunatha, L., Saabale, P., Parida, S., Chaturvedi, S.K., Thakro, V. and Singh, N.P. (2021). Association mapping of genomic loci linked with Fusarium wilt resistance (Foc2) in chickpea. *Plant Genetic Resources*, 19(3), 195-202

Formatted: Font: Italic