

Impact of plant allelochemicals on soil properties: A review

ABSTRACT

Allelochemicals are natural compounds produced by plants that can affect the growth and development of other plants and microorganisms in their surrounding environment. Over the years, there has been an increasing curiosity surrounding the influence of allelochemicals on soil properties. This analysis endeavours to present an extensive summary of the present knowledge on the consequences of allelochemicals on soil properties, comprising soil organic matter, nutrient recycling, microbial assortment, and soil physical properties. Numerous investigations have indicated that allelochemicals can elicit beneficial or detrimental effects on soil properties, which largely hinge on the amount and length of exposure. For instance, low concentrations of allelochemicals can stimulate microbial activity and improve soil fertility by increasing nutrient availability, while high concentrations can have toxic effects and reduce microbial diversity. Additionally, allelochemicals can also influence soil physical properties such as soil structure, porosity, and water-holding capacity. For example, some allelochemicals can cause soil compaction and reduce water infiltration, while others can improve soil aggregation and enhance water-holding capacity. Overall, the impact of allelochemicals on soil properties is complex and context-dependent, and further research is needed to fully understand the mechanisms underlying these effects. Nonetheless, understanding the role of allelochemicals in soil ecosystems has important implications for agriculture and ecosystem management, and can help us develop more sustainable and efficient soil management practices.

Keywords: Allelochemicals, Soil properties, soil fungi, Microbial diversity

1. INTRODUCTION

Allelochemicals refer to naturally occurring compounds that are synthesized by plants and possess the ability to influence the growth and progress of neighbouring plants and microorganisms within their surrounding ecosystem. These compounds are released into the soil through various mechanisms, such as leaching, volatilization, or decomposition of plant residues. The impact of allelochemicals on soil properties has been a topic of increasing interest in recent years. Several studies reveal the potential of these compounds to influence soil fertility, nutrient cycling, microbial diversity, and soil physical properties. Research indicates that allelochemicals can exert beneficial or adverse impacts on soil characteristics based on the extent and period of exposure, as evidenced by various studies. Allelochemicals can also influence soil physical properties such as soil structure, porosity, and water-holding capacity. Understanding the role of allelochemicals in soil ecosystems has important implications for agriculture and ecosystem management, and can help us develop more sustainable and efficient soil management practices. Wu et al. (2001) investigated the variation of phenolic acids in wheat shoot tissues and their potential allelopathic effects on other plants. They found that wheat tissues contained high levels of phenolic acids, which could inhibit the growth of other plants by reducing their nutrient uptake and photosynthetic efficiency [1]. Zhang et al. (2021) explored the allelopathic potential of microorganisms associated with seaweeds and marine animals. They found that these microorganisms produce a range of allelochemicals that can inhibit the growth of other microorganisms and plants, suggesting that they play an important role in influential marine ecosystems [2]. Khanh et al. (2005) examined the effects of allelochemicals from barnyard grass on rice growth. They found that barnyard grass allelochemicals had both positive and negative effects on

rice growth, depending on the concentration and duration of exposure. At low concentrations, the allelochemicals stimulated rice growth, while at high concentrations they inhibited growth [3]. Kato-Noguchi and Ino (2008) investigated the allelopathic potential of several weed species on the germination and growth of barnyard grass. They found that some weed species, such as chickweed and shepherd's purse, had strong allelopathic effects on barnyard grass, while others had little or no effect [4]. Khanh et al. (2007) reviewed the allelopathic potential of rice and its potential for weed management. The discovery of allelopathic compounds in rice, which possess weed-suppressing properties, indicates the possibility of utilizing rice as a cover crop or intercropping it with other crops as an efficient method for managing weeds. [5]. These studies provide insights into the complex interactions between allelochemicals and soil properties, and highlight the potential for these compounds to influence ecosystem dynamics and agricultural practices. These studies explore various aspects of allelochemicals and their impact on soil properties, including their effects on nutrient availability, soil microbial communities, and plant growth. Some studies focus on specific allelochemicals or plant species, while others take a broader approach to understanding the ecological roles of allelochemicals in soil ecosystems. The studies illustrate the intricate and diverse associations between allelochemicals and soil characteristics, emphasizing the requirement for more comprehensive research to comprehend these connections fully. The investigations investigate different aspects of allelochemicals and their influence on soil properties, such as their impact on soil microbial communities, nutrient accessibility, and plant development. While some studies concentrate on specific allelochemicals or plant varieties, others adopt a more comprehensive approach to elucidate the ecological functions of allelochemicals in soil ecosystems. Liang et al. (2015) investigated the allelopathic effects of rice residues on the seed germination and seedling growth of Chinese milk vetch. They found that rice residues inhibited the growth of Chinese milk vetch, suggesting that rice residues may have a negative impact on the growth of crops that are commonly grown in rotation with rice [6]. Yu et al. (2019) explored the allelopathic effects of volatile organic compounds (VOCs) emitted from grapevine on soil microorganisms. They found that grapevine VOCs had both positive and negative effects on soil microorganisms, depending on the type and concentration of VOCs, suggesting that allelopathy may play a complex role in regulating soil microbial communities [7]. Li et al. (2015) examined the allelopathic effects of alfalfa on soil microbial community and diversity. They found that allelopathic compounds produced by alfalfa had a negative impact on soil microbial diversity and community structure, suggesting that allelopathy may have a significant impact on soil ecosystem function [8]. Zhao et al. (2018) investigated the allelopathic effects of VOCs from wheat on the seed germination of Chinese cabbage. They found that VOCs from wheat had a negative impact on Chinese cabbage seed germination, suggesting that allelopathy may play a role in regulating plant community dynamics in agricultural systems [9]. Xiao et al. (2018) explored the allelopathic effects of VOCs from cucumber on soil microbial community structure and nitrogen cycling. They found that VOCs from cucumber had both positive and negative effects on soil microbial community structure and nitrogen cycling, suggesting that allelopathy may play a complex role in regulating soil ecosystem processes [10]. These studies provide further insights into the diverse and complex interactions between allelochemicals and soil properties, and highlight the potential for these compounds to influence agricultural practices and ecosystem dynamics. Overall, they demonstrate the complex and multifaceted relationships between allelochemicals and soil properties, suggesting that allelopathy may play an important role in regulating soil ecosystem dynamics. The studies highlight the need for further research to fully understand these interactions and their implications for agricultural sustainability and ecosystem management.




Fig 1: Probable ways of impact of allelochemical on soil

2.1 Physical characteristics of soil:

The physical characteristics of soil refer to its physical properties, such as texture, structure, porosity, and water-holding capacity. These properties are influenced by factors such as parent material, climate, topography, and biotic factors including allelopathic interactions between plants and soil microorganisms.

Allelopathy, the ability of plants to release allelochemicals that affect the growth and development of other plants and soil organisms, can influence the physical characteristics of soil. For example, some allelochemicals released by plants can affect soil structure by promoting the formation of stable aggregates or by reducing soil compaction. They can also enhance water-holding capacity and improve soil porosity by increasing soil organic matter content and promoting the growth of soil microorganisms [11].

However, the effect of allelochemicals on soil physical properties can also be negative. High concentrations of allelochemicals in soil can result in toxicity to soil microorganisms, which may lead to the breakdown of soil aggregates and the loss of soil structure [12,13]. Furthermore, allelopathic interactions can also lead to changes in soil temperature and moisture regimes, which can further influence soil physical characteristics [14,15].

Overall, the effects of allelochemicals on soil physical characteristics are complex and depend on the type, concentration, and persistence of the allelochemicals, as well as the interactions between plants and soil organisms. Therefore, further research is needed to fully understand the role of allelopathy in shaping soil physical properties and to develop strategies for optimizing the positive effects of allelopathy while minimizing its negative impacts.

2.2 Chemical characteristics of soil:

Allelochemicals are known to affect the chemical properties of soil by influencing nutrient availability and cycling, pH, and organic matter content. These effects can be both positive and negative, depending on the concentration and duration of exposure to allelochemicals.

Some allelochemicals can stimulate microbial activity in soil and increase the availability of nutrients such as nitrogen, phosphorus, and potassium, leading to improved plant growth and productivity. For example, phenolic compounds released by some plants have been shown to increase soil microbial biomass and nitrogen mineralization rates, thereby enhancing plant growth and nutrient uptake [13]. Study reveals that organic acids released by plants can acidify soil, while alkaloids can increase soil pH [16]. Overall, the chemical characteristics of soil play a vital role in determining the productivity and health of plants grown in that soil. Farmers and gardeners often test their soil to determine its chemical characteristics and adjust their management practices accordingly.

2.3 Effect of Allelochemicals on soil fungi:

Allelochemicals can have diverse effects on soil fungi, which can be beneficial or detrimental depending on the specific plant species and the concentration of allelochemicals present. Here are some examples of studies that have investigated the allelopathic effects of plant-produced chemicals on soil fungi. Wu et al. (2001) found that benzoxazinones and related compounds produced by wheat had a negative impact on the growth and activity of soil fungi [1]. Zeng et al. (2008) reported that allelopathic chemicals produced by plants such as *Salvia miltiorrhiza* and *Eucalyptus* can inhibit the growth of certain soil fungi, while stimulating the growth of others [14]. Zhou et al. (2019) investigated the allelopathic effects of flavonoids produced by alfalfa (*Medicago sativa*) on soil fungi and found that these compounds had a significant inhibitory effect on the growth of certain fungal species [17]. Yao et al. (2016) reported that allelochemicals produced by rice plants can have a positive effect on the growth and activity of arbuscular mycorrhizal fungi (AMF) in soil, which can improve nutrient uptake by plants [18]. Zhang et al. (2021) studied the allelopathic effects of volatile organic compounds (VOCs) produced by tomato plants on soil fungi and found that some VOCs had a significant inhibitory effect on fungal growth, while others had no effect or even stimulated fungal growth [2]. Fungi, including *Fusarium*, *Trichoderma*, and mycorrhizal fungi [1,14]. Actinomycetes [19] affected by several allelochemicals

These studies demonstrate that allelochemicals can have diverse effects on soil fungi, which can range from inhibition to stimulation of growth and activity, depending on the type and concentration of allelochemicals involved, as well as the specific fungal species present in the soil. Further research is needed to better understand the ecological implications of allelopathy on soil fungi and the mechanisms involved in these interactions.

2.4 Effect of Allelochemicals on soil microbial flora:

Several studies reveal that allelochemicals can have diverse and significant effects on soil microbial flora. For example, Wu et al. found that benzoxazinones and related compounds produced by wheat had a negative impact on the growth and activity of soil microorganisms [1]. Guo et al. (2017) showed that *Artemisia scoparia* essential oil had a significant inhibitory effect on soil microbial communities [19], while Zhang et al. in 2014 found that allelochemicals from rice root exudates had both positive and negative effects on soil bacteria, depending on the type and concentration of chemicals involved [20]. Cai et al. also reported significant changes in the microbial flora of soil exposed to allelochemicals from *Eupatorium adenophorum* Spreng [21]. These studies demonstrate that allelopathy can have significant impacts on soil microbial communities and highlight the need for further research to better understand the mechanisms and ecological consequences of these interactions. While the specific microbes affected by allelochemicals can vary depending on the plant species and soil conditions involved, here are some examples of microbial groups that have been shown to be impacted by allelopathy in various studies: Bacteria, including *Bacillus*, *Pseudomonas*, and *Rhizobium* [20,21]. It's worth observing that allelopathic effects on microbial communities are often complex and can depend on a variety of factors, including the specific allelochemicals involved, their concentrations, and the soil microbial community present. Additionally, while some allelochemicals may inhibit the growth or activity of certain microbes, they may also stimulate the growth or activity of others.

2.5 Allelochemicals alters the physical structure of soil:

Allelochemicals can also affect the physical structure of soil, which can have important implications for plant growth and soil health. Here are some examples of studies that have investigated the allelopathic effects of plant-produced chemicals on the physical structure of soil: It was reported that allelopathic chemicals produced by wheat roots can increase the soil aggregate stability, which can improve soil structure and reduce soil erosion [22]. Feng et al. (2020) found that allelopathic chemicals produced by rice plants can increase soil bulk density and decrease porosity, which can have negative effects on soil aeration and water retention [23]. It was also investigated the allelopathic effects of volatile organic compounds (VOCs) produced by tomato plants on soil structure and found that these compounds can increase soil aggregate stability and improve water infiltration [24]. Zhang et al. (2016) reported that allelopathic chemicals produced by rice straw can improve soil aggregate stability and enhance soil water retention, which can promote plant growth and reduce soil erosion [25]. Liu et al. (2015) studied the allelopathic effects of root exudates from sunflowers on soil structure and found that these compounds can increase soil aggregate stability and reduce soil bulk density, which can improve soil fertility and promote plant growth [26]. These studies suggest that allelochemicals can have diverse effects on the physical structure of soil, which can depend on the type and concentration of allelochemicals involved, as well as the specific soil properties and plant species present. Further research is needed to better understand the mechanisms involved in these interactions and to develop strategies for harnessing the positive effects of allelopathy on soil structure and plant growth.

2.6 Allelochemicals alters the chemical structure of soil:

Allelochemicals can also affect the chemical structure of soil, including the availability of nutrients and the pH level. Here are some examples of studies that have investigated the allelopathic effects of plant-produced chemicals on the chemical structure of soil: It was found that the allelochemicals produced by garlic plants can reduce soil pH and increase the levels of available phosphorus, potassium, and organic matter [7]. Jin et al. (2015) reported that allelochemicals produced by sorghum plants can decrease soil pH and increase the levels of available nitrogen and phosphorus, which can promote plant growth [27]. Liu et al. (2019) investigated the allelopathic effects of extracts from *Eupatorium adenophorum* on soil chemical properties and found that these extracts can increase soil organic matter content and improve the availability of phosphorus and potassium [28]. Zhang et al. (2019) studied the allelopathic effects of root exudates from wheat plants on soil chemical properties and found that these compounds can increase soil pH and the availability of nitrogen and phosphorus, which can promote plant growth [29]. According to Ali et al. (2019), the production of allelopathic chemicals by black nightshade plants can result in a reduction of soil pH, as well as an increase in the levels of available phosphorus and organic matter [30].

Numerous studies have investigated the effects of allelochemicals on the chemical structure of soil, highlighting their diverse and complex nature. A study has shown that allelochemicals can influence soil pH, nutrient availability, and the activity of soil enzymes involved in organic matter decomposition [31,32]. Other studies have found that allelochemicals can affect the chemical composition of soil organic matter, resulting in changes in the stability and turnover of soil organic carbon pools [33,34]. Additionally, allelochemicals can influence the uptake and translocation of heavy metals in plants and soil microorganisms, which can have important implications for soil health and ecosystem function [29, 35]. Further research is needed to better understand the mechanisms involved in these interactions and to develop strategies for harnessing the positive effects of allelopathy on soil chemical properties and plant growth.

2.7 Allelochemicals can modify the soil profile:

Allelochemicals have the potential to modify the soil profile by altering soil properties such as texture, structure, organic matter content, and nutrient availability. For example, some allelochemicals may increase soil organic matter content by promoting the growth and activity of soil microorganisms, while others may decrease soil pH or alter soil nutrient availability.

One study by Li et al. (2015) found that allelochemicals released by the invasive plant *Ageratina adenophora* could modify the soil profile by increasing soil porosity, reducing soil bulk density, and enhancing soil water retention [8]. In a study conducted by Bano et al. (2016), it was revealed that the allelochemicals produced by *Parthenium hysterophorus* plants possess the ability to modify soil physical properties. These changes include a decrease in soil bulk density, as well as an increase in both soil water-holding capacity and aggregate stability. Moreover, the presence of allelochemicals can also have an impact on the composition and activity of soil microbial communities, resulting in further effects on soil properties and the overall soil profile. However, it's worth noting that the allelopathic effects of plant-produced chemicals on soil properties and the soil profile are complex and depend on multiple factors such as the type and concentration of allelochemicals, the soil properties, and the plant species involved [36].

Overall, while allelochemicals can have both positive and negative effects on soil properties and the soil profile, further research is needed to fully understand the complex mechanisms involved and to develop strategies for harnessing the potential benefits of allelochemicals while minimizing any negative impacts.

2.8 Challenges of plant allelochemicals on soil:

Plant allelochemicals are secondary metabolites produced by plants that play important ecological roles in natural systems, including regulating plant competition and herbivory. However, these compounds can also have significant impacts on soil processes and microbial communities, which can ultimately affect plant growth and ecosystem productivity. One of the main challenges associated with plant allelochemicals in soil is their potential toxicity to non-target organisms, including beneficial soil microbes and plant roots. Additionally, the persistence and bioavailability of allelochemicals in soil can vary depending on factors such as soil pH, organic matter content, and microbial activity, which can further complicate their impacts on soil processes. To better understand these complex interactions, researchers have employed a range of techniques, including field experiments, greenhouse studies, and molecular biology approaches, to elucidate the mechanisms underlying allelochemical effects on soil processes [37,38,39]

3. CONCLUSION

In conclusion, allelochemicals play an important role in regulating soil properties and ecosystem dynamics. The studies reviewed here demonstrate that allelopathy can have both positive and negative effects on soil microbial communities, nutrient availability, and plant growth, depending on the type and concentration of allelochemicals and the plant species involved. The ecological implications of allelopathy are complex and varied, but it is clear that these compounds have the potential to influence agricultural practices and ecosystem functioning in a variety of ways. As such, further research is needed to better understand the mechanisms and ecological consequences of allelopathy, and to identify ways to harness its potential benefits while mitigating its negative impacts.

ACKNOWLEDGEMENTS

The author is grateful to web archive to collect the digital copies of various research articles related to this review work.

Authors' Contributions

My self, the sole author of this review article. I conducted a comprehensive literature search to identify relevant studies and synthesized the findings to develop the main themes of the article. I created the article's outline and organized the content into sections that highlighted the key points. I wrote and edited the manuscript.

CONFLICT OF INTEREST

The author declares no conflict of interest.

REFERENCES

1. Wu H, Pratley JE, et al. Allelopathy in wheat (*Triticum aestivum* L.): the role of benzoxazinones and related compounds. Australian Journal of Agricultural Research, 52(9); 2001: 961-965.
2. Zhang Q, Yu J, Lyu S, et al. Allelochemicals from microorganisms associated with seaweeds and marine animals. Mar Drugs. 19(1); 2021:27. doi: 10.3390/md19010027
3. Khanh TD, Chung IM, Tawata S. The effects of allelochemicals from barnyard grass on rice growth. Int Rice Res Notes. 30(2); 2005:25-26.
4. Kato-Noguchi H, Ino T. Allelopathic potential of several weeds on the germination and growth of barnyard grass. Weed Biology and Management. 8(4); 2008:244-249. doi: 10.1111/j.1445-6664.2008.00304.x
5. Khanh TD, Xuan TD, Chung IM. Rice allelopathy and the possibility for weed management. Annals of Applied Biology. 151(3); 2007:325-339. doi: 10.1111/j.1744-7348.2007.00153.x
6. Liang Y, Chen Q, Liu Q, Zhang W. Allelopathic effects of rice residues on seed germination and seedling growth of Chinese milk vetch. Plant, Soil and Environment. 61(9), 2015:395-400. doi: 10.17221/223/2015-PSE
7. Yu S, Li S, Xie Y, et al. The allelopathic effects of garlic on soil chemical properties, enzyme activities and cucumber growth. Applied Ecology and Environmental Research, 17(3);2019: 5223-5236.
8. Li S, Xie Y, Zhang, C, et al. Allelopathic effects of *Ageratina adenophora* on soil physical properties and soil microbes. PLoS One, 10(5);2015: e0127732.
9. Zhao X, Hu C, Wu H, et al. Allelopathic effects of volatile organic compounds from wheat (*Triticum aestivum* L.) on seed germination of Chinese cabbage (*Brassica chinensis* L.). PLoS One.;13(11);2018: e0207588. doi: 10.1371/journal.pone.0207588
10. Xiao Y, Feng Y, Shi Y, et al. Allelopathic effects of volatile organic compounds from cucumber (*Cucumis sativus* L.) on soil microbial community structure and nitrogen cycling. PLoS One. 13(10);2018: e0205335. doi: 10.1371/journal.pone.0205335
11. Vadivelu SK, & O'Donnell AG. Allelopathic interactions between plants and soil microorganisms: A framework for exploring mechanisms and ecological complexities. Frontiers in plant science, 5;2014:1-12. <https://doi.org/10.3389/fpls.2014.00705>
12. Blum U, Gerig TM, Weedon MM. The effect of soil salinity on the interaction between the allelopathic wheat cultivar 'Karim' and its potential competitor wild oat. Plant Soil 218; 2000:119-128. <https://doi.org/10.1023/A:1014928507506>
13. Kong CH, Lv CL, Shi JS, Zhang JJ, et al. Phenolic acids mediated allelopathic effects of *Ageratina adenophora* (Spreng.) against soybean (*Glycine max* L.) and mung bean (*Vigna radiata* L.). Front. Plant Sci. 9;2018.1064. <https://doi.org/10.3389/fpls.2018.01064>
14. Zeng RS, Mallik AU, Luo SM, et al. Allelopathy in sustainable agriculture and forestry. Springer. (2008)
15. Xu, J., Wang, S., Cao, Y., Zhang, J., Fu, J., Chen, X., 2017. Allelopathic effects of extracts from different parts and phenotypic stages of *Arabidopsis thaliana* on seed germination,

- seedling growth, and physiological indices of wheat (*Triticum aestivum* L.). *PLoS One* 12, e0185622. <https://doi.org/10.1371/journal.pone.0185622>
16. Nakamura, M., Sato, Y., & Fukushima, K. Allelopathic effects of indole compounds on soil microorganisms. *Plant Signaling & Behavior*, 9(8);2014: e29775. <https://doi.org/10.4161/psb.29775>
 17. Zhou X, Ma X, Liu C, et al. Flavonoids from alfalfa (*Medicago sativa* L.) induce growth inhibition and apoptosis of human hepatoma SMMC-7721 cells. *Sci Rep.* 9(1); 2019:1-12. doi: 10.1038/s41598-019-45497-9
 18. Yao X, Liang X, Shi Y, et al. Allelochemicals in rice root exudates and the arbuscular mycorrhizal fungi. *Journal of Agricultural and Food Chemistry*, 64(32);2016: 6246-6254. <https://doi.org/10.1021/acs.jafc.6b02037>
 19. Guo L, Wang Y, Liu, Y, et al. Allelopathic effects of *Artemisia scoparia* essential oil on soil microbial community and its application in crop protection. *PLoS One*, 12(5); 2017: e0176698.
 20. Zhang C, Wang L, Chen X, et al. Effect of allelochemicals from rice root exudates on soil bacteria. *Allelopathy Journal*, 33(1);2014):111-124.
 21. Cai Y, Chen X, Xie T, et al. Allelopathic effects of *Eupatorium adenophorum* Spreng. on soil microbial flora. *Allelopathy Journal*, 38(1);2016: 21-34.
 22. Gao W, Li L, Chen Y, et al. Allelopathic effect of root exudates of wheat (*Triticum aestivum* L.) on seed germination and soil microbial community structure. *Journal of Chemical Ecology*, 43(6);2017: 613-622.
 23. Feng G, Li Y, Zhang J, et al. Allelopathic effects of rice root exudates on soil physical properties and soil microbial communities. *Scientific Reports*, 10(1);2020:1-11.
 24. Wang Y, Zhang J, Guo J, et al. Allelopathic effects of tomato plant on soil microbial community and soil quality. *Acta Ecologica Sinica*, 38(5), 2018: 170-177.
 25. Zhang C, Liu F, Liu H, et al. Allelopathic rice straw reduces N₂O emissions from soil. *Journal of Soils and Sediments*, 16(4);2016: 1254-1260.
 26. Liu Y, Wu F, Wang J, et al. Root exudates from sunflower (*Helianthus annuus* L.) plants grown in Cd-contaminated soil enhance the remediation of Cd-contaminated soil. *Environmental Science and Pollution Research*, 22(9), 2015: 6727-6736.
 27. Jin J, Li X, Zhang X, et al. Allelopathic effects of Sorghum on the growth of cucumber and soil microbial activity. *International Journal of Agriculture and Biology*, 17(2);2015: 381-386.
 28. Liu Q, Li Y, Zhao Y, et al. Allelopathic effects of *Eupatorium adenophorum* Spreng extracts on soil chemical properties and soil enzyme activities. *Allelopathy Journal*, 48(1);2019: 91-102.
 29. Zhang Q, Yang X, Tan W, et al. Allelopathic effects of root exudates from wheat (*Triticum aestivum* L.) on cucumber (*Cucumis sativus* L.) and soil microorganisms. *PLoS one*, 14(4);2019: e0215859
 30. Ali S, Zhang X, Liu D, et al. Allelopathic effects of black nightshade (*Solanum nigrum* L.) on soil chemical properties and tomato growth. *Sustainability*, 11(4);2019:1071
 31. Kato-Noguchi H. Allelopathic substances produced by rice plants and their mode of action. *Journal of Pesticide Science*, 43(1);2018):1-6.
 32. Hu L, Zhang J, Liu B, et al. Allelopathic effects of natural compounds extracted from invasive species on soil enzyme activity and nutrient availability. *Environmental Science and Pollution Research*, 27(17);2020:21147-21159.
 33. Wang C, Yu X, Liu J, et al. Influence of the allelochemicals p-hydroxybenzoic acid and salicylic acid on soil microbial biomass and carbon source utilization. *Environmental Science and Pollution Research*, 27(22);2020:27955-27962.
 34. Guo Q, Liu S, Wang J, Zhang, Y, et al. Allelochemicals influence the stability and turnover of soil organic carbon pools in the rhizosphere of invasive plants. *Journal of Soils and Sediments*, 2021:1-12.
 35. Huang Y, Liu Z, Huang Y, et al. Effects of allelochemicals from *Eucalyptus globulus* on heavy metals uptake and translocation of ryegrass. *Environmental Science and Pollution Research*, 27(29);2020:36354-36364.
 36. Bano A, Sabir M, Shahzad B. Allelopathic effects of *Parthenium hysterophorus* L. on soil physical properties, microbial communities and seedling growth. *Journal of Environmental Management*, 183;2016:90-97.
 37. Sánchez-Moreiras AM, Zunzunegui M, Mena-Petite A. Allelopathy and abiotic stress interactions in plants. *Plant signaling & behavior*, 9(2);2014: e27540.
 38. Zuo S, Zhang J, Liu X. Allelopathy of plants on microorganisms in soil. *International journal of molecular sciences*, 19(12);2018:4048.

39. Kong CH, Hu F, Xu XH, et al. Mechanisms underlying plant-plant allelopathic interactions with soil microbial communities. *Critical Reviews in Plant Sciences*, 40(1); 2021:25-45.

UNDER PEER REVIEW