

Establishing Climatic Change on Rainfall Trend, Variation and Change Point Pattern in Benin City, Nigeria.

ABSTRACT

This study examined the effect of changing climate for thirty-six years (1982-2017) on 24-hourly annual maximum time series (AMS) rainfall data for Benin City. The results show the existence of a trend, variation in magnitude, and change point dates. Trend analysis was performed using Mann-Kendall (MK) test and Sen's slope estimator (SSE) applied to obtain the trend magnitude. A statistically significant trend was found for Benin City at a 5% level of significance. The MK $|Z|$ statistic varied from 2.0710 to 2.1550 for IMD and 2.0844 to 2.0995 for MCIMD downscaling models, which were both greater than the critical Z-value of 1.96. The SSE gave the magnitude of the trend variation rate due to climate change at 24-hour duration as 1.008 mm/year (10.08 mm/decade) and 1.1376 mm/year (11.376 mm/decade) for the IMD and MCIMD models, respectively. Also, the trend change point analysis was conducted using the distribution-free cumulative sum test (CUSUM) and the sequential Mann-Kendall test (SQMK) which has rainfall pattern definite change point date for Benin City as 2005 from where continued a positive trend which intensified through the year 2015. Thus, proving the existence of increasing changing climatic condition for the study area.

Keywords

Rainfall data, 24-hourly AMS, trend analysis, change point, changing climate.

1. INTRODUCTION

Rainfall is an indicator for climate change which could be an increasing or decreasing phenomena. The rainfall measurements constitutes series of data that can be analyzed as a time series. The characteristic of given set of time series observations could possess a gradual process in pattern termed as trend or possess abrupt changes that are described as trend change point. Recent studies on climate change have mostly focused on long-term variability of temperature and rainfall which are, as it were, considered the most important climate change indicators. However, evidence of climate change elsewhere and in Nigeria's meteorological system indicates an increasing rainfall trend in the coastal region and a decreasing rainfall inland (Adejuwon,

2012; Olaniyi et al., 2013; Nwaogazie and Ologhadien, 2014; Ebele and Emodi, 2016; Elisha et al., 2017; Sam, et al., 2021).

Time-series analysis is an applied tool for the detection and quantitative description of the generating process which is characteristic of a given set of observations. The generative process of the time series possesses both gradual and abrupt changes that are described as either a trend or a trend change point. Therefore, trend analysis deal with evaluating gradual future events from past measured data, while change point detection deal with determining the unexpected, structural, changes in the time series data properties like mean or variance (Mann, 1945; Zhang et al., 2006). There are different approaches in the literature used in calculating both the trend and change point. A parametric test is applied as a powerful tool, wherein the condition for its application required in trend analysis for a given data set are normal distribution and independent. While the non-parametric test data should be independent, and tolerant of outliers. The non-parametric test for trend analysis in time series data that are most commonly applied also adopted for this study is the Mann-Kendall test (Mann, 1945; Kendall, 1975, Lehmann, 1975; Longobardi and Villani, 2010; Yanming et al., 2011) and the Sen Slope estimator is used for evaluating the trend magnitude and its intercept (Sen, 1968).

Different techniques in the literature for the analysis of trend change points available are Wild binary segmentation, the Bayesian analysis of change points, the E-Agglomerative algorithm, and Iterative robust detection methods (Baranowski & Frylewicz, 2014; Fryzlewicz, 2013; Erdman & Emerson, 2007). However, other methods of trend change-point analysis also available that is applied in this study is the Distribution-free cumulative sum (CUSUM) and the Sequential Mann-Kendall (SQMN) tests (McGilchrist & Woodyer, 1975; Taubenheim, 1989; Sneyers, 1990).

The purpose of this study is to determine the effect of changing climate in Benin City, Nigeria using measured time series rainfall data recorded for 36 years (1982-2017) in terms of trend, its variation and magnitude, and the trend change point dates. Trend analysis was performed using the Mann-Kendall (MK) test, Sen's slope estimator (SSE) was applied to obtain the trend magnitude, while the trend change point analysis was conducted using the distribution-free cumulative sum test (CUSUM) and the sequential Mann-Kendall test (SQMK). The 'python statsmodel library-pymannkendall' (Seabold and Perktold, 2010) and 'trendchange' (R Core Team, 2021; Patakamuri, 2022) were the two different open-source software packages adopted.

2. METHODOLOGY

2.1 Study Area

The study station Benin City is located in Edo State in the South-South region of Nigeria. The study area lies between latitudes $6^{\circ} 52' N$ - $6^{\circ} 21' N$ and longitudes $5^{\circ} 34' E$ - $5^{\circ} 44' E$, as shown in Figure 1. The area is located at an altitude/elevation of Nonem (0 ft) above sea level. Benin City has a tropical Savanna climate which is dry and wet. The driest month takes place in December period with 22.44 mm (0.88 inches) rainfall and the wettest month occurs in July with 332.95 mm (13.11 inches) of rainfall. The study area receives about 189.68 mm (7.47 inches) of rainfall with about 274.89 rainy days annually (ie., 75.31 % of the period). The annual highest and lowest temperature of $33.89^{\circ} C$ ($93.0^{\circ} F$) and $24.5^{\circ} C$ ($76.1^{\circ} F$) are recorded, respectively.




Figure 1. Map of Edo State showing Benin City (Edo State Govt. M L S, 2009).

2.2 Data Collection

The measured rainfall data for this study was collected for 36 years (1982-2017) from the Nigeria Meteorological Agency (NIMET) meteorological gauge station at Benin City. The collected data on rainfall amount was recorded in mm against corresponding durations recorded in minutes. Thereafter, the data were sorted out by extraction of the maximum daily (24-hourly) rainfall amount for each month from where the maximum for each year was extracted to form the 24-hourly annual maximum series (AMS) for the rest of the 36 years interval.

2.3 Downscaling of 24-hourly Rainfall Data into Shorter Durations Time Series Data

Maximum daily rainfall data were collated for each month for the 36 years for various analysis of the time series data. The maximum 24-hourly annual maximum series (AMS) rainfall

data extracted were further downscaled into shorter duration rainfall as detailed in Sam, et al., (2021) for the trend study. The formula applied were the proposed Indian Meteorological Department (IMD) initially proposed by Ramaseshan (1996) and the Modified Chowdhury Indian Meteorological Department (MCIMD) (Rashid et al., 2012) as given in Equations (1) and (2), respectively, while Equation (3) was used for conversion of rainfall amount to its intensity equivalent;

$$R_t = R_{24} \left(\frac{t}{24} \right)^n \quad (1)$$

$$R_t = R_{24} \left(\frac{t}{24} \right)^n + C \quad (2)$$

$$I = \frac{R_t}{t} \quad (3)$$

Where: R_t is the required rainfall depth in mm for durations less than 24 hours, R_{24} is the daily rainfall depth (mm), t is the required duration (hours), n is an exponential constant = 1/3; while for the MCIMD method the n and C are determined constants, where I denotes the rainfall intensity (mm/hr).

2.4 Mann-Kendall Trend & Sen' Slope Estimation Test

Mann Kendall test (Mann 1945, Kendall 1975, Gilbert 1987) is a rank-based statistical test that helps to check if there is a monotonous upward or downward trend existing in a time series data. Mann Kendall (MK) test is a non-parametric test in which the distribution of the data set must not be auto-correlated (no serial correlated). The conditions required to achieve a relatively error-free MK test result are well treated in the literature (Von Storch and Navarra, 1995; Hamed and Rao, 1998; Yue et al., 2002; Yue and Wang, 2002). Trend Free Pre-Whitening

(TFPW) is used to remove serial correlation where it exists in the rainfall intensity. Sam, et al., (2022) documented the procedure used in performing the Mann-Kendall test to find if a monotonous positive or negative trend exist in the time series data. Where serial correlation exist in the data set the TFPW can be applied to remove the serial correlation. Also, where the ACF is not significant, the MK test is applied directly to the original data set. The python statsmodel library- pymannkendall (Seabold and Perktold, 2010) was used to compute both the ACF at lag 1 and in applying TFPW to the time series data set before MK test was carried out on the pre-whitened data set. The trend magnitude was computed using the Sen Slope estimator analyzed with microsoft Excel stat 2016 and python pymannkendall library.

2.5 Trend Change-Point Test

Identifying trends and time of trend change, play a significant part in precipitation parameters and time-series studies. To determine the trend change-point, two non-parametric testing methods, the Distribution-free cumulative sum (CUSUM) test (McGilchrist and Woodyor, 1975) and the Sequential Mann-Kendall test (Taubenheim, 1989; Sneyers, 1990) were applied. The free CUSUM test utilizes a cumulative sum chart, while the sequential Mann-Kendall (SQMK) test uses each sample point sequence sequentially treated in both prograde and retrograde procedures. Sam, et al. (2022a) also demonstrated in detail the application of both non-parametric approaches. The package “trendchange” was used for change-point analysis. The software package is an open-source library freely available via the CRAN repository and version 1.2 (R Core Team, 2021; Patakamuri, 2022).

3 Results and Discussion

3.1 Results

3.1.1 Different Data Sets Used for the Time Series Analysis

Detecting trend in the time series data was carried out for three different sets of data. The first set of data was rainfall measurements obtained from downscaling 24-hourly rainfall data for Benin City (see Table 1) using the IMD formula in Equations (1) and (3), while the second set of data was the rainfall intensities obtained from downscaling the same 24-hourly rainfall data using the MCIMD model in Equations (2) and (3). The third data set is the initial sorted 24-hourly monthly maximum series (MMS) for the 36 years (1982-2017 inclusive) study period.

Table 1: 24-hourly annual maximum series (AMS) rainfall data for Benin City (1982-2017)

Year	Rainfall Amount (mm)	Year	Rainfall Amount(mm)	Year	Rainfall Amount(mm)
1982	164.3	1994	156.1	2006	109.8
1983	93.7	1995	93.3	2007	122.6
1984	54.7	1996	121.1	2008	125.3
1985	61.6	1997	122.6	2009	84.1
1986	59.6	1998	87.8	2010	183.6
1987	112.4	1999	78.9	2011	169.6
1988	88.5	2000	99.1	2012	83.8
1989	133.4	2001	96.9	2013	131.6
1990	114.4	2002	95.8	2014	119.2
1991	110.4	2003	102.5	2015	106.6
1992	120.8	2004	101.7	2016	111.0
1993	79.9	2005	114.6	2017	266.6

3.1.2 Trend Change Analysis Using Mann-Kendall (MK) Statistic

The performance of the MK test require the computation of the autocorrelation function (ACF) which result was insignificant for ACF at lag 1 at 95% Confidence interval (see Figure 2 plot). Therefore, no trend free pre-whitening (TFPW) was applied, hence, the MK test was conducted directly on the original time series data. The results of the MK test are presented in Table 2. The $|Z|$ value for MCIMD model data varied for the durations ranging from 2.0844 to 2.0986 with slight variations for the IMD model data ranging from 2.0710 to 2.1550. All the MK statistic $|Z|$ values were greater than the Critical Z-value of 1.96. Similarly, the p-value for the MCIMD method data produced a range of value varying from 0.0359 to 0.0371 for the durations, while the IMD method data p-value were 0.0312 to 0.0384. These p-values obtained were all also below alpha, $\alpha = 0.05$ level of significance. These results reinforces the existence of statistically significant trend in the various downscaled short duration data.




Figure 2: Correlogram of ACF at lag-1 for various 24-hourly AMS downscaled durations using MCIMD model for Benin City.

3.1.3 Evaluation of Trend Magnitude & Variation

Evaluation of the magnitude of the trend was carried out using Sen's slope estimator, as the trend line intercept and slope, Q_i indicated in Figures 3 to 5 and values presented in Table 2. The result of the slope shows decreasing positive values of 0.9574 (at 0.25 hour) to 0.0420 (at 24 hour) and 1.3893 (at 0.25 hour) to 0.0474 (at 24 hours) for IMD and MCIMD models, respectively. The intercept for the IMD method also decreased from 79.3963 to 3.8647 at 0.25 and 24 hours, respectively. The MCIMD data produced higher intercept values which similarly decreased from 139.0875 to 4.0197 at 0.25 to 24 hours, respectively. From Figure 3 graph, the result obtained for the 24-hourly MMS data confirmed the trend variation magnitude test results by giving slope value of 0.020 mm/hr/year and intercept of 1.5168. The p-value of 0.00118 which is far less than alpha value at 5% significant level was obtained.




Figure 3: 24-hourly MMS rainfall intensities trend pattern for Benin City (1982-2017).

Table 2: Mann-Kendall test and Sen Slope estimates result for Benin City

Time (hrs)	Statistic	IMD	MCIMD
		Value	Value
0.25	Z	2.0978	2.0978
	p-value	0.0359	0.0359
	Q_i	0.9574	1.3893
	Intercept	79.3963	139.0875
0.5	Z	2.0978	2.0978
	p-value	0.0359	0.0359
	Q_i	0.5999	0.8274
	Intercept	50.1000	80.5208
0.75	Z	2.0978	2.0978
	p-value	0.0359	0.0359
	Q_i	0.4609	0.6095
	Intercept	38.1838	58.6346
1	Z	2.0844	2.0844
	p-value	0.0371	0.0371
	Q_i	0.3741	0.4929
	Intercept	31.6538	46.825
2	Z	2.0710	2.0844
	p-value	0.0384	0.0371
	Q_i	0.2352	0.2935
	Intercept	19.93	27.314
6	Z	2.0986	2.0986
	p-value	0.0359	0.0359
	Q_i	0.1155	0.1232
	Intercept	9.5292	11.7938
12	Z	2.0718	2.0986
	p-value	0.0383	0.0359
	Q_i	0.0707	0.0768
	Intercept	6.0625	6.8562
24	Z	2.1550	2.0995
	p-value	0.0312	0.0359
	Q_i	0.0420	0.0474
	Intercept	3.8647	4.0197

Level of significant $\alpha = 0.05$, where Z = standardized Mann Kendall statistic, Q_i = Sen Slope (mm/hr/year), Critical Z-value = 1.96.

Table 3: Distribution Free CUSUM Result for Benin

Parameters	Values
Maximum CUMSUM value	5
Critical value at 90% CI	7.32
Critical value at 95% CI	8.16

Critical value at 99% CI
Date of Change

9.78
2004 and 2006




Figure 4: 24-hourly AMS rainfall intensities trend pattern for IMD model downscaled durations for Benin City (1982-2017).




Figure 5: 24-hourly AMS rainfall intensities trend pattern for MCIMD model downscaled durations for Benin City (1982-2017).

3.1.4 Evaluation of Time Series Data for Trends Change-Point

The result for the trend graph plots of the 24-hourly AMS rainfall intensities for Benin City for both long (24 hours) and short (0.25 hours) durations are presented in Figures 4 and 5,

respectively. Also carried out was trend change-point analysis for all downscaled durations and the results of the distribution-free CUSUM and the Sequential Mann-Kendall tests are presented in Tables 3 & 4. Figures 6 and 7 show distribution-free CUSUM plot and Sequential Mann-Kendall plot for 24-hourly AMS rainfall intensities, respectively, which were obtained via the “trendchange” software package (R Core Team, 2021; Patakamuri, 2022).

Table 4: Trend change point for different downscaled duration rainfall intensities for various stations

Rainfall Durations (mins)	CUSUM	SQMK
15	2004 and 2006	2005
30	2004 and 2006	2005
45	2004 and 2006	2005
60	2004 and 2006	2005
120	2004 and 2006	2005
360	2004 and 2006	2005
720	2004 and 2006	2005
1440	2004 and 2006	2005




Figure 6: Distribution-free CUSUM plot for 24-hourly AMS rainfall intensity for Benin.




Figure 7: Sequential Mann-Kendall plot for 24-hourly AMS rainfall intensity for Benin

3.2 Discussion of Results

3.2.1 Analysis of 24-Hourly AMS Non-stationary Rainfall Trend

This investigation established the existence of meteorological trends and variation in 24-hourly AMS extracted data from the 24-hourly Monthly Maximum Series (MMS) historical precipitation (rainfall) data (HPD). Three sets of meteorological time-series data were applied. The IMD model data and the MCIMD model data downscaled from Table 1 using Equations (1) and (2), respectively, with those of the initially extracted MMS recorded rainfall intensities.

3.2.2 Analysis of 24-Hourly AMS Rainfall Trend Change-Point

Change point analysis identifies the most probable date for occurrence of significant change in the time series. Analysis of trend change points was for all the 24-hourly AMS time series downscaled durations. Results of the distribution-free cumulative sum (CUSUM) and sequential Mann-Kendall (SQMK) tests are presented in Table 4. Figures 6 and 7 provide the distribution-free CUSUM and SQMK plots, respectively, for different durations of the 24-hourly AMS rainfall intensity. Interpreting the result for Benin, the distribution-free CUSUM plot showed two change dates of 2004 and 2006 at the maximum CUSUM value of 5.00 less than the critical value of 8.16 at 95% CI shown in Table 3. Contrastingly, the SQMK plot produced a

definite intersection between the pro-grade and retrograde in 2005 at 95% CI. Therefore, it could be deduced that the trend change point process intensified for 3 years, while the ideal change point year for Benin City took place in the year 2005 that is, the intersection point between the pro-grade and retrograde at 95% confidence interval as indicated in Figure 7. The positive trend continued and intensified through the year 2015, another point of intersection in the SQMK plot.

3.2.3 Autocorrelation and TFPW of 24-Hourly AMS Rainfall

The statistical tool adopted for a non-stationarity check if trends exist in a time-series data is the Mann-Kendall (MK) test after autocorrelation analysis at a 95% Confidence interval. The Correlogram in Figure 2 showed that the ACF at lag 1 was statistically insignificant for all downscaled rainfall durations. The 24-hourly MMS data subjected to autocorrelation also exhibited the same features. Therefore, ACF values for the time-series data at a 95% Confidence interval were insignificant. This result does not require any trend-free pre-whitening (TFPW) because the time series data had no significant serial autocorrelation. Hence, the MK test was applied directly to the time series data.

3.2.4 Trend and Variations in MK and Sen's Slope Test Results

The Mann-Kendall test results for Benin are presented in Table 2. The indices measured by the MK test for the trend are the p-value or alpha value of 0.05 (5% level of significance) and the standardized MK statistics compared to the critical Z-value at 1.96. The hypothesis examined for the MK test is that for H_a : Alternate hypothesis, there is a trend in the rainfall intensity over time, and for H_o : Null hypothesis, there is no trend in the rainfall intensity over time. Therefore considering the various result for the MK and Sen's Slope tests in Table 2, we have the following observations:-

The calculated p-value varied from 0.0312 to 0.0383 and 0.0359 to 0.0371 for IMD and MCIMD time series data, respectively. The p-values were less than the alpha value of 0.05 at a 5% level of significance. The MK statistic varied from 2.0710 to 2.1550 for both IMD and MCIMD data series and were all greater than the critical Z-value of 1.96. The alternate hypothesis is accepted that a trend exists in the time series data.

Sen's Slope test also showed that the magnitude of the slope varied from 0.0420 to 0.9574 and 0.0470 to 1.3893 for IMD and MCIMD time series data, respectively. The intercept, likewise varied from 3.8647 to 79.3963 and 4.0197 to 139.0875 for the IMD and MCIMD data series, respectively. The data series for Benin indicates the same trend pattern and variation in the values of magnitudes of the slope and the intercept as those of sister stations in the region such as Uyo (Sam, et al., 2022). That is an increase in value from a higher duration of 24 hours to a lower duration of 0.25 hours. The result obtained for Benin did not conform with the opinion of Ologhadien's (2019) report of insignificant trends on observed time series data. The Mann-Kendall result suggests that when developing the intensity-duration-frequency (IDF) model, the change in the behavioral parameter extremes such as shape, location and scale in the rainfall time series data must be put in-view in the IDF modeling.

The Sen's Slope results also proved that the magnitude of the trend decreases as the duration of rainfall increased that is to say that shorter durations tend to exhibit higher values than longer durations (Sam et al., 2022).

3.2.5 Analysis of Variations in Magnitude and Rainfall Trend Pattern

The results of the MK trend and Sen Slope tests result presented in Tables 2 also indicated that both tests exhibited a high degree of consistency. They both showed a statistically significant positive trend increase in their values of computed variables for 24 hours higher

duration to 0.25 hours lower durations. The observed performance in the consistency of both tests conforms with some earlier publications (Yue *et al.*, 2002; Shadmani *et al.*, 2012; Ahmad *et al.*, 2015).

The Sen's Slope estimator (SSE) used for the quantification of the rate of change in the magnitude of rainfall intensity showed a positive trend variation statistically significant for the study area.. For instance, the slope, Q , and the rate of variation observed for the station from Table 2 is given as 0.0420 and 0.0474 mm/hr/year for the IMD and the MCIMD data series, respectively, which translates to variation rate of 1.008 mm/year (10.08 mm/decade) and 1.1376 mm/year (11.376 mm/decade) for the IMD and MCIMD models, respectively.

The trend test for the 24-hourly monthly maximum series (MMS) in Figure 3 also confirmed the test statistic, with a p-value of 0.00118 which is far less than the alpha value at a 5% significant level. Therefore, the 24-hourly AMS data could adequately serve as a representative sample for the time series data in IDF modeling for the prediction of rainfall intensity. The trend variation rate in extreme precipitation is brought about by behavioral parameter extreme changes occasioned by climatic change. Human activity also generates greenhouse gas emissions with resultant land use changes. These factors result in extreme precipitation events that lead to landslides and flooding.

These results agreed with the publications of earlier research along the coastal stations of the Gulf of Guinea. For instance, Ayensu (2004,) got 13mm/decade for Ghana, and Nwaogazie and Ologhadien (2014) also had 55.2 mm/decade, and Sam et al. (2022) had a variable rate of 21.288 mm/decade for Uyo, for gauge stations in the Niger Delta. The results did not agree with those of Okafor *et al.* (2017), whose work was on the Nigerian hinterland and found a decreasing trend in the rate of the variability of rainfall and in the MK test statistic. These are evidence of climatic

change variability in Nigeria whose trend shows rainfall increase in the coastal region (Olaniyi *et al.*, 2013; Nwaogazie & Ologhadien, 2014, Sam *et al.*, 2022) and decrease in the continental interiors (Ebele & Emodi, 2016; Elisha *et al.*, 2017).

4.2 CONCLUSION

The consequences of changing climate were found to exist for Benin City in this study. Statistically significant trends were found for Benin City with the p-values less than the alpha value of 0.05 at 5% level of significance. The calculated p-value varied from 0.0312 to 0.0383 and 0.0359 to 0.0371 for IMD and MCIMD time series downscaled data, respectively. Also, the MK $|Z|$ statistic varied from 2.0710 to 2.1550 for IMD and 2.0844 to 2.0995 for MCIMD downscaling model series, which were both greater than the critical Z-value of 1.96.

The magnitude of the trend and variation rate contributed by climate change at 24-hour duration was 1.008 mm/year (10.08 mm/decade) and 1.1376 mm/year (11.376 mm/decade) for the IMD and MCIMD models, respectively. The rainfall pattern change point dates for Benin City were found to be from 2005 from where there continued to be a positive trend that intensified through the year 2015.

References

- Adejuwon, J. O. (2012). *Rainfall seasonality in the Niger delta belt, Nigeria*. Journal of Geography and Regional Planning, Vol. 5, No. 2, pp. 51-60.
- Ahmad, I., Tango, D., Wang, T., Wang, M. and Wagan, B (2015), "Precipitation Trends over Time Using Mann-Kendall and Spearman's rho Tests in Swat River Basin, Pakistan", *Advances in Meteorology*, 2015 (431860), 1-15.
- Ayensu, A. (2004). *Assessment of Climate Change and Vulnerability Trends in Temperature and Rainfall*. J. Appl. Sci. Technol. (JAST), Vol. 9, No. 112, pp. 21-27.
- Baranowski, R. and Fryzlewicz, P. (2014). WBS: Wild Binary Segmentation for Multiple Change-Point Detection. R Package Version 1.1

- Cheng, L. and AghaKouchak, A. (2014). *Non-Stationarity Precipitation Intensity-Duration-Frequency Curves for Infrastructure Design in a Changing Climate*. Science Reports, Vol. 4. No. 7093, pp. 1-6.
- Cheng, L., A. AghaKouchak, Gilleland, E Katz, R. W. (2014). *Non-stationary Extreme Value Analysis in a Changing Climate*. Climate Change, Vol. 127, No. 2, pp. 353-369.
- Duhan, D. and Pandey, A. (2013). *Statistical Analysis of Long Term Spatial and Temporal Trends of Precipitation during 1901–2002 at Madhya Pradesh, India*. Atmospheric Research, Vol. 122, pp. 136–149
- Ebele, N. E. and Emodi, N. V. (2016). *Climate Change and its Impact in Nigerian Economy*. Journal of Scientific Research and Reports, Vol. 10, No. 6, pp. 1-13.
- Edo State Government (2009). Ministry of Lands & Survey, Benin City, Edo State
- Elisha, I., Sawa, B. A. and Lawrence, E. U. (2017). *Evidence of Climate Change and Adaptation Strategies among Grain Farmers in Sokoto State, Nigeria*. IOSR Journal of Environmental Science, Toxicology and Food Technology (IOSR-JESTFT), Vol. 11, No. 3, pp. 1-7.
- El Adlouni, S., Ouarda, T. B. M. J., Zhang, X., Roy, R., and Bobee, B. (2007). *Generalized Maximum Likelihood Estimators for the Non-Stationary Generalized Extreme Value Model*. Water Resour. Res., Vol. 43, No. W03410, <https://doi.org/10.1029/2005WR004545>.
- Erdman, C. and Emerson, J. W. (2007). BCP: An R Package for Performing a Bayesian Analysis of Change Point Problems. Journal of Statistical Software 23(3):1–13
- Fryzlewicz, P. (2013). Wild Binary Segmentation for Multiple Changepoint Detection. Department of Statistics, London School of Economics, London
- Ganguli, P. and Coulibaly, P. (2017). *Does Non-stationarity In Rainfall Require Non-stationary Intensity-Duration-Frequency Curves?* Hydrology and Earth System Sciences, Vol. 21, pp. 6461-6483.
- Gilbert, R. O. (1987). *Statistical Methods for Environmental Pollution Monitoring*. John Wiley and Sons, New York.
- Hall, M. J. and O’Connell, B. E. (1972). Time Series Analysis of Mean Daily River Flow. *Water and Water Engineering*, Vol. 5, pp. 125-133.
- Hamed, K. H. and Rao, A. R. (1998). *A Modified Mann-Kendall Trend Test for Auto-correlated Data*. Journal of Hydrology, Vol. 204, pp. 182-196.

- Hirsch, R. M., Slack, J. R. and Smith, R.A. (1982). *Techniques of Trend Analysis for Monthly Water Quality Data*. Water Resources Research, Vol.18, pp. 107-121.
- IPCC (2007). *Climate Change 2007, the Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the IPCC*. Solomon, S., Qin, D., Manning, M., Chen, Z., Marquis, M., Averyt, K. B., Tignor, M. and Miller, H. L (Eds). Cambridge University.
- Jung, M., Kim, H., Mallari, K. J. B., Pak, G and Yoon, J. (2015). *Analysis of Effect of Climate Change on Runoff in an Urban Drainage System: A Case Study from Seoul, Korea*. Water and Science Technology, Vol. 75, No. 5, pp. 653-660.
- Kendall, M. G. (1975). *Rank Correlation Methods*. Griffin, London, UK.
- Lehmann, E. L. (1975). *Non-parametrics, Statistical Methods Based on Ranks*. Holden-Day, San Francisco, Calif., USA.
- Longobardi, A. and Villani, P. (2010). *Trend Analysis of Annual and Seasonal Rainfall Time Series in the Mediterranean Area*. International Journal of Climatology, Vol. 30, No. 10, pp. 1538–1546
- Matteson, D. S. James, N. A. (2013). A Nonparametric Approach for Multiple Change Point Analysis of Multivariate Data. *Journal of American Statistical Association* 109(505):334–345. doi:10.1080/01621459.2013.849605
- Mann, H. B. (1945). *Nonparametric Tests against Trend*. *Econometrica*, Vol. 13, pp. 245–259. <https://doi.org/10.2307/1907187>
- McGilchrist, C.A. and Woodyer, K.D. (1975). *Note on a distribution-free cusum technique*. *Technometrics*, Vol. 17, pp. 321–325.
- Nwaogazie, I. L. and Ologhadien, I. (2014). *Trend in Climate Change and Vulnerability Assessment of Nigerian Gulf of Guinea*. *Standard Scientific Research and Essays*, Vol. 2, No. 10, pp. 516 -523.
- Okafor, G.C., Jimoh, O.D and Larbi, K.I. (2017). *Detecting changes in Hydro-climatic Variables during the Last Four Decades (1975 – 2014) on Downstream Kaduna River Catchment, Nigeria*. *Atmospheric and Climate Sciences*, Vol. 7, pp. 161-175.
- Okon, K.E., Oworen, U., Daniel, K. S., and Okon, I. K. (2019). *Occupational hazard and work safety among chainsaw operators in Nigeria*. *Environment*. Vol. 3, pp. 8-20. DOI: 10.31058/j.envi.2019.33002.
- Olaniyi, O. A., Funmlayo, O. A., and Olutimehin, I. O. (2014). *Review of Climate Change and its Effect on Nigeria Ecosystem*. *International Journal of Environment and Pollution Research*. Vol. 2, No. 3, pp. 70-81.

- Patakamuri, S. K. (2022). *Trendchange: Innovative Trend Analysis and Time-Series Change Point Analysis*; The R project for statistical computing; Viena, Austria.
- Peterson, B. J., Holmes, R. M., and McClelland, J. W. (2002). *Increasing River Discharge to the Arctic Ocean*. *Science*, Vol. 298, No. 5601, pp. 2171–2173.
- Rashid, M., S. Faruque & J. Alam, 2012. Modelling of Short Duration Rainfall Intensity Duration Frequency (SDRIDF) Equation for Sylhet City in Bangladesh. *ARPN J. Sci. Technol.* 2, 92-95.
- Ramaseshan, S. 1996. *Urban Hydrology in Different Climate Conditions*. Lecture Notes of the International Course on Urban Drainage in Developing Countries, Regional Engineering College, Warangal, India.
- R Core Team (2021). *R: A language and environment for statistical computing*. R Foundation for Statistical Computing, Vienna, Austria.
- Sam, M. G., Nwaogazie, I. L. and Ikebude, C. (2021). *Improving Indian Meteorological Department Method for 24-hourly Rainfall Downscaling to Shorter Durations for IDF Modeling*. *International Journal of Hydrology*, Vol. 52, pp. 72-82. DOI: 10.15406/ijh.2021.05.00268.
- Sam, M. G., Nwaogazie, I. L. and Ikebude, C. (2022). *Non-Stationary Trend Change Point Pattern Using 24-Hourly Annual Maximum Series (AMS) Precipitation Data*. *Journal of Water Resources and Protection*, 14, 592-609. <https://doi.org/10.4236/jwarp.2022.148031>.
- Sam, M. G., Nwaogazie, I. L. and Ikebude, C. (2022a). *Climate Change and Trend Analysis of 24-Hourly Annual Maximum Series Using Mann-Kendall and Sen Slope Methods for Rainfall IDF Modelling*. *International Journal of Environment and Climate Change*, 12(2):44-60. DOI:10.9734/IJECC/2022/v12i230628.
- Seabold, S., and Perktold, J. (2010). *Satsmodel: Econometric and statistical modeling with python*. In 9th Python in Science Conference.
- Sen, P. K. (1968). *Estimates of the Regression Coefficient Based on Kendall's tau*. *Journal of the American Statistical Association*, Vol. 63, No. 324, pp.1379–1389. <https://doi.org/10.1080/01621459.1968.10480934>
- Shadmani, M., Maron, S. and Roknian, M. (2012). *Trend Analysis in Reference Evapotranspiration Using Mann-Kendall and Spearman's Rho Test in Arid Regions of Iran*. *Water Resources Management*, 26(1):211-224
- Shi, Y. F., Shen, Y. P. and Hu, R. J. (2002). *Preliminary Study on Signal, Impact and Foreground of Climatic Shift from Warm-dry to Warm-humid in Northwest China*. *Journal of Glaciology and Geocryology*, Vol. 24, pp. 219–226.
- Sneyers, R. (1990.). *On the Statistical Analysis of Series of Observations*. World Meteorological Organization: Geneva, Switzerland,

- Sule, B. F. and Ajala, O. F. (2016). *Investigating the Non-stationarity of Historical Data on Rainfall at Illorin North Central Nigeria*. Journal of Scientific and Engineering Research, Vol. 3, No. 3, pp. 537-548.
- Taubenheim, J. (1989). *An easy procedure for detecting a discontinuity in a digital time series*. Zeitschrift für Meteorol, Vol. 39, pp. 344–347.
- Von Storch, V. H. (1995). *Misuses of Statistical Analysis in Climate Research*. In: Storch, H. V. and Navarra, A. Eds., *Analysis of Climate Variability: Applications of Statistical Techniques*, Springer, Berlin, pp. 11-26.
- Von Storch, H. and Navarra, A. (1995). *Analysis of Climate Variability: Applications of Statistical Techniques*. Springer Verlag, Berlin, pp. 352.
- Yanming, Z., Jun. W. and Xinhua, W. (2011). *Study on the Change Trend of Precipitation and Temperature in Kunming City based on Mann-Kendall Analysis*, in *Future Computer, Communication, Control and Automation*, vol. 119 of *Advances in Intelligent and Soft Computing*, pp. 505-513, Springer, Berlin, Germany.
- Yue, S. and Wang, C. Y. (2002). *The Applicability of Pre-Whitening to Eliminate the Influence of Serial Correlation on the Mann-Kendall Test*. Water Resources Research, Vol. 38, No. 41, pp. 4-7
- Yue, S., Pilon, P., Phinney, B. and Cavadias, G. (2002) *The Influence of Autocorrelation on the Ability to Detect Trend in Hydrological Series*. Hydrological Processes, Vol. 16, pp. 1807-1829. <https://doi.org/10.1002/hyp.1095>.
- Zhang, Q., Liu, C., Xu, C. Y., Xu, Y. and Jiang, T. (2006). *Observed Trends of Annual Maximum Water Level and Streamflow during Past 130 Years in the Yangtze River Basin, China*. Journal of Hydrology, Vol. 324, No. 1-4, pp. 255–265.