

Effect of Biofertilizers and foliar application of zinc on growth and yield of Groundnut (*Arachis hypogea- L.*)

Abstract

A field experiment was conducted during Zaid 2022 at Crop Research Farm, Department of Agronomy, SHUATS, Prayagraj (U.P). The soil of experimental plot was sandy loam in texture, nearly neutral in soil reaction (pH 7.1), low in organic carbon (0.36 %), available N (171.48 kg/ha), available P (15.2 kg/ha) and available K (232.5 kg/ha). The experiment was laid out in Randomized Block Design with ten treatments each replicated thrice on the basis of one year experimentation. [\(The purpose of the work was missing\) \(Insert the statistical analysis used in the work\)](#). The results showed that application of PSB 20g/kg seed + 0.75% ZnSO₄ was recorded significantly higher plant height (97.37 cm), [n](#)Number of nodules/plant (112.60), [p](#)Plant dry weight (18.41 g/plant), [p](#)Pods/plant (31.2), [s](#)Seeds/pod (2.20), [f](#)Fest weight (38.46 g), [p](#)Pod yield (3.00 t/ha), [s](#)Seed yield (2.46 t/ha), [h](#)Haulm yield (4.94 t/ha), [h](#)Harvest index (49.83 %), [s](#)Shelling percentage (82.13 %), gross returns ([Rs.98,508.00/ha](#)), net return ([Rs.71,388.00/ha](#)) and benefit cost ratio (2.63). [\(Maybe it is better to quote in dollars, because it is an international article\)](#)

Key words: *Rhizobium*, PSB, Zinc sulphate, ~~Groundnut~~[\(It is better choose another word because is the same in the title\)](#).

INTRODUCTION

The peanut (*Arachis hypogaea L.*), also known as the groundnut, goober (US), (Domonoske 2014) pindar (US) or monkey nut (UK), is a Legume crop grown mainly for its edible seeds. It is widely grown in the tropics and subtropics, being important to both small and large commercial producers. It is classified as both a grain legume and, due to its high

oil content, an oil crop. World annual production of shelled peanuts was [44 million tonnes in 2016, led by China with 38% of the world total](#)[\(Cite here the reference of this information\)](#). Atypically among legume crop plants, peanut pods develop underground rather than above ground. With this characteristic in mind, the botanist Carl Linnaeus gave peanuts the specific epithet

Formatted: Font color: Red

Formatted: Font color: Red

Formatted: Font: Font color: Red

hypogaea, which means "under the earth". The peanut belongs to the botanical family Fabaceae, commonly known as the legume, bean, or pea family. Like most other legumes, peanuts harbor symbiotic nitrogen-fixing bacteria in root nodules. The capacity to fix nitrogen means peanuts require less nitrogen-containing fertilizer and improve soil fertility, making them valuable in crop rotation.

Zinc is required in various metabolic processes as catalysts. Zinc also increases the content of protein, calorific value, amino acid and fat in oilseed crop. Balanced fertilization helps to improve the quality of the produce. Zinc entered into the constituents of enzyme system that regulate initial metabolic reactions in the plant body. Zinc catalyses the process of oxidation in plant cells and is vital for the transformation of carbohydrates. It regulates the consumption of sugars and increases the source of energy for the production of chlorophyll. Zinc also aids in the formation of auxin and synthesis of protein. [\(Cite here the reference of this information\).](#)

Bio-fertilizers are known to play a number of vital roles in soil fertility, crop productivity and production in agriculture as they are eco-friendly but cannot at any cost replace chemical fertilizers that are indispensable for getting maximum crop

yields. They supplement chemical fertilizers for meeting the integrated nutrient demand of the crops. Application of bio-fertilizers results in increased mineral and water uptake, root development, vegetative growth and nitrogen fixation. Bio-fertilizers offer an economically attractive and ecologically sound means of reducing external inputs and improving quality and quantity of crop. They contain microorganisms which are capable of mobilizing nutrient elements from unavailable form to available form through different biological processes **(Hadiya et al. 2017)**. *Azotobacter* inoculants when applied to many non-leguminous crop plants, promote seed germination and initial vigor of plants by producing growth promoting substance **(Kalita et al. 2019)**.

[The purpose of the work was missing. Write here](#)

MATERIAL AND METHODS:

The experiment was conducted to know the effect of different doses of bio-fertilizers and foliar application of zinc on growth and yield attributes of Groundnut cv. K1812 was carried out at Crop Research Farm of Sam Higginbottom University, Prayagraj, Uttar Pradesh during 2022 [\(Cite here the period \(months\) of the experiment\)](#). The experiment was laid out in a [Randomized Block Design \(RBD\)](#) consisting of ten treatments including [Control](#) with 3

Formatted: Font: Italic

replications, [\(what is the plot size? Number of plants per plot?\)](#)with the treatment combinations

T₁: *Rhizobium* 20g/kg seeds + 0.25% ZnSO₄,

T₂: Rhizobium 20g/kg seeds + 0.50% ZnSO₄,

T₃: Rhizobium 20g/kg seeds + 0.75% ZnSO₄,

T₄: [\(describe the meaning of PSB only once\)](#)PSB 20g/kg seed + 0.25% ZnSO₄,

T₅: PSB 20g/kg seed + 0.50% ZnSO₄,

T₆: PSB 20g/kg seed + 0.75% ZnSO₄,

T₇: Rhizobium 10g/kg seeds + PSB 10g/kg seeds + 0.25% ZnSO₄,

T₈: Rhizobium 10g/kg seeds + PSB 10g/kg seed + 0.50% ZnSO₄,

T₉: *Rhizobium* 10g/kg seeds + PSB 10g/kg seeds + 0.75% ZnSO₄,

T₁₀:Control.

[It is missing the described statistical analysis used in this work! Statistical tests.](#)

RESULTS AND DISCUSSION:

PARAMETERS:

PRE-HARVEST:

The perusal of data indicate that **plant height** measured at final stage (i.e., 60 DAS) was not influenced markedly by the application of different levels of phosphorous, though numerical increase in plant height was recorded in dose dependent manner. At 60 DAS, significantly higher plant height (53.55 cm)

was recorded with the treatment PSB 20g/kg seed + 0.75% ZnSO₄. However, the treatments PSB 20g/kg seed + 0.50% ZnSO₄ (53.45 cm), PSB 20g/kg seed + 0.25% ZnSO₄ (52.75 cm) and *Rhizobium* 20g/kg seeds + 0.75% ZnSO₄ (52.25 cm) and Rhizobium 10g/kg seeds + PSB 10g/kg seeds + 0.75% ZnSO₄ (53.35 cm) were found to be statistically at par with PSB 20g/kg seed + 0.75% ZnSO₄ and there was significant difference among the treatments. However, highest Number of nodules/plants (112.60) was recorded with the application of PSB 20g/kg seed + 0.75% ZnSO₄. At 60 DAS, there was significant difference among the treatments. However, highest dry weight (28.47 gm) was recorded in treatment T₆.

POST - HARVEST:

Significantly Maximum Number of Pods/plant (31.2), Seeds per Pod (2.20) and test weight (38.46 gm) were recorded with the treatment of application of PSB 20g/kg seed + 0.75% ZnSO₄ over all the treatments.

DISCUSSIONS:

Muhammad (2011)observed **M**maximum plant height (88.00cm) ~~was observed~~ in the inoculated plants, while less plant height (56.13cm) was recorded for uninoculated plants [Groundnut plants? Cite](#). The reason for this might be the adequate amount of nitrogen, fixed by the *Rhizobium*~~Rhizobia~~.

Formatted: Font: Italic

Formatted: Font: Italic, Font color: Auto

Formatted: Font: Italic

Formatted: Font: Font color: Red

Formatted: Font color: Red

Formatted: Font: Italic

More number of leaves per plant may also be positively contributed to more plant height in the inoculated plants. The mean values for cultivars also show significant effect for plant height. Maximum plant height (80.43 cm) was achieved by cv. Chakori, while minimum plant height (64.04 cm) was observed in cv. KS-1. Comparing means for cultivars for plant height, it is evident that variation observed in plant height was probably due to the difference in the degree of adaptation to the environment of experimental site.

Rhizobium inoculation treatments also gave superior results to the un-inoculated ones in plant height; stem dry weight, number of nodules/plant and nodule dry weight. Potdaret *al.* (2019) concluded that application of 60 kg P₂O₅ ha⁻¹, FYM (Needs to describe the meaning) 5 t ha⁻¹, and PSB + VAM (Needs to describe the meaning) 20 g/kg seed inoculum increased the number of pods/plants. (281,20), seeds/pods (10.95), test weight (4.35 g), seed yield 1956 kg/ha, Indian mustard (*Brassica juncea* L.) vs oil yield of each control (40.2%). Kalitaet *al.* (2019) in their experiment reported that seed inoculation with *Azotobacter* and PSB 40 g kg⁻¹ seed + 75% NPK in toria (*Brassica campestris* L.) recorded maximum grain yield (11.15 qha⁻¹ is the same as kg ha⁻¹?) due to the siliquae plant⁻¹ (164.76), root growth (2.30 g plant⁻¹), seeds siliquae⁻¹(10.97) and seed weight (4.82 g).

CONCLUSION:

It is concluded that application of PSB 20g/kg seed + 0.75% ZnSO₄ was recorded significantly higher seed yield (2.46 t^{ha}⁻¹), and benefit cost ratio (2.63) as compared to other treatments. Since, the finding based on the research done in one season (This phrase needs to messed up, it is confused and it is better rewrite).

REFERENCES:

- Domonoske, Camila (April 20, 2014). "A Legume with Many Names: The Story Of 'Goober'". NPR. National Public Radio. Archived from the original on June 6, 2020.
- Hadiyal, J.G., Kachhadiya, S. P., Ichchhuda, P. K. and Kalsaria, R. N. 2017. Response of Indian mustard (*Brassica juncea* L.) to different levels of organic manures and biofertilizers. *Journal of Pharmacognosy and Phytochemistry*. 6(4): 873-875.
- Meena, J., Singh, V., Kumar, S. and Sagar, A. 2018. Effect of Biofertilizers and Levels of Sulphur on Growth and Yield attributes of Mustard [*Brassica juncea* (L.) Czern.&Coss]. *International Journal of Current Microbiology and*

Formatted: Superscript

Formatted: Font color: Red

Formatted: Font: Italic

Formatted: Font color: Red

Formatted: Font color: Red

Formatted: Font color: Red

Formatted: Font color: Red

Formatted: Superscript

Applied Sciences; **7** (10): 2242-2249. (Check if it is mentioned in the text)

Sajid, M., A. Rab, F. Wahid, S.N.M. Shah, I. Jan, M. A. Khan, S.A. Hussain, M.A. Khan and Z. Iqbal. 2010. Influence of rhizobium inoculation on growth and yield of groundnut cultivars. *Sarhad J. Agric.* **27**(4): 573-576.

Kalita, N., Bhuyan, S., Maibangsa, S. and Saud, R. K. 2019. Effect of Biofertilizer seed treatment on

Muhammad

(2011)???)

Growth, yield and economics of toria (*Brassica campestris*L.) under rainfed condition in hill zone of

Assam. *Current Agriculture Research Journal*, **7**(3); 332-336.

Potdar, D. S., Purohit, H. S., Meena, R. H., Kaushik, M. K., Jain, H. K. and Ameta, K. D. (2019). Effect of integrated phosphorus management on growth, yield and quality of mustard (*Brassica juncea* L.). *Journal of Pharmacognosy and Phytochemistry*, **8** (4): 1700-1704.

missing citation

Formatted: Font color: Red

Table 1: Effect of biofertilizers and foliar application of Zinc on growth of Groundnut at 60 days after sowing.

Treatments	Plant Height (cm)	No. of Nodules	Dry weight (g)	Pods/plant(No)	Seed/pod(No)	Test weight (g)
T1	50.25	108.18	24.03	26.0	1.73	30.24
T2	51.70	109.39	25.22	26.2	1.73	30.24
T3	52.25	109.80	26.12	27.1	1.80	31.46
T4	52.75	110.18	26.32	28.2	1.87	32.69
T5	53.45	111.38	28.00	30.3	2.00	37.00
T6	53.55	112.60	28.47	31.2	2.20	38.46
T7	52.95	110.43	26.75	28.2	1.87	32.69
T8	52.75	110.74	26.98	28.5	1.93	33.74
T9	53.35	111.29	27.14	29.0	2.00	34.96
T10 (Control)	48.30	105.70	23.08	24.1	1.73	30.24
F- Test	S? or ns?	S	S	S	S	S
SEm(±)	0.72	1.218	0.365	0.43	0.03	0.53
CD(p=0.05)	2.15	3.61	1.056	1.29	0.09	1.58

[Describe the meanings of acronyms. Example. SE= ???; CD = ???; S = ??; Check the journal rules.](#)

If for the statistical analysis of the data the test of means between treatments was performed, it is necessary to insert the letters next to the numerical values.

Formatted: Indent: First line: 0"

UNDER PEER REVIEW