

# Antitrypanosomal efficacy of *Annona muricata* leaf extracts in *Trypanosoma brucei brucei* experimentally infected rats

## ABSTRACT

**Aims:** Trypanosomiasis is an important protozoan disease that affects domestic animals and man. It is caused by the tsetse fly-transmitted extracellular hemo-flagellates that belong to the genus, *Trypanosoma*. In East and Southern Africa, Human African trypanosomiasis (HAT) is caused by the tsetse fly-transmitted hemo-flagellates *Trypanosoma brucei rhodesiense* while in West and Central Africa, it is caused by *T. b. gambiense*. Animal trypanosomiasis on the other hand is caused by *T. b. brucei*, *T. vivax*, and *T. congolense*. The incidence of sleeping sickness as a major health problem in many African countries has been on the rise in recent years. In sub-Saharan Africa, about sixty million people are at risk of infection. This current study evaluates the antitrypanosomal efficacy of extracts of *Annona muricata* leaf in *Trypanosoma brucei brucei* infected albino rats.

**Place and Duration of Study:** Animal parasitology and microbiology research unit, Department of Animal production and Health, Federal University of Technology, Akure, Nigeria, between February and June, 2021.

**Methodology** *In vitro* antitrypanosomal analysis was done in varied concentrations of 2.5mg/ml, 5mg/ml and 10mg/ml using various solvent extracts (ethanolic, ethyl acetate, n-hexane, chloroform and aqueous). Diminazine aceturate and normal saline were used as positive and negative controls respectively. The *in vivo* assay was carried out through intraperitoneal administration of graded doses (200, 400 and 600mg/kg) of ethyl acetate and chloroform extracts of the plant for three consecutive days.

**Results:** The n-hexane, chloroform and ethyl acetate extracts yielded high percentage DPPH free radical scavenging activity of 85.52, 80.00 and 78.49% respectively. Decrease in motility of the parasites at different times were observed in all extracts tested *in vitro*. These responses were positively concentration-dependent. 10mg/ml concentration of chloroform and ethyl acetate showed complete cessation of parasite motility at 35 and 45 minutes respectively. These two extracts (ethyl acetate and chloroform extracts) which showed the best *in vitro* responses, were subjected to *in vivo* analysis. Both extracts caused decrease in trypanosome parasitemia and prolongation of mean survival days of the rats to 14.67days as compared with 6.83days in the negative control group. The extracts displayed dose-dependent significant ( $p \leq 0.05$ ) antitrypanosomal activities when compared with the negative and positive controls.

**Conclusion:** The chloroform and ethyl acetate extracts of *Annona muricata* showed a relatively higher antitrypanosomal activity over other solvent extracts used in this study. Further fractionation, purification and isolation should be done to confirm the active components in this plant that is responsible for the antitrypanosomal activities recorded.

**Keywords:** *Annona muricata*, antitrypanosomal activities, solvent extracts, *Trypanosoma brucei brucei*,

## 1. INTRODUCTION

Trypanosomiasis, though neglected sub-Sahara African tropical infectious disease, its medical and veterinary importance cannot be overemphasized. It is caused by a protozoan

parasite of the genus *Trypanosoma*, transmitted by the tsetse fly [3] within the distributional limits. *Trypanosoma* species *T. brucei brucei* responsible for African Animal Trypanosomiasis (AAT) called Nagana in West Africa, *T. b. rhodesiense* and *T. b. gambiense* cause Human African Trypanosomiasis (HAT) or sleeping sickness [1, 2]

African animal trypanosomiasis is a parasitic disease which results in serious economic losses in livestock following anemia, lethargy and effects on reproduction with more prominence in cattle. Animals other than livestock, including dogs, can also be affected as untreated cases can be fatal, with very high mortality rate in some outbreaks. In domestic animals, Trypanosomiasis caused by *T. brucei*, *T. equiperdum* and *T. evansi* pose a great socio-economic influence as well as global decreased reproductive efficiency [4]. With about 57 million people are at risk of infection [5], Trypanosomiasis can be zoonotic with symptoms such as anemia, fever, weakness and weight loss. Some proposed interventions to mitigate the disease's effect include the World Health Organization (WHO) programs to eliminate Trypanosomiasis as a public health issue by 2020 and interrupt its transmission by 2030 [5].

Bumduuren *et al.*, [6], suggests *Trypanosoma* species are transmitted not only by tsetse flies (*Glossina* spp.) but by other biting insects. Trypanosomiasis control in Africa has been mainly through chemotherapy and chemoprophylaxis, coupled with vector control programmes, as there has not been any reported effective vaccine against the disease, with diminazene and isometamidium chloride as the main drug control of animal trypanosomiasis in Africa. Their continued use and effectiveness have been threatened by serious limitations, ranging from high cost, long-course of parenteral administration, serious side effects, varied efficacy and emergence of drug resistant trypanosome strains [7]. The presence of drug resistant trypanosomes has recently risen to alarming proportions [8], hence Osho and Lajide [9] supports the search for new chemical entities which are effective against trypanosomes, safe yet affordable for disease-endemic countries is rational.

Countless number of secondary molecules that have pharmacological effects present in plants increases their chances as potential sources of new drugs [9,10] which are pharmacologically active and can provide an alternative to chemically synthesized drugs to which many infectious microorganisms have become resistant [11]. Exploring traditionally appraised medicinal plant for the biological activity has provided humankind with a number of antiprotozoal medications which pastoralists all over the world, have used in old tradition for the treatment of several animal diseases, including trypanosomiasis.

*Annona muricata* Lin., commonly called soursop, is part of the Annonaceae family. *A. muricata* L. contains various compounds with pharmacological activity. This plant is widely grown in tropical and subtropical areas such as Southeast Asia, South America, and the rainforests of Africa. The plant produces edible fruit all year round and is widely used as a traditional medicine for skin disease, respiratory disease, fever, bacterial infections, diabetes, hypertension, and cancer [12].

Validation of medicinal plant for their antitrypanomal activity will direct the society on the best approach to employ their indigenous knowledge and at the same time provide hit compounds to feed future regime for antitrypanomal drug development [13]. This study therefore provides a valid scientific proof for the invitro and invivo therapeutic efficacy of extracts of *Annona muricata* against *Trypanosoma brucei brucei* infected wistar albino rats.

## **2. MATERIAL AND METHODS**

### **2.1 Collection of plant leaves**

Leaves of *Annona muricata* Linn (Annonaceae) were harvested locally during the daytime from their natural habitats in Akure, Ondo State, Nigeria. They were authenticated by the Department of Pharmacognosis, Faculty of pharmacy, Obafemi Awolowo University, Ile ife with voucher number FPI 2400.

## **2.2 Plants extraction**

Fresh leaves of *Annona muricata* were air-dried and crushed with the aid of a mechanical grinder to powdery form. Extraction was done using five different solvents (chloroform, ethanol, hexane, ethyl acetate and distilled water). 200 g dried weight of powdered sample of leaf was soaked in 1.0 liter of individual solvents in order to extract the bioactive components that are present in the plant samples. The sample-solvent combination was agitated at hourly intervals for 72 hours after which mixtures were filtered. The resultant extracts were lyophilized using rotary evaporator (R110) at 40°C to obtain a dry powder extract. Percentage yield of each plant extracts were calculated. The respective extracts obtained were stored in a sealed container and refrigerated until use.

## **2.3 Phytochemical Analysis of Plant Extracts**

Extracts were subjected to quantitative and qualitative phytochemical screening. Methods described by Trease and Evans [14] and Ejikeme *et al.*, [15] were adopted to detect the absence or presence of secondary metabolites such as; saponins, tannins, alkaloids, flavonoids, and phenol in the plant.

## **2.4 DPPH free radical scavenging activities**

Based on the scavenging activities of the stable 1,1-diphenyl 2-picrylhydrazyl (DPPH), the free radical scavenging activity of the extracts were determined adopting the method described by Villano *et al.*, [16].

## **2.5 Trypanosomes and experimental animals**

*Trypanosoma brucei brucei* was obtained from the National Institute of Trypanosomiasis Research, Vom, Jos, Nigeria. Serial passaging of the parasite was done in a donor rat. Parasitaemia of 'donor rat' was estimated using "rapid matching" method. [17].

## **2.6 Experimental rats' acclimatization**

Both male and female healthy adult wistar albino rats of whose weight ranges between 130-180g were procured from the Animal facility of Obafemi Awolowo University Ile Ife, Osun State, Nigeria. They were housed in clean dry cages and fed standard pellets and watered ad libitum. They were kept for seven (7) days to allow them acclimatise and cared for following the international guidelines for the use and maintenance of experimental animals [18].

## **2.7 Evaluation of invitro antitrypanosomal activity**

Invitro test was carried out in triplicates to assess the motility of trypanosomes using a 96 well microplate. 20µL of blood containing about 16-32 organisms per field were mixed with 5µL of the test substance at concentrations of 2.5, 5, 10mg/mL to produce test concentrations of 0.5, 1, 2.0mg/mL, respectively. The standard trypanocidal drug, diminazene aceturate (DA) and phosphate buffer saline (pH 7.2) were used to serve as

treated and untreated controls, respectively. At 37°C, the mixtures were incubated for up to 2 hours. Motility of the parasites was checked in intervals of 5 minutes under microscope (X40 objective lens). In summary, about 2 µL of test mixtures was placed on microscope slide and covered with cover slips. The parasites were observed for reduced motility or complete cessation of motility. Chloroform and ethyl acetate extracts yielded the best results in this invitro experiment and were further subjected to invivo analysis.

## 2.8 Evaluation of invivo antitrypanosomal activities

Forty-eight wistar albino rats of both sexes weighing between 130g-180g were used for the invivo phase of this study. The animals were randomly divided into eight groups (I-VIII) consisting 6 rats each. All groups were infected intraperitoneally with 0.2ml of *T. brucei* (at 7.2 antilog) suspension. Groups I-III were treated with chloroform extracts while groups IV-VI were treated with the ethyl acetate extracts of *Annona muricata* at daily doses of 200, 400 and 600mg/kg body weight for three consecutive days intraperitoneally. Group VII was administered Diminacene aceturate DA at 3.35mg/kg as a single dose intraperitoneally to serve as treated control. 0.3 mL normal saline was administered intraperitoneally to Group VIII for three consecutive days to serve as untreated control. Treatment was initiated on fourth day post inoculation in all groups. Body weight, parasitemia, and rectal temperature were monitored daily for 21 days [19].

Parasitemia was monitored through examination of blood from the tail of rats under microscope at × 40 magnifications using the “Rapid Matching” method as illustrated by Herbert and Lumsden 1979. Parasitemia was monitored on a daily basis until the 21st day from the onset of treatment [20, 21].

Body weight of experimental animals were recorded on the day of parasite inoculation and everyday thereafter for 21 days [22].

With the aid of rectal thermometer (Mettler Toledo, Switzerland), rectal temperature was recorded on the day of parasite inoculation and everyday thereafter for 21 days [23].

## 3. RESULTS

### 3.1 Yield

Ethanol gave the highest yield followed by n-hexane, chloroform, aqueous and ethyl acetate in that order.

**Table I: Percentage yield with the different solvents of extraction**

S/N	EXTRACTS	% YIELD
1	EEAM	13.61
2	EAAM	18.11
3	NHAM	15.31
4	AEAM	13.59
5	CEAM	16.30

EEAM: Ethanolic extract of *A. muricata*

EAAM: Ethyl acetate extract of *A. muricata*

NHAM: N- hexane extract of *A. muricata*

AEAM: Aqueous extract of *A. muricata*  
 CEAM: Chloroform extract of *A. muricata*

### 3.2 DPPH free radical scavenging activities of extracts of *Annona muricata* leaves

N-Hexane extract gave the highest DPPH free radical scavenging percentage followed by chloroform, ethyl acetate, ethanolic and aqueous extracts in that order

**Table II: Percentage DPPH free radical scavenging activities with the different solvents of extraction**

Sample	DPPH (%)
EEAM	77.38
EAAM	78.49
NHAM	85.52
AEAM	55.56
CEAM	80.00

EEAM: Ethanolic extract of *A. muricata*  
 EAAM: Ethyl acetate extract of *A. muricata*  
 NHAM: N- hexane extract of *A. muricata*  
 AEAM: Aqueous extract of *A. muricata*  
 CEAM: Chloroform extract of *A. muricata*

### 3.3 Phytochemical screening

Phytochemical screening revealed the presence of saponins, tannins, phenol, flavonoids and alkaloids in varying quantities with the different extraction solvents used (Table III and IV).

**Table III: Qualitative phytochemical analysis of extracts of *Annona muricata* leaves**

Sample	Alkaloids	Saponin	Tannin	Flavonoid	Terpenoids	Steroid
EEAM	+	+	+	+	-	+
EAAM	++	++	+	+	+	+
NHAM	++	++	+	+	+	+
AEAM	+	++	±	+	±	+
CEAM	++	+	+	+	+	+

Foot note:

+: Present      - : Absent      +: Slightly present      ++: Highly present

**Table IV: Quantitative phytochemical analysis of extracts of *Annona muricata* leaves**

Sample	Alkaloids %	Saponin %	Flavonoids mg/100g	Phenol mg/100g	Tannin mg/100g
EEAM	6.98	9.59	7.64	3.43	5.00
EAAM	27.77	16.40	3.14	3.78	5.08
NHAM	21.03	26.49	3.86	3.68	4.63
AEAM	3.50	32.08	6.64	0.07	0.08
CEAM	23.16	7.89	2.50	3.11	3.68

### 3.4 *Invitro antitrypanosomal activity*


Chloroform and ethyl acetate extracts of *A. muricata* leaves ceased motility of the trypanosomes within 40 minutes at 10mg/ml concentrations (Table V).

**Table V: Effect of different extracts of *Annona muricata* on motility cessation time of *Trypanosoma brucei brucei* invitro**

Conc.	Time (min) After Which Motility Ceased						DA	NC
	EEAM	EAAM	NHAM	AEAM	CEAM			
10mg/ml	75	40	65	95	30	15	NE	
5mg/ml	80	55	70	105	35	20	NE	
2.5mg/ml	95	65	80	115	60	35	NE	

### 3.5 *Effect of treatment on parasitemia*

Parasitemia level assessed at day 5 post-treatment in all experimental animals for the different groups showed that all doses of the extract including the standard drug suppressed parasitemia ( $p < 0.05$ ). The best result was obtained with chloroform extract of *A. muricata* at 400mg/kg on day 11 (Figure I).



**Figure I: Effect of *A. muricata* on parasitaemia in rats infected with *T. brucei brucei***

Notes

AMC1: Chloroform extract of *A. muricata* at 200mg/kg

AMC2: Chloroform extract of *A. muricata* at 400mg/kg  
 AMC3: Chloroform extract of *A. muricata* at 600mg/kg  
 AME1: Ethyl acetate extract of *A. muricata* at 200mg/kg  
 AME2: Ethyl acetate extract of *A. muricata* at 400mg/kg  
 AME3: Ethyl acetate extract of *A. muricata* at 600mg/kg  
 DM : Diminazene aceturate at 3.35mg/kg  
 INT: Infected not treated

### 3.6 Effect of treatment on body weight

There is statistically significant ( $p < 0.05$ ) body weight changes in 600mg/kg and DA 3.35mg/kg treated groups compared with 200mg/kg, 400mg/kg, treatment groups with both chloroform and ethyl acetate extracts from day 8 to day 14 post-treatment initiation and with untreated control on day 5 post-treatment (Figure II).




Figure II: Effect of *Annona muricata* leave extracts on weight of *T. brucei* infected rats

### 3.7 Effect of treatment on mean survival days

The average survival days of infected rats were increased to 13.33 and 14 days in 600mg/kg of ethyl acetate and chloroform extracts respectively compared with 6.83 days in negative control group (Figure III).




Figure III: Effect of *A. muricata* leaf extracts on mean survival days of rats infected with *T. brucei brucei*

### 3.8 Effects of treatment on rectal temperature

The rectal temperatures of the rats fluctuated throughout the duration of the experiment independent of the environmental temperatures. There is no observable significant difference nor pattern.

## 4. DISCUSSION

The affinity of chemical composition of substances for the different solvents of extraction is responsible for variations observed in yield of extracts. Solubility of compounds tend to be higher in polar solvents such as methanol in comparison with non-polar solvent such as chloroform [24]. Nguyen et al., [25] also reported variation in yield of *Annona muricata* seeds using distilled water, ethanol, ethyl acetate and chloroform, where chloroform gave the highest yield.

Different roles of phytochemicals have been reported ranging from plants' protection from harmful insects, ultraviolet (UV) rays, microbes and extreme temperatures. They also play important roles in man and animals. Most phytochemicals have been observed to display antioxidant activities as they help in mopping up free radicals and also protect against diseases that manifest as a result of free roaming reactive oxygen species (ROS) [26, 27, 28]. Phytochemicals have been documented to help prevent and ameliorate diseases such as diabetes, cancer, hyperlipidaemia, Alzheimer, cardiovascular diseases, liver toxicity, cataract, age related function decline, stroke and others. Various phytochemicals that have been isolated from plants include alkaloids, glycosides, saponins, flavonoids, kaempferol, phytol, gallic acid, kolaviron and essential oils [29].

The qualitative and quantitative phytochemicals obtained from both plants differ with the various solvent used for the extraction. Significant pharmacological functions such as; antimalarial, anticancer, analgesic, and anti-hyperglycemic and antibacterial functions have been reported for alkaloids [30]. Steroids are used to relief inflammatory conditions while terpenoids have antioxidant activities. Saponins have been evidenced to possess

anticoagulant, anti-carcinogenic, hypoglycemic, immunomodulator, neuroprotective, anti-inflammatory and antioxidant potentials. Glycosides have antimicrobial and anticancer activities. [31].

Traditional uses of *Annona muricata* in the treatment of various diseases can be credited to the availability of these constituents in its leaf

Antitrypanosomal and therapeutic properties of some medicinal plants have been associated with the significant levels of various classes of phytochemicals compounds or secondary metabolites present in them. The leaf extract of *Annona muricata* contains alkaloids, saponins, tannins, phlobatannins, and flavonoids. These phytochemicals have been reported earlier to be present in the seeds of *A. muricata* [32].

The decrease in body weight of experimental rats over a period might be due to anorexia which is one of the clinical features of trypanosomosis. This can further be described by the inverse relationship between level of parasitaemia and weight loss of gain in the treatment groups; a relationship that is positively dose dependent.

Chloroform extracts of *A. muricata* exhibited higher antitrypanosomal activities than its ethyl acetate extract both invitro and invivo. This fact may suggest the concentration of the plant's pharmacological active ingredients responsible for its antitrypanosomal activities by chloroform solvent to a large extent. Diminazene aceturate treatment group displayed the highest antitrypanosomal activities in the infected rats because the parasite has been tested sensitive to diminazene being a standard antitrypanosomal drug. The effectiveness of diminazene aceturate at very low dosages is because its purity when compared with the crude extracts of *Annona muricata* that were used in this study in which relapse was recorded.

## 5. CONCLUSION

The plant antitrypanosomal activities may be associated to the presence of active secondary metabolites in the leaf and this affirms its use in Nigerian local communities towards the treatment of parasitic infection. The chloroform and ethyl acetate extracts showed a relatively higher activity over other solvent extracts used in this study.

The extracts of *A. muricata*, in this study can be beneficial to human and animal health in the treatment of trypanosomosis.

This study provides a lead for researchers to uncover the specific bioactive components of *Annona muricata* that exerts the antitrypanosomal efficacy.

## CONSENT

Not applicable

## ETHICAL APPROVAL

Not applicable

## REFERENCES

1. D'Archivio S, Medina M, Cosson A, Chamond N, Rotureau B, Minoprio P. Genetic engineering of *Trypanosoma* (*Dutonella*) *vivax* and in vitro differentiation under axenic

conditions. PLoS Negl Trop Dis. 2011;5(12):e1461.  
<https://doi.org/10.1371/journal.pntd.0001461>.

2. Cayla M, Rojas F, Silvester E, Venter F, Mathews KR. African trypanosomes. *Parasit. Vectors*. 2019;12:190. <https://doi.org/10.1186/s13071-019-3355-5>.
3. WHO. Interim guidelines for the treatment of gambiense human African trypanosomiasis. Geneva: World Health Organization; 2019. Licence: CC BY-NC-SA 3.0 IGO. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK545514/>
4. Desquesnes M, Holzmüller P, Lai D, Dargantes A, Lun Z, Jittaplapong S. *Trypanosoma evansi* and surra: a review and perspectives on origin, history, distribution, taxonomy, morphology, hosts, and pathogenic effects. *Biomed Res Int*. 2013: 10.1155
5. Franco JR, Simarro PP, Diarra A, Jannin J. Epidemiology of human African trypanosomiasis. *Clin Epidemiol*. 2014;6:257-275. doi: 10.2147/CLEPS39728. pmid:25125985
6. Bumduuren T, Thillaiampalam S, Naoaki Y, Ikuo I. Development of unstable resistance to diminazene aceturate in *Babesia bovis*. *Int J Parasitol Drugs. Drug Resist*. Apr; 9: 87–92. doi: 10.1016/j.ijpddr. 2019.02.001 PMID: PMC6382846
7. Rasool-Hassan BA. Medicinal plants (importance and uses). *Pharmaceut Anal Acta*. 2012;3:139. <https://doi.org/10.4172/21532435.1000e139>.
8. Burri C. Chemotherapy against human African trypanosomiasis: is there a road of success? *Parasitology*. 2010;137:1987–94.
9. Osho IB, Lajide L. Prescreening evaluation of some plant extracts used in ethno-veterinary practices as antitrypanosomal agents. *Journal of Medicinal Plants Research* 2012;6(11):2056-2060, Available online at <http://www.academicjournals.org/JMPR> DOI: 10.5897/JMPR10.045 ISSN 1996-0875.
10. Liu WJH. Traditional herbal medicine research methods: identification, analysis, bioassay, and pharmaceutical and clinical studies. Hoboken: Wiley. 2011;477.
11. Akinpelu DA, Abioye EO, Aiyegoro OA, Akinpelu OF, Okoh AI. Evaluation of antibacterial and antifungal properties of *Alchornea laxiflora* (Benth.) Pax. & Hoffman. *Evid-Based Compl Alternat Med* 2015: 1 - 6.
12. Moghadamtousi SZ, Fadaeinasab M, Nikzad S, Mohan G, Ali HM, Kadir HA. *Annona muricata* (Annonaceae): A Review of Its Traditional Uses, Isolated Acetogenins and Biological Activities. *Int. J.Mol. Sci*. 2015,16, 15625–15658.
13. Ayawa NG, Ramon-Yusuf SB, Wada YA. Toxicity study and antitrypanosomal activities of aqueous and methanol whole plant extracts of *Brillantaisia owariensis* on *Trypanosoma brucei* infection in BALB/c mice. *Clin Phytosci* 2012;7, 39. <https://doi.org/10.1186/s40816-021-00267-3>.
14. Trease and Evans. W. C. Trease and Evans Pharmacognosy (14th Edition), W. B Saunders Company Limited, London, 2005;357-358. 13.
15. Ejikeme CM, Ezeonu CS, Eboatu AN. Determination of physical and phytochemical constituents of some tropical timbers indigenous to Niger Delta Area of Nigeria. *European Scientific Journal*. 2014;10(18);247–270.
16. Villano D, Fernandez-Pachon MS, Moya ML, Troncoso AM, Garcia-Parrilla MC. Radical scavenging ability of polyphenolic compounds towards DPPH free radical, *Talanta* 2007;71; 230.
17. Herbert WJ, Lumsden WH. *Trypanosoma brucei*: A rapid “matching” method for estimating the host’s parasitemia. *Exp Parasit*; 1976;40: 427- 431.
18. OECD (Organization for Economic Cooperation and Development) revised draft guidelines 423. OECD Guidelines for the testing of chemicals; 2001.
19. Ene A, Atawodi S, Ameh D, Nnamani C, Apeh Y. Antitrypanosomal effects of petroleum ether, chloroform and methanol extracts of *Artemisia maciverae* Linn. *Indian J Exp Biol*. 2009;47:981–6.

20. Maikai V. Antitrypanosomal activity of avonoid extracted from *X. americana* stem bark. *Int J Biol*;2011;1:115–21.
21. Angela O, Mustapha L, David MY. Anti-trypanosoma activity of the ethanolic leaf extract of *Senna occidentalis* (Fabaceae) on *T. brucei brucei* infected mice. *Int J Basic Appl Sci*;2013; 2:32–7.
22. Nweze N, Anene B, Asuzu I. Investigation of the anti-trypanosomal activity of *Buchholzia coriacea* seed extract against a eld strain of *T. congolense*. *Afr J Tradit Complement Altern Med*;2011;8:175– 80.
23. Ngulde S, Tijjani M, Ihopo J, Ya'uba M. Anti-trypanosomal potency of methanolic extract of *Cassia arereh* Delile root bark in albino rats. *Int J Drug Res Tech*. 2013;3:1–7.
24. Oladunmoye MK, Kehinde FY. Ethnobotanical survey of medical plants used in treating viral infections among Yoruba tribe of South Western Nigeria. *African. Journal of Microbiology Research*. 2011;5(19):2991–3004.
25. Omoya FO, Oloruntuji AB. Prophylactic and Curative Potency of *Alchornea laxiflora* Extract on *Plasmodium berghei* Infected Swiss Albino Mice *International Journal of Tropical disease & Health* 2019;36(1): 1-11. Article no.IJTDH.48675 ISSN: 2278–1005, NLM ID: 101632866.
26. Mith H, Dure R, Delcenserie V, Zhiri A, Daube G, Clinquart A. Antimicrobial activities of commercial essential oils and their components against food-borne pathogens and food spoilage bacteria. *Food Sci. Nutr.*, 2014;2: 403-416.
27. Tilaoui M, Mouse HA, Jaafari A, Ziad A. Comparative phytochemical analysis of essential oils from different biological parts of *Artemisia herba alba* and their cytotoxic effect on cancer cells. *Plos One*, 2015;10.
28. Oksana S, Marian B, Mahendra R, Bo SH. Plant phenolic compounds for food, pharmaceutical and cosmetics production. *J. Med. Plants Res.*, 2012;6: 2526-2539.
29. Swamy MK, Akhtar MS, Sinniah UR. Antimicrobial properties of plant essential oils against human pathogens and their mode of action: An updated review. *Evidence-Based Complementary Altern. Med.*, 2016.
30. Osabiya OJ, Bada SO, Ibitoye FO, Aribisala OK. Assessment of physical constituents, antibacterial and cytotoxic activities of *A. laxiflora* (Ewe pepe) extracts. *Journal of Applied Life Sciences Intl*. 2017;12(2):1-9.
31. Rao AV, Gurfinkel DM. The Bioactivity of saponins: Triterpenoid and steroidal glycosides. *Drug Metab Drug Inter*. 2000;17(1-4): 211–235.
32. Nguyen VT, Nguyen MT, Tran QT, Thinh PV, Bui LM, Le THN, Le VM, Linh HTK. Effect of extraction solvent on total polyphenol content, total flavonoid content, and antioxidant activity of soursop seeds (*Annona muricata* L.). *Energy Security and Chemical Engineering Congress IOP Conf. Series: Materials Science and Engineering*. 2020. IOP Publishing doi:10.1088/1757-899X/736/2/022063