

# Artificial Propagation of Indigenous Ornamental Fish Species via Nutritional Manipulation for Stock Enhancement and Conservation as a Step towards Environmental Sustainability: A Review

---

## ABSTRACT

Indigenous flora and fauna of a particular habitat greatly impacts the environment in maintaining an ecological balance and equilibrium for the normal functioning of the ecosystem as a whole. Native fishes play a major role and holds a major niche in the ecosystem which needs attention at the moment for their conservation and protection, failure of which may prove to be detrimental to the environment. The rapid developmental activities and industrialization taking place currently are leaving a negative impact upon the natural ecosystem. These activities are a major reason for loss of habitat, deterioration, pollution, and degradation. The ornamental fish industry is a budding industry in India and the indigenous fish fauna of the country offers a good market but unfortunately, so far, the requirements and demand are fulfilled from the natural stock through manual collection. Rigorous exploitation of these species may result into loss of species diversity or reduction of stock, provided, anthropogenic causes necessitated by development and industrialization are already acting upon the ecosystem. Therefore, concrete measures such as development of captive breeding techniques should be standardized for such species so that the wild stock is not made vulnerable and instead, the same shall be enhanced through ranching and this shall bring us nearer towards the goal of sustainability. Development of captive breeding techniques may be carried out by domesticating the selected species in indoor facilities, manipulate the habitat in a way that will mimic its natural habitat, provide optimum nutrition by supplementing the best available feeding matter and eventually under controlled and favorable conditions maturation and breeding may be achieved successfully.

*Keywords: Biodiversity, Captive breeding, Conservation, Habitat, Nutrition, Sustainability.*

## 1. INTRODUCTION

Fish are a vital component of our ecosystems and are key to preserving the harmony of the natural world. Particularly native fish species are essential to the general health of aquatic ecosystems and are an essential component of their own habitats. But because of human activities like overfishing, habitat destruction, and the introduction of invasive species, many native fish species are in danger of going extinct. Therefore, protecting these species is

essential to ensuring the survival of our aquatic environments and the advantages they offer. For several reasons, it is essential to preserve native fish species. Fish species that are native to the area are a vital source of food for other creatures that live there. The survival of several predators, such as birds, animals, and other fish, depends on the presence of local fish species. Additionally, by consuming and excreting organic matter, native fish help to distribute nutrients throughout aquatic ecosystems, where they play a crucial role in the nutrient cycle. Furthermore, in their respective habitats, these native fish species frequently serve as keystone species. Keystone species are those that, in relation to their abundance, have an outsized influence on their ecosystem. An ecosystem can be severely disrupted by the removal of a keystone species, which may have a domino effect on other species. Because of their function in controlling the numbers of other animals and preserving the ecosystem's balance, native fish species are frequently keystone species in aquatic ecosystems. Therefore, it is crucial to preserve native fish species in order to keep these ecosystems ecologically balanced.

Native to India, ornamental fish comprise 700 marine species and 374 freshwater species. Of the marine species, 150 occurs in Andaman & Nicobar Islands and 300 in Lakshadweep Islands [1]. The demand for these fish is very high in foreign export markets. Freshwater biodiversity is abundant in NER and the Western Ghats, in contrast to marine biodiversity, which is mostly found in Lakshadweep, Andaman & Nicobar Islands, Gulf of Kutch, Gulf of Mannar, Palk Bay, Kerala coast, etc. 90% of ornamental fish marketed originates from the port of Kolkata, with the remaining 8% and 2% coming from Mumbai and Chennai, respectively [2]. In both domestic and foreign markets, demand is limited to the local fish *Sahyadria denisonii*. Even though native ornamental fish from India are highly sought-after on international markets, [3] noted that only a small amount is exported for a variety of reasons.

Maintaining biodiversity depends on protecting native fish species. Biodiversity is the term used to describe the diversity of life on Earth, including the range of species, genes, and ecosystems. Native fish species are a crucial component of this biodiversity, and their extinction could have a big effect on how resilient and healthy an ecosystem is. Additionally, protecting indigenous fish species can aid in limiting the spread of invasive species, which can harm native ecosystems and species. The most important concern is sustainability, and breeding native species that are unpopular in the domestic market is likewise a low priority. Even though the nation has made scientific advancements in the breeding techniques for a few native ornamental fishes, their mass production has not yet begun. If government organizations can set up substantial facilities and provide breeders with specialized instruction and assistance, more indigenous ornamental fishes can be produced to increase exports from the country. According to the research done by [4], native fish species found in India, specifically the native fish fauna found in the Ganges and its tributaries, prefer shallow, clear water streams with a sandy substratum, along with the presence of gravels and substantial growths of aquatic plants like *Hygrophilia*, *Limnophila*, *Ottelia*, *Rotala*, and *Vallisneria*. And if we do a reality check, we might discover that natural water bodies are gradually losing quality as a result of ongoing industrial and development activities, endangering the abundance, existence, and biodiversity of indigenous species, and ultimately having a negative effect on the environment. Therefore, the prime need of the hour is to work on this aspect by developing captive breeding techniques for these important species and once achieved, the seeds can be ranches for stock enhancement and traders may acquire the seeds from breeders rather than collecting from the wild which will enormously reduce the pressure upon the natural ecosystem.

## **2. IMPORTANCE OF CAPTIVE BREEDING FOR PROPAGATION OF INDIGENOUS ORNAMENTAL FISHES**

In order to lessen the strain placed on nature by wild-catch and preserve the natural population of these living treasures, [1] have stressed the importance of developing captive broodstock for indigenous ornamental fish species. According to [5], both climatic and human-caused factors have contributed to the loss of fish variety. Additionally, according to their study, 63 species that were present in the Hakaluki Haor wetland in north-eastern Bangladesh were still available in 83% of cases in 2008, but a drop of 51% was noted in 2018. The wild populations of fish are at great risk due to overfishing and habitat destruction happening due to ongoing developmental activities, which is gradually reducing the species' natural supply.

Furthermore, [6] reported that only 15% of native fish species are raised in captivity, out of the total indigenous ornamental fish trade that takes place in India. If technological advancements in captive breeding are not made, this could ultimately result in over-exploitation, endangerment, and eventually lead to the extinction of native fish species, provided, climatic and anthropogenic factors are already acting upon this to create a synergistic effect.

## **3. HABITAT MANIPULATION FOR BREEDING OF ORNAMENTAL FISHES IN CAPTIVITY**

To speed up the maturation process, it is crucial to mimic the natural environment as closely as possible in captivity. This is accomplished by supplying a thicket of plants [7]. Males being territorial in nature require objects for hideouts to enable them to create distinct boundaries for their specific territories, which may include the use of plants, woods, or even mud pots and a substratum basically formed by laying gravel or sand. The presence of floating plants can be advantageous in the way that it blocks excess light which might negatively impact the breeding performance of the fish. A dense cushion of moss that is ideally 4 inches (10 cm) deep and without gaps is also very effective [8]. The fish feel at home in this habitat, which also serves as a shelter and an initial supply of food for the brood. Eventually, this improves the fish's ability to reproduce. For the fish to mature in captivity, Dutta *et al.*, 2020, underlined the need to provide a semi-natural setting. Light maintenance from low to moderate is recommended for enhancing plant growth.

## **4. NUTRITIONAL REQUIREMENTS FOR REPRODUCTION OF ORNAMENTAL FISHES**

Enhancing the external appearance and reproductive performance of invertebrates, including fish, and finally determining the reproductive outcome in them are all influenced by nutrition, which is one of the fundamental factors. However, nutritional information for ornamental fish is primarily based on extrapolating findings from research on food fish produced in intensive farming environments [9, 10]. In their thorough investigation of the effects of dietary supplementation on fish reproduction, [9], noted that the reproduction process can only achieve success, if necessary, conditions are met; however, it may also lead to reproductive failure if devoid of optimal conditions. For example, when female fishes are malnourished, they frequently exhibit serious reproductive malfunctions which even lead to the occurrence of follicular atresia conditions and inhibition of other mechanisms in the system of the fish, which evidently will lead to the complete failure of breeding of the fish.

Due to the challenges of conducting studies concerning the optimal feeding and reproduction of broodstock, the nutrition of broodstock is currently little known [11, 12]. The types of feed had a substantial impact on *Betta splendens* feed intake, growth, ovary weight, and fertility

[13]. Enhancing reproductive success necessitates adequate nutrient availability as per the requirement of the fish to provide and maintain the calorific requirements for the growing cells, which are important factors in the design and final acceptance of aquatic feeds [14, 9]. An otherwise nutritionally balanced meal may function poorly or ineffectively if it contains only a small amount of nutrients that cause a specific aquatic species to react positively to stimuli [14]. Fishes in the natural ecosystem frequently experience food scarcity or shortages throughout their life cycles, especially during seasonal variation and migratory phase. In response to these intrinsic and extrinsic effects, the sensory and endocrine systems, react, and the reproduction process is then carried out based on the signal produced [9].

#### **4.1. Protein requirement**

According to [15], ornamental fishes need a crude protein content between 30 and 40% to mature. To promote reproductive activities, female freshwater ornamental swordtails, *Xiphophorus helleri*, need a diet with at least 30% protein [12]. Feeding *B. splendens* with a 35% CP diet resulted in higher growth and spawning performance [16]. Protein requirements can range from 30 to 50% for growing fishes based on the type of which wherein omnivorous goldfish, *Carassius auratus*, requires the former level and carnivorous discus, *Symphysodon aequifasciata* requires the latter [17]. According to [18], for guppy, protein requirements range from 30 to 40 % for initial sizes of 0.10 g.

Furthermore, [19] revealed that most ornamental fish need diets with 25–50% protein, while [20] found that the highest fertility in female dwarf gouramis was reported from dietary supplementation of 25–45% protein. Additionally, [21] observed that *X. helleri* females fed with a very high quantity of protein (> 30%) increased their relative fecundity and produced more fry. According to [22], gourami brooders need 30–40% protein content, with the primary protein sources being fish meal, squid meal, etc. Blue gourami, *Trichogaster trichopterus*, was fed with various protein and lipid levels, and [23] reported that treatment with 35 % CP and 8 % EE produced the best results. It was also noted that blue gourami, swordtail, molly, and guppy require a 30-40% dietary protein supplementation.

#### **4.2. Fatty acids and lipids requirement**

The lipid and fatty acid composition of the diet of broodstock is thought to be one of the key nutritional elements, useful for effective breeding performance and the survival of spawns [9]. Dietary lipids are prime sources of energy and fatty acids, thereby playing a significant role in the regular growth and survival of fishes [17]. In addition to playing a crucial role in the flavour and texture of fish feed, lipids also play a crucial role in maintaining the structure of biological membranes, are carriers of hormones, effectively transport carotenoids and deposits in chromatophores, and also act as precursors for the synthesis of essential metabolites, including prostaglandins [24]. Fish mainly need long-chain fatty acids consisting of high levels of unsaturation as reported by [17]. A lipid content of 9–10% is added to feed for the Nile tilapia broodstock development [25].

According to [21], muscle lipid content serves as an ovarian lipid supply, making it a useful measure of reproductive function. Swordtail displayed enhanced growth performance with dietary fat increases from 8 to 16% and the same protein level. With the largest accumulation being shown with the highest dietary fat, the muscle lipid content followed the same trajectory as the protein level. According to [26], 7-8% lipid was included in the diet of milkfish broodstock. According to [23], blue gourami, swordtail, molly, and guppy need a 6–10% dietary lipid supplement. A direct incorporation of highly unsaturated fatty acids such as DHA, EPA, and ARA, to female broodstock diets, has been shown to directly benefit the reproduction of zebrafish, *Danio rerio* [27]. Fish with higher levels of some important fatty acids and vitamins, including n-3 and n-6 HUFA, vitamin E, vitamin C, and carotenoids, have better reproductive performance and better gamete and larvae quality [28]. To get the

desired outcomes, it is essential to include these ingredients in tiny amounts in broodstock diets.

#### **4.3. Carbohydrate requirement**

One of the most vital elements in all living beings is carbohydrates, commonly known as sugars or saccharides. They serve as rapidly metabolized energy source molecules that help the body transmit energy, and structural elements. There is no evidence that fish has a demand for carbs in its diet. However, because they are a cheap source of energy, carbohydrates can "spare" the catabolism of other substances like protein and lipids [17]. According to [9], lower limits of carbohydrate incorporation do not have a significant effect on maturation and breeding.

#### **4.4. Vitamin-Mineral Supplements**

While some organisms cannot synthesize vitamins, which are organic compounds needed in relatively small amounts by most forms of life and contribute to oogenesis and spermatogenesis in aquatic animals, minerals are inorganic compounds required by fish for the formation of tissue as well as aids in numerous metabolic and regulatory processes [17, 9]. Vitamin E and C supplements were given to milkfish brooders, and they produced more spawns with greater percentages of viable eggs, hatching, and cumulative survival rates [26]. Additionally, [26] made a point of mentioning the requirement for vitamin C incorporation to increase the number of spawns with high-quality eggs and larvae. In rainbow trout, vitamin C has been linked to vitellogenesis and has been found to sustain sperm quality [29]. Fishes are incapable of naturally synthesizing vitamin E in their system, therefore, an external supplementation via incorporation in feedstuffs is an important step as it determines fish fertility and reproduction, as stated by [30] and [9]. Diet deficient in vitamin E inhibited ovarian growth in common carp, *Cyprinus carpio* [31], and it is also responsible for egg quality.

For their essential physiological and biochemical activities as well as to maintain their regular life processes, all aquatic species need minerals [32]. The presence of calcium in the water medium is necessary as it activates and hardens the eggs, as the eggs come into contact with water, and embryonic formation requires phosphates which are attained through vitellogenin in the yolk. Low female fecundity, hatchability, and high rates of abnormal deformity are consequences of phosphorus deficiency [9].

### **5. IDEAL DIET FOR GONADAL MATURATION AND REPRODUCTION**

According to [13], in majority of Asian countries, broodstock development is still carried out with the aid of live foods which are the primary sources of optimum nutrition, apart from other minor supplementation, such as feeding with a paste-based diet prepared from fish meal and skimmed milk powder. The use of formulated meals in place of live food is essential for reducing production costs and maintaining the generation of juveniles of high and consistent quality [33]. Furthermore, according to [34], freshwater fish production has been constrained due to the scarcity of live food. Formulated diets would mitigate this issue and enhance a consistently high output of high-quality seeds.

According to [13], the composition and palatability of meals had a substantial impact on *B. splendens* reproduction. The availability of proper type of feed is crucial since, according to [16], the short reproductive cycle of ornamental fish may cause continuous oogenesis in the adult female. Some species, like the voracious and indiscriminate feeder ruby barb (*Puntius nigrofasciatus*), have less preference for artificial diets, and highly accept and prefer live feed [35]. Breeders, involved in large-scale production of Discus (*S. aequifasciata*), feed the

developing fry primarily with live foods including Tubifex, blood worms, and *Artemia* nauplii as reported by [36].

Furthermore, the dissociation of nitrogenous compounds from feedstuffs during protein metabolism (mostly ammonia) would immediately damage their existing habitat and cause stress in the fish, therefore this fact needs to be taken into consideration and ornamental fish kept in confinement must use their dietary protein as effectively as possible, hence the selection of an ideal diet is necessary [37]. Protein and lipid content of discussed ingredients is given in Table 1.

**Table 1. Protein and lipid content of feed ingredients**

| <b>Ingredient</b>  | <b>CP (%)</b> | <b>CL (%)</b> | <b>Reference</b> |
|--------------------|---------------|---------------|------------------|
| Tubifex            | 59.70         | 17.50         | [13]             |
| Fish meal          | 65.42         | 5.58          | [39]             |
| Squid meal         | 72.19         | 10.19         | [47]             |
| Chicken liver meal | 60.37         | 23.60         | [48]             |

\*Abbreviation – CP: crude protein, CL: Crude lipid

### **5.1. Tubifex**

Tubifex's crude protein and crude lipid contents are high enough, i.e., 59.7% and 17.5%, respectively [13], which will facilitate easy maturation and enhanced reproductive performance in fishes. [38] mentioned that tubifex is a nutritionally complete diet for maturation and reproduction. Tubifex is utilized as a gourami broodstock feed because it has a good amount of iron, carotenoids, and haemoglobin [22].

### **5.2. Fish Meal**

Fish Meal is a good source of nutrients for fish maturation, with 65.42 % crude protein and 5.58 % lipid [39]. Fish meal is the best source for supplying the proper nutrition in terms of growth, feed conversion, and gonadal development, according to the report of [18] on guppy, *P. reticulata*.

### **5.3. Squid Meal**

The presence of free amino acids in squid meal makes it one of the most potent attractants that pique aquatic animals' gustatory (taste) and olfactory (smell) senses. Its recommended inclusion level in broodstock diet is 20–30%. [40], [41] also defined squid meal as a good source of dietary lipids and protein. Glycine and betaine, two chemo-attractant qualities of squid meal and other squid products, are crucial stimulators of the feeding behaviour [42]. Biological research has revealed that squid contains some undiscovered growth factors, which are referred to as "squid factor," and squid meal appears to have additional growth-promoting qualities [43, 44].

When squid meal was added to the diet of Kuruma shrimp (*Penaeus japonicus*), growth was increased by 30 to 50%. Animals' serum cholesterol is reduced by the squid's lipid content [45]. Furthermore, 30% of the maturation diets for tiger shrimp broodstock contain both fresh and squid meal [46]. Squid meal is a feasible source of nutrients for fish growth and maturation, with a crude protein level of 72.19 % and a lipid content of 10.19 % [47].

### **5.4. Chicken Liver Meal**

A very good source of nutrients for the maturity and reproductive performance of ornamental fishes is chicken liver meal, which has a crude protein content of 60.37% and a lipid content of 23.6% [48]. The best results were shown in swordtail fed a diet that included chicken liver meal in terms of weight increase, SGR, and FCR [23]. In their study, [49] found that feeding black molly a diet including chicken liver meal produced better results than feeding fish waste meal, earthworm meal, etc. Additionally, the performance of chicken liver meal is comparable to that of squid meal and prawn meal.

## 6. THE IMPACT OF SELECTIVE MEAL TYPE ON GROWTH AND SURVIVAL

SGR in larval rearing of *Chitala chitala* was found to be higher in post-hatchlings fed on live tubifex worms as measured by [50] in their study to compare different diet performances, including fish eggs, dry tubifex, chironomid larvae, spirulina, daphnia, and planktons. [51] found that young South American catfish surubim (*Pseudoplatystoma fasciatum*) fed with tubifex had the best survival rate (100%) and the highest level of 20:4n-6 deposition in the fish tissue. When compared to alternative diets like formulated feed, vermicompost worms, etc., tubifex-fed angel fish showed the highest weight gain [52]. In terms of weight increase, SGR, and FCR, [53] found that goldfish fed a diet based on fish meal showed improved growth response and feed performance. However, Discus and Red Head Cichlid fed a diet based on fish meal with 40–50% protein content displayed superior SGR [54, 55].

According to [47], adding 4% squid meal to de-hulled solvent extracted soybean meal as a secondary source of protein will effectively act as an attractant and boost the intake of plant-based diet for pompano. According to [23], goldfish fed squid meal had the second-highest growth performance after fish-fed tubifex, and their cost per unit of production was about 20 times lower than that of tubifex. In addition, [23] found that feeding blue gourami (*T. trichopterus*) and guppy (*P. reticulata*) squid meal and chicken liver resulted in the best growth and feed performance as measured in terms of weight gain, SGR, FCR, and PER, whereas feeding swordtail (*X. helleri*) with chicken liver meal resulted in the best performance.

## 7. CONCLUSION

In conclusion, local fish species are essential for preserving the harmony and health of aquatic ecosystems. These species need to be conserved for a variety of reasons, such as their value as a food source, their part in preserving ecosystems' ecological balance, their cultural relevance, and their contribution to biodiversity. To protect these species and assure their survival for future generations, action must be taken. This can be done by taking a variety of steps, including as habitat restoration, sustainable fishing methods, and the control of invasive species. Enhancing the fish population's natural stock can be accomplished through captive breeding. Captive breeding can be used to supplement natural populations and boost their size in light of the decline of many fish species brought on by overfishing, habitat degradation, and other human influences. Genetic variety may be preserved in fish by controlled mating, and the progeny can then be released into the wild to aid in the recovery of wild populations. Furthermore, captive breeding can aid in the production of fish with desirable characteristics, such as disease resistance, which can aid in boosting the general health and resilience of wild populations. Overall, improving the natural fish resource and aiding in the conservation of these significant species can be accomplished through captive breeding.

## REFERENCES

1. Pandey PK, Mandal SC. Present status, challenges and scope of ornamental fish trade in India. In: Conference: Aqua Aquaria India, At Mangalore. 2017.
2. Sekharan N. Market trends in Indian ornamental fish trade, INFOFISH International, 2017, 42-44. 3/2017, [www.infofish.org](http://www.infofish.org)
3. Raja K, Aanand P, Padmavathy S, Sampathkumar JS. Present and future market trends of Indian ornamental fish sector. International Journal of Fisheries and Aquatic Studies. 2019;7(2):6-15.
4. Hamilton F. An account of the fishes found in the river Ganges and its branches. Edinburgh: A. Constable and Company; 1822.
5. Aziz MSB, Hasan NA, Mondol MMR, Alam MM, Haque MM. Decline in fish species diversity due to climatic and anthropogenic factors in Hakaluki Haor, an ecologically critical wetland in northeast Bangladesh. Heliyon. 2021;7(1). p.e05861.
6. Mahapatra BK, Lakra WS. Ornamental fishes of East Kolkata wetland, West Bengal, India. Zoology. 2014;3(12):406-408.
7. Mahadevi N, Felix S, Ahilan B, Rajagopalasamy CBT. Evaluation of reproductive performance in indigenous endemic ornamental fish *Sahyadria denisonii* using hormones under captive environment. Journal of Environmental Biology. 2020;41(3): 549-555.
8. Kullander SO, Britz R. Revision of the family Badidae (Teleostei: Perciformes), with description of a new genus and ten new species. Ichthyol Expl Freshw. 2002;13 (4):295–372.
9. Singh SK, Baidya S, Das P, Biswas P. Functional Role of Dietary Supplements on Reproductive Physiology of Fishes. In: Sundaray JK, Rather MA, Kumar S, Agarwal D, editors. Recent updates in molecular Endocrinology and Reproductive Physiology of Fish (pp. 243-258). Springer, Singapore; 2021.
10. Keshavanath P, Patil P. Nutrition in ornamental fishes. Fishing chimes. 2006;26(8):13-18.
11. Izquierdo MS, Fernandez-Palacios H, Tacon AGJ. Effect of broodstock nutrition on reproductive performance of fish. Aquaculture. 2001;197(1-4): 25-42.
12. Chong AS, Ishak SD, Osman Z, Hashim R. Effect of dietary protein level on the reproductive performance of female swordtails *Xiphophorus helleri* (Poeciliidae). Aquaculture. 2004;234(1-4):381-392.
13. Mandal SC, Sahu NP, Kohli MPS, Das P, Gupta SK, Munilkumar S. Replacement of live feed by formulated feed: effect on the growth and spawning performance of Siamese fighting fish (*Betta splendens*, Regan, 1910). Aquaculture Research. 2010;41(11):707-1716.
14. Meyers SP. Utilization of shrimp processing waste. INFOFISH marketing digest. 1986; 4:18-19.
15. Lewbart GA. Clinical nutrition of ornamental fish. In: Seminars in Avian and Exotic Pet Medicine. WB Saunders. 1998;7(3): 154-158.
16. James R, Sampath K. Effects of meal frequency on growth and reproduction in the ornamental red swordtail, *Xiphophorus helleri*. Isr. J. Aquac. Bamidgeh. 2003;55(3): 197–207.
17. Sales J, Janssens GP. Nutrient requirements of ornamental fish. Aquatic Living Resources. 2003;16(6): 533-540.
18. Shim KF, Chua YL. Some studies on the protein requirement of the guppy, *Poecilia reticulata* (Peters). Journal of Aquaculture and Aquatic Science. 1986;4: 79-84.
19. Yanong RP. Nutrition of ornamental fish. Veterinary Clinics of North America: Exotic Animal Practice. 1999;2(1): 19-42.
20. Shim KF, Landesman L, Lam TJ. Effect of dietary protein on growth, ovarian development and fecundity in the dwarf gourami *Colisa lalia* (Hamilton). J. Aquat. Trop. 1989;4:111-123.

21. Ling S, Hashim R, Kolkovski S, Chong Shu-Chien A. Effect of varying dietary lipid and protein levels on growth and reproductive performance of female swordtails *Xiphophorus helleri* (Poeciliidae). *Aquacult Res.* 2006;37(13): 1267-1275.
22. Jena AK, Biswas P, Pattanaik SS. Recent trends in breeding and trade of ornamental gourami in India. *World Aquaculture.* 2019;61.
23. Mohanta KN, Subramanian S. Nutrition of Common Freshwater Ornamental Fishes. Technical Bulletin No: 27, ICAR Research Complex for Goa (Indian Council of Agricultural Research), Old Goa - 403 402, Goa, India; 2011.
24. National Research Council. Nutrient Requirements of Warmwater Fishes. National Academy Press, Washington, DC, USA; 1977.
25. El-Sayed AFM, Mansour CR, Ezzat AA. Effects of dietary lipid source on spawning performance of Nile tilapia (*Oreochromis niloticus*) broodstock reared at different water salinities. *Aquaculture.* 2005;248(1-4): 187-196.
26. Emata AC, Borlongan IG, Damaso JP. Dietary vitamin C and E supplementation and reproduction of milkfish *Chanos chanos* Forsskal. *Aquaculture Research.* 2000;31(7): 557-564.
27. Jaya-Ram A, Kuah MK, Lim PS, Kolkovski S, Shu-Chien AC. Influence of dietary HUFA levels on reproductive performance, tissue fatty acid profile and desaturase and elongase mRNAs expression in female zebrafish *Danio rerio*. *Aquaculture.* 2008;277(3-4): 275-281.
28. De Silva SS, Anderson TA. Fish nutrition in aquaculture (Vol. 1). Springer Science & Business Media; 1994.
29. Ciereszko A, Liu L, Dabrowski K. Effects of season and dietary ascorbic acid on some biochemical characteristics of rainbow trout (*Oncorhynchus mykiss*) semen. *Fish Physiology and Biochemistry.* 1996;15(1): 1-10.
30. Mandal SC, Singh SK, Das P, Barman D, Gupta SK. Trace and heavy metal content in diets and their effect on the growth and survival of Siamese fighting fish, *Betta splendens*. *Indian Journal of Animal Nutrition.* 2013;30(1): 80-86.
31. Watanabe T, Takashima F. Deficiency Symptoms and Changes of Fatty Acid and Triglyceride Distributions in Adult Carp. *Bulletin of the Japanese Society of Scientific Fisheries.* 1977;43(7): 819-830.
32. Lall SP, Kaushik SJ. Nutrition and metabolism of minerals in fish. *Animals.* 2021;11(09): 2711
33. Cahu C, Infante JZ. Substitution of live food by formulated diets in marine fish larvae. *Aquaculture.* 2001;200(1-2): 161-180.
34. Lim LC, Dhert P, Sorgeloos P. Recent developments in the application of live feeds in the freshwater ornamental fish culture. *Aquaculture.* 2003;227(1-4): 319-331.
35. Weerasooriya AC, Edirisinghe U, Silva P, Dematawewa CMB. Feeding behaviour and acceptability of different feeds by *Puntius nigrofasciatus*. *Trop. Agric. Res.* 1999;11: 221-228.
36. Chong A, Hashim R, Ali AB. Assessment of soybean meal in diets for discus (*Symphysodon aequifasciata* HECKEL) farming through a fishmeal replacement study. *Aquaculture Research.* 2003;34(11): 913-922.
37. Earle KE. The nutritional requirements of ornamental fish. *Veterinary Quarterly.* 1995;17(sup1): 53-55.
38. Tamaru CS, Ako H, Paguirigan R. Essential fatty acid profiles of maturation feeds used in freshwater ornamental fish culture. *Hydrobiologia.* 1997;358(1): 265-268.
39. Nasim Al Mahmud MD, Hossain MB, Minar MH. Proximate composition of fish feed ingredients available in Lakshmipur region, Bangladesh. *Am. J. Agric. Environ. Sci.* 2012;12: 556-560.
40. Hertrampf JW, Piedad-Pascual F. Handbook on ingredients for aquaculture feeds. Springer Science & Business Media; 2003.

41. Vassallo-Agius R, Watanabe T, Imaizumi H, Yamazaki T, Satoh S, Kiron V. Effects of dry pellets containing astaxanthin and squid meal on the spawning performance of striped jack *Pseudocaranx dentex*. *Fisheries Science*. 2001;67(4): 667-674.
42. Xue M, Cui Y. Effect of several feeding stimulants on diet preference by juvenile gibel carp (*Carassius auratus gibelio*), fed diets with or without partial replacement of fish meal by meat and bone meal. *Aquaculture*. 2001;198(3-4): 281-292.
43. New MB. Feed and feeding of fish and shrimp. A manual on the preparation and presentation of compound feeds for shrimp and fish in aquaculture. UNDP/FAO/ADCCP/REP/87/26, Rome, Italy; 1987.
44. Guillame J, Cruz-Ricque E, Cuzon G, Warmhoudt A van, Revol A. Growth factors in penaeid shrimp feeding. *Aquacop Ifremer Actes de Colloque*. 1989;9: 327-338.
45. Stansby ME. Fish oils in nutrition (Vol. 313). New York: Van Nostrand Reinhold; 1990.
46. Millamena OM, Pudadera RA, Catucatan MR. Effects of diet on reproductive performance of ablated *Penaeus monodon* broodstock. In: Proceedings of the First International Conference on the Culture of Penaeid Prawns/Shrimps, 4-7 December 1984, Iloilo City, Philippines (pp. 178-179). Aquaculture Department, Southeast Asian Fisheries Development Center; 1985.
47. Novriadi R, Spangler E, Rhodes M, Hanson T, Davis DA. Effects of various levels of squid hydrolysate and squid meal supplementation with enzyme-treated soy on growth performance, body composition, serum biochemistry and histology of Florida pompano *Trachinotus carolinus*. *Aquaculture*. 2017;481: 85-93.
48. Mohanta KN, Subramanian S, Korikanthimath VS. Evaluation of different animal protein sources in formulating the diets for Blue Gourami, *Trichogaster trichopterus* Fingerlings. *Journal of Aquaculture Research and Development*. 2013;4(2): 1-7.
49. Pai IK, Altaf MS, Mohanta KN. Development of cost effective nutritionally balanced food for freshwater ornamental fish Black Molly (*Poecilia latipinna*). *Journal of Aquaculture Research and Development*. 2016;7:401.
50. Sarkar UK, Lakra WS, Deepak PK, Negi RS, Paul SK, Srivastava A. Performance of different types of diets on experimental larval rearing of endangered *Chitala chitala* (Hamilton) in recirculatory system. *Aquaculture*. 2006;261(1): 141-150.
51. Arslan M, Dabrowski K, Portella MC. Growth, fat content and fatty acid profile of South American catfish, surubim (*Pseudoplatystoma fasciatum*) juveniles fed live, commercial and formulated diets. *Journal of Applied Ichthyology*. 2009;25(1): 73-78.
52. Jayalekshmi JN, Abraham KM, Sobhanakumar K. Growth performance of Angelfish, *Pterophyllum scalare* fed with different live worm diets. *Journal of Aquatic Biology & Fisheries*. 2017;5: 116-12.
53. Lochmann RT, Phillips H. Dietary protein requirement of juvenile golden shiners (*Notemigonus crysoleucas*) and goldfish (*Carassius auratus*) in aquaria. *Aquaculture*. 1994;128(3-4): 277-285.
54. Chong ASC, Hashim R, Ali AB. Dietary protein requirements for discus (*Symphysodon* spp.). *Aquaculture Nutrition*. 2000;6(4): 275-278.
55. Olvera-Novoa MA, Gasca-Leyva E, Martinez-Palacios CA. The dietary protein requirements of *Cichlasoma synspilum* Hubbs, 1935 (Pisces: Cichlidae) fry. *Aquaculture Research*. 1996;27(3): 167-166.