

Influence of Foliar Application of Zinc on Growth and Yield in Different Varieties of Rice (*Oryza sativa* L.) and Yield Validation Using SPSS Model

ABSTRACT

Agronomy Department, Farm, Sam Higginbottom University of Agriculture, Technology, and Sciences, Prayagraj (U.P.), India. to research the effects of zinc treatment on rice (*Oryza sativa* L.) growth and yield. The medications include the varieties MTU-7029, BPT -5204, and RNR15048, as well as zinc concentrations of 0.2%, 0.3%, and 0.4%. The pH of the soil in the experimental plot was 7.8, the texture was sandy loamy, and the amount of organic carbon was low (0.35%). The larger plant height (117.68 cm), greater number of tillers per hill (13.64), Application of RNR-15048 + ZnSo₄ - 0.4% significantly increased plant dry weight (55.47 g/plant), number of panicles/hill (10.01), number of grains/panicle (141.37), 1000 seed weight (22.27 gm), grain yield (6.43 t/ha), and stove yield (12.11 t/ha). Higher B:C ratio (2.37) and higher gross return (INR 1,52,860/ha), net return (INR 1,07,531/ha), and net return (INR 1,07,531/ha) were also noted in treatment-9 (RNR-15048 + ZnSo₄ - 0.4%).

Keywords: Rice; varieties; zinc; growth parameters; yield attributes and economics.

1. INTRODUCTION

The most important cereal food crop in India is rice (*Oryza sativa* L.), which accounts for around 24% of the nation's total cropped land. It contributes 42% of the nation's overall production of food grains and 45% of the nation's total production of cereals. After China, India is the country with the largest rice-growing area and the second-highest rice production [1].

About 31% of India's GDP is derived from agriculture, and 25% of its exports are composed of agricultural goods. More than 65% of people rely on it as a staple meal, so the expansion and stability of its production are crucial to the safety of our country's food supply. The biggest area of rice cultivation is in India, where there are 44.6 million hectares of rice planted in four primary eco-systems: irrigated, rain fed lowland, rain fed upland, and flood prone, which take up 21, 14, 6, and 3 million hectares, respectively, and produce about 90 million tonnes of rice annually. Saha et al. [2].

With the introduction and cultivation of semi-dwarf, fertilizer-responsive, and non-inns moderate yielding varieties in the early 1970s, which was crucial to the "Green Revolution," rice

production and productivity significantly improved. Yatheesh and others, 2022). The output and quality of rice, which is utilised for feed and industrial uses, were significantly improved by rice hybrids. If the management level is greater than 60%, hybrid rice growing is commercially viable. Hybrids are non-lodging, have considerable pest and disease resistance, are short-lived, respond well to stress and various climatic conditions, and have a longer shelf-life. Out of the 43 million hectares that are being grown for rice, 3 million of those are hybrids, according to Sarkar SC [3]. Because of the pressing need for high yielding rice types due to population growth, rice hybrids break yield barriers and produce 15-20% more.

Around 50% of farmed soils worldwide have low Zn concentrations during cereal-growing seasons [4]. Zn deficiency in plants is characterised by decreased membrane integrity, increased vulnerability to thermal stress, and decreased synthesis of auxin, chlorophyll, cytochromes, carbohydrates, and nucleotides.

A few of the Zn-containing enzymes that are inhibited include alkaline phosphatase, carbonic anhydrase, Cu-Zn-superoxide dismutase, phospholipase, carboxypeptidase, and RNA

polymerase [5]. Zinc interacts with more than 500 different kinds of proteins. Chelated zinc (EDTA) Zinc sulphate heptahydrate, applied as a foliar fertiliser and in the soil, increased the zinc content of plant parts like hull, bran, white rice kernels, and straw [6]. Foliar fertilisation has a stronger effect on raising Zn concentration in rice than soil application, according to the rice crop's immediate response to nutrient treatment. The production of cereal and straw from different types of rice differed substantially when Zn was applied and when it wasn't. Grain yields of diverse rice kinds increased by 29% and 22% on average with soil + foliar and merely soil applications of Zn, respectively (shah et al 2013).

These factors were taken into consideration as the current study, "Foliar application of zinc on growth and yield in different varieties of rice (*Oryza sativa* L.) and yield validation using SPSS model", was carried out at the Kharif season of 2022 at the Crop Research Farm, Department of Agronomy, Naini Agriculture Institute, Sam Higginbottom University of Agriculture Technology and Sciences, Prayagraj, Uttar Pradesh.

2. MATERIALS AND METHODS

At the Crop Research Farm, Department of Agronomy, Naini Agriculture Institute, Sam Higginbottom University of Agriculture Technology and Sciences, Prayagraj, Uttar Pradesh, the experiment was carried out during the Kharif of 2022. which is situated 98m above mean sea level (SL) at 25.24' 42" N latitude, 81.50' 56" E longitude. Ten treatments, each replicated three times, were used in the experiment's Randomised Block Design. Each treatment's plot was 3m x 3m in size. There are three levels of zinc factors[ZnSo4 -0.2% (30,60 DAS), ZnSo4 -0.3% (30,60 DAS), ZnSo4 -0.4% (30,60 DAS)], and three different rice varieties (MTU-7029, BPT-5204, RNR-150480). The different levels of zinc supplied during 30 and 60 DAS. The Rice varieties were sown on 26 June 2022 by maintaining a spacing of 22.5 cm x 10 cm. Each plot was used for 1m² of the harvest. Five plants were then at random chosen from it to record the yield and growth parameters. The treatment details are as follows, T₁-(MTU-7029 + ZnSo4- 0.2%), T₂ -(MTU-7029 + ZnSo4- 0.3%), T₃ – (MTU-7029 + ZnSo4- 0.4%), T₄ -(BPT -5204 + ZnSo4- 0.2%), T₅ -(BPT -5204 + ZnSo4- 0.3%), T₆ -(BPT -5204 + ZnSo4- 0.4%), T₇ -(RNR15048 + ZnSo4- 0.2%), T₈ -(RNR15048 + ZnSo4- 0.3%), T₉ -(RNR15048 + ZnSo4- 0.4%), T₁₀-(N 120 Kg/ha + P 60 kg/ha +k

60 kg/ha) Control. Plant height, dry weight, number of tillers per hill, crop growth rate, number of panicles per hill, number of grains per panicle, test weight, grain yield, and straw yield were all observed. By using the analysis of variance approach, the data were statistically analysed [7].

3. RESULTS AND DISCUSSION

3.1 Growth Parameters

3.1.1 Plant height

At 100 DAS, treatment-9 (RNR-15048 + ZnSo4 - 0.4%) had the noticeably taller plant height (117.68 cm). However, statistically speaking, treatment-8 (RNR-15048 + ZnSo4 -0.3%) was comparable to treatment-9 (RNR-15048 + ZnSo4 -0.4%). Different zinc levels over the control had a substantial impact on plant height, a key growth metric. In comparison to the other treatments, plants at 0.4% were found to be noticeably taller. In comparison to the other treatments, plants at 0.4% were found to be noticeably taller. High levels of photosynthetic and hormonal activity led to meristematic growth in the apical region, which caused the plants to elongate sufficiently as a result of the application of Zn. According to Ghatak et al. [8] and Haque et al. [9], a significant contributing aspect is the genetic makeup of the variety. The synchronised availability of all important plant nutrients, particularly nitrogen for a longer period during growth phases, may also contribute to an increase in plant height [10].

3.1.2 Number of tillers/hill

Treatment-9 (RNR-15048 + ZnSo4 - 0.4%) had the considerably larger number of tiller/hill (13.64) than the other treatments. However, statistically speaking, treatment-8 (RNR-15048 + ZnSo4 -0.3%) was comparable to treatment-9 (RNR-15048 + ZnSo4 -0.4%). The application of zinc may be the cause of the much more tillers per hill. The dynamics of zinc in soil may have been improved by zinc-enriched urea and Zn-FYM incubation to extend its availability for enhanced rice growth and tillering. Superiority of Zn-EDTA. Gangloff et al. [11]. Additionally, the likely cause of high tillering capacity in high yielding varieties. Yadav et al. [1] similarly reported similar results. According to Wang et al. [12], the heterogeneity in individual tiller yields was caused by the uneven distribution of photo- synthetic active radiation

(PAR), in which the early-emerging superior tillers took priority over the light source at the top and shaded the later-emerging tillers in low-light situations.

3.1.3 Dry weight/plant

Treatment 9 (RNR-15048 + ZnSo4 - 0.4%), which had the significantly greater plant dry weight (55.47 gm), was the one that was noticed. However, statistically speaking, treatment-8 (RNR-15048 + ZnSo4 -0.3%) was comparable to treatment-9 (RNR-15048 + ZnSo4 -0.4%). Chlorosis, decreased or absent tillering, a slower rate of crop maturity, and increased spikelet sterility are all symptoms of severe zinc deficiency in rice. Zinc insufficiency is linked to calcareous soils in lowland rice-growing regions and is exacerbated by protracted flooding. Slaton et al. [13] observed a positive dry matter production response to Zn treatment. Additionally, the rate of photosynthesis and respiration determines how much dry matter accumulates at a given time, which in turn affects how quickly plants grow in terms of their height, leaf area, tillers/hills, etc. As a result, Kumar [14] also noted that the treatment that had the highest growth also gathered more dry matter.

3.2 Yield

3.2.1 Number of panicles/hill

The significant and higher number of panicles/hill (10.01) were observed in treatment-9 with (RNR-15048 + ZnSo4 - 0.4%), which was significantly superior over rest of the treatments. However, treatment-8 (RNR-15048 + ZnSo4 - 0.3%), was found to be statistically at par with treatment-9 (RNR-15048 + ZnSo4 - 0.4%). An adequate Zn supply may have increased the delivery of other nutrients and spurred the growth of the entire plant, which could explain why there were more panicles/hills. Many earlier researchers, including Veeranagappa [15], have also reported an increase in panicles per square metre. The advantage of hybrid rice may also be that it grows long roots and broad leaves, which allow it to absorb more nutrients and yield more grains. It is appropriate for the local climate, particularly during the grain-filling stage of panicle development. Bhuiyan et al. [16] reported similar outcomes as well.

3.2.2 Number of seeds/pod

The significant and higher number of seeds/pod (1.94) were observed in treatment-9 with (RDP

40 kg/ha+ Sulphur 30 kg/ha), which was significantly superior over rest of the treatments. However, treatment-8 (RDP 40 kg/ha+ Sulphur 20 kg/ha), was found to be statistical at par with treatment-9 (RDP 40 kg/ha+ Sulphur 30 kg/ha). The significant rise in seed/pod production could be attributed to enhanced phosphorus fertilisation, which ensures the availability of other plant nutrients, increasing carbohydrate buildup and their re-mobilization to the plant's reproductive organs, which are the closest sink. Because phosphorus is known to promote flowering and fruiting, the plants may have been stimulated to produce more pods on each plant and more seeds per pod. Shah et al. (2000) reported similar results.

3.2.3 Number of grains/panicle

The significant and higher number of grains/panicles (141.37) were observed in treatment-9 with (RNR-15048 + ZnSo4 - 0.4%), which was significantly superior over rest of the treatments. However, treatment-8 (RNR-15048 + ZnSo4 - 0.3%), was found to be statistically at par with treatment-9 (RNR-15048 + ZnSo4 - 0.4%). The increase in the number of grains panicle-1 may have been caused by its stimulation of physiological processes, including photosynthesis, the movement and assimilation of photosynthetic material, and the formation of more spikelets during the spikelet initiation process, all of which led to a greater number of grains/panicle. These conclusions concur with those made by Muhammad et al. [17].

3.2.4 Test weight (g)

Treatment 9 with (RNR-15048 + ZnSo4 - 0.4%) had the considerable and higher test weight (22.27 gm), making it significantly better than the other treatments. Nevertheless, it was discovered that treatment-8 (RNR-15048 + ZnSo4 - 0.3%) was statistically equivalent to treatment-9 (RNR-15048 + ZnSo4 - 0.4%). Higher rates of photosynthetic activity and assimilate translocation during the post-flowering phase may have contributed to an increase in the number of grains in panicle-1, improved panicle weight, and higher grain weight per panicle. These outcomes support the conclusions made by Naik and Das [18].

3.2.5 Grain yield (t/ha)

The significant and higher grain yield (6.43 t/ha) was observed in treatment-9 with (RNR-15048 + ZnSo4 - 0.4%), which was significantly superior

over rest of the treatments. However, treatment-8 (RNR-15048 + ZnSo4 - 0.3%), was found to be statistically at par with treatment-9 (RNR-15048 + ZnSo4 - 0.4%). The greatly increased grain output could be attributed to zinc. Zn treatments may have contributed to the enhanced grain output of rice by starting the early emergence of panicles, which may have allowed for more assimilate storage in rice grains. Khan and others [19]. Additionally, the extended yield characteristics are likely the result of an extended boom and improvement parameters, which ultimately led to an extended grain. These outcomes are in line with Yatheesh Kumar et al. [20].

3.2.6 Straw yield (t/ha)

The significant and higher stover yield (12.11 t/ha) was observed in treatment-9 with (RNR-

15048 + ZnSo4 - 0.4%), which was significantly superior over rest of the treatments. However, treatment-8 (RNR-15048 + ZnSo4 - 0.3%), was found to be statistically at par with treatment-9 (RNR-15048 + ZnSo4 - 0.4%). Zn application increased vegetative development, which was reflected in greater accumulation of dry matter and grain yield per pot. This may have occurred as a result of Zn's involvement in metabolic, hormonal, and photosynthetic processes. Higher straw yield was obtained as a result of stimulating vegetative growth. Ghatak and others [8]. The research by Padmavathi from 1997 [21] also supports the idea that hybrid rice has the capacity to use more nitrogen through the expression of superior growth brought about by the favourable influence on nutrient uptake and physiological growth increase straw yield [22-24].

Table 1. Influence of foliar application of zinc on growth parameters of different rice varieties

S. No.	Treatment combinations	Plant height	Number of tillers/hill	Plant Dry weight
1.	MTU-7029 + ZnSo4- 0.2%	108.88	14.25	47.58
2.	MTU-7029 + ZnSo4- 0.3%	109.37	14.51	48.48
3.	MTU-7029 + ZnSo4- 0.4%	111.73	15.17	49.55
4.	BPT -5204 + ZnSo4- 0.2%	111.23	15.54	51.30
5.	BPT -5204 + ZnSo4 - 0.3%	113.27	15.55	52.13
6.	BPT -5204 + Znso4 - 0.4%	116.41	16.27	53.88
7.	RNR15048 + ZnSo4- 0.2%	115.99	14.60	54.21
8.	RNR-15048 + ZnSo4- 0.3%	116.91	15.48	54.97
9.	RNR-15048 + ZnSo4 - 0.4%	117.68	16.72	55.47
	F test	S	S	S
	S Em.(±)	0.29	0.25	0.33
	CD (P=0.05)	0.88	0.76	0.99

Table 2. Influence of foliar application of zinc on yield and yield attributes of different rice varieties

S. No.	Treatment combinations	No. of Panicles/hill	No. of Grains/panicle	Test weight	Grain yield (t/ha)	Straw yield (t/ha)
1.	MTU-7029 + ZnSo4- 0.2%	7.55	120.56	19.08	5.06	9.67
2.	MTU-7029 + ZnSo4- 0.3%	8.36	126.20	19.67	5.31	9.90
3.	MTU-7029 + ZnSo4- 0.4%	8.57	131.78	20.02	5.56	10.09
4.	BPT -5204 + ZnSo4- 0.2%	7.72	127.65	19.66	5.30	9.83
5.	BPT -5204 + ZnSo4 - 0.3%	8.67	132.04	20.29	5.76	10.29
6.	BPT -5204 + Znso4 - 0.4%	9.08	135.72	21.06	6.16	11.06
7.	RNR15048 + ZnSo4- 0.2%	7.89	132.00	21.42	5.84	10.19
8.	RNR-15048 + ZnSo4- 0.3%	9.37	136.32	22.03	6.19	11.57
9.	RNR-15048 + ZnSo4 - 0.4%	10.01	141.37	22.27	6.43	12.11
	F test	S	S	S	S	S
	S Em (±)	0.23	1.98	0.21	0.07	0.22
	CD (P=0.05)	0.68	5.92	0.62	0.24	0.67

Table 3. Yield validation using SPSS model

S. No.	Treatments	Predicted yield through SPSS model (t/ha)	Obtained yield in experiment (t/ha)	Percentage increase over predicted yield(%)
1.	MTU-7029 + ZnSo4- 0.2%	2.88	5.06	43.08
2.	MTU-7029 + ZnSo4- 0.3%	2.88	5.31	45.76
3.	MTU-7029 + ZnSo4- 0.4%	2.88	5.56	48.20
4.	BPT -5204 + ZnSo4- 0.2%	2.88	5.30	45.20
5.	BPT -5204 + ZnSo4 - 0.3%	2.88	5.79	50.26
6.	BPT -5204 + Znso4 - 0.4%	2.88	6.16	53.25
7.	RNR15048 + ZnSo4- 0.2%	2.88	5.84	50.68
8.	RNR-15048 + ZnSo4- 0.3%	2.88	6.19	53.47
9.	RNR-15048 + ZnSo4 - 0.4%	2.88	6.43	55.21



3.3 Economic Analysis

3.3.1 Gross returns (INR/ha)

In comparison to other treatments, treatment 9 (RNR-15048 + ZnSo4 - 0.4%) had the highest gross return (1,55,870.00 INR/ha).

3.3.2 Net returns

Net return (1,10,542.00 INR /ha) was found to be highest in treatment-9 (RNR-15048 + ZnSo4 - 0.4%) as compared to other treatments.

3.3.3 Benefit cost ratio

Benefit Cost ratio (2.44) was found to be highest in treatment-9 with (RNR-15048 + ZnSo4 - 0.4%) as compared to other treatments.

4. CONCLUSION

It was determined that the variety RNR-15048 (Treatment-9) performed well after receiving Zinc 0.4% application, which enhanced growth and yield characteristics. With the application of Zinc 0.4% on the variety of RNR-15048 (Treatment-9), maximum grain yield, gross returns, net returns, and benefit cost ratio were also noted. Further studies may be necessary to confirm these results since they are based on only one season.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Yadav P, Rangare NR, Anurag JP, Chaurasia AK. Quantitative analysis of rice (*Oryza sativa* L.) in Prayagraj Agro Climate Zone. Journal of Rice Research. 2004;3(1).

2. Saha B, Saha S, Roy PD, Hazra GC, Das A. Zinc fertilization effects on agromorphological and quality parameters of commonly grown rice SAARC J. Agric. 2013;11:105–120.
3. Sarkar SC, Akter M, Islam MR, Haque M. Md. Performance of five selected hybrid rice-varieties in aman season. Journal of Plant Sciences. 2016;4(2):72-7.
4. Prasad R, Shivay YS, Kumar D. Agronomic bio-fortification of cereal grains with iron and zinc Adv. Agron. 2014;125:55–91.
5. Marschner H. Mineral nutrition of higher plants. Academic Press London UK; 1995.
6. Stein AJ, Nestel P, Meenakshi JV, Qaim M, Sachdev HPS, Bhutta ZA. Plant breeding to control zinc deficiency in India: how cost-effective is biofortification. Public Health Nutr. 2007;10:492–501.
7. Gomez KA, Gomez AA. Three or more factor experiment. In: Statistical Procedure for Agricultural Research 2nd edition. 1976;139 -141.
8. Ghatak R, Jana PK, Sounda G, Ghosh RK, Bandopadhyay P. Responses of transplanted rice to Zn fertilization at farmers' field on red and laterite soils of West Bengal. Journal of Inter Academician. 2005;9(2):231- 234.
9. Haque MD, Elora PMD, Romel B. Identification of potential hybrid rice variety in Bangladesh by evaluating the yield potential. World Journal of Agricultural Sciences. 2015;11(1):13-18.
10. Singh V, Rachana Mithare P, Kumar S, Mishra JP, Singh SN, Tiwari D, Sanodiya LK. Performance of hybrid rice cultivar (*Oryza sativa* L.) on growth and yield attributes under agro-climatic conditions of Prayagraj Uttar Pradesh in aman season of planting. International Journal of Current Microbiology and Applied Sciences. 2019;8(9):2970-2982.
11. Gangloff WJ, Westfall DG, Peterson GA, Mortvedt JJ. Relative availability coefficients of organic and inorganic Zn fertilizers. Journal of Plant Nutrition. 2002;25:259-273.
12. Wang Y, Ren T, Lu J. W, Ming R, Li PF, Saddam H, Cong RH, Li XK. Heterogeneity in rice tillers yield associated with tillers formation and nitrogen fertilizer. Agronomy Journal. 2016;108: 1717–1725.
13. Slaton NA, Gbur EE, Wilson CE, Norman RJ. Rice of response to granular zinc sources varying in water-soluble zinc. Soil Science Society of America Journal. 2005a;69:443-452.
14. Kumar SN. Evaluate the establishment techniques on growth and yield of rice Agricultural Research Communication Centre Agricultural Science Digest. 2016;36 (2):110- 113.
15. Veeranagappa P, Prakasha H. C, Basavaraja M. K. and SaqeebullaM.H. 2010. Effect of zinc enriched compost on yield and nutrient uptake of rice (*Oryza sativa* L.) European Journal of Biological Sciences (EJBS) 3 (1): 23-29.
16. Bhuiyan HS, Zahan A, Khatun H, Iqbal M, Alam F, Manir R. Yield performance of newly developed test crossed hybrid rice variety. International Journal of Agronomy and Agricultural Research. 2014;5(4):48-54.
17. Muhammad A, Niaz A, Maqsood J, Iftikar A. Effect of zinc iron and manganese on growth and yield of rice (*Oryza sativa* L.). Pakistan Journal of Agricultural Sciences. 2002;39(3):177-180.
18. Naik SK, Das DK. Relative performance of chelated zinc and zinc sulphate for lowland rice (*Oryza sativa* L.). Nutrient Cycling in Agroecosystems. 2008;81(3): 219-227.
19. Khan MU, Qasim M, Subban M, Ahmad R, Ali L. Effect of zinc application by different methods on the chemical composition and grain quality of rice. Pakistan Journal of Applied Sciences. 2003;3(7): 530- 536.
20. Yatheesh Kumar K, Vikram Singh, Shrish Kumar Singh, Shruti Grace. George assessment of rice (*Oryza sativa* L.) hybrids on growth and yield under agro-climatic conditions of Prayagraj U. P. International Journal of Plant & Soil Science. 2022;34(18):45-49.
21. Padmavathi P. Studies on relative performance of conventional hybrid rice varieties under various levels of nitrogen plant population and planting patterns. phd thesis Indian agricultural research institute New Delhi; 1997.
22. Biswajit Saha, Parimal Panda, Partha Sarathi Patra, Ranajit Panda, Arindam Kundu, Singha Roy AK, Nabakishor Mahato. Effect of different levels of nitrogen on growth and yield of rice (*Oryza sativa* L.) Cultivars under terai-agro climatic situation. Int. J. Curr. Microbiol. App. Sci 2017;6(7):2408-2418.
23. Fatma YS, Rusmana I, Wahyudi AT, Hamim W. Community structure of total bacteria and methane emission-related

prokaryotes in the rice fields applied with urea and biofertilizer. Asian J. Agric. Biol. 2021;2021:202005273.

24. Imran M, Ali A, Safdar ME. The impact of different levels of nitrogen fertilizer on maize hybrids performance under two different environments; 2021.
